



*THE
MARCH
OF THE
PHYLA!*

Invertebrate Phyla



(Figure 18.4)



Phylum Arthropoda

The most abundant phylum!

- There are > 1 million species of **arthropods** (including crayfish, lobsters, crabs, barnacles, spiders, ticks, and insects, many zooplankton).
 - Arthropods have
 - an **open circulatory system** and
 - an **exoskeleton of chitin**, an external skeleton that protects the animal but must be shed in the process of **molting** to permit growth.
 - The body of most arthropods includes a **head, thorax, and abdomen**, although these segments may be fused.
-


What are the numbers?

2/3 of described species of animals are arthropods. The vast majority of known animal species (> 1 million! spp)


On the criteria of species diversity, distribution, and sheer numbers, **arthropods are the most successful animal phylum!** Found in nearly all habitats of the biosphere



Think **JOINTED EXOSKELETON!**



The Exoskeleton



- made of a tough fibrous material called **chitin**.
- gives the skeleton **strength** and **flexibility**.
- several pairs (5 or more) of **jointed appendages** (movable arms and legs).

Arthropod Diversity



- The body of an arthropod
 - is completely covered by the cuticle, an exoskeleton made of chitin
- When an arthropod grows
 - it molts its exoskeleton in a process called ecdysis
 - ecdysiast = stripper

Arthropod Limbs



Arthropod Molting



Arthropod Flight





Types of metamorphosis in insects

- Insect life cycles often include **metamorphosis**, during which the animal takes on different body forms as it develops from larva to adult.
 - More than 80% of insect species undergo **complete metamorphosis** in which a free-living larva transforms from a pupa into an adult.

Other insect species undergo **incomplete metamorphosis** in which the transition from larva to adult is achieved through multiple molts, but without forming a pupa.

Incomplete metamorphosis (direct development)




Complete metamorphosis (indirect development)

(example: butterflies and beetles); not pillbugs

Phylum Nematoda




Have a pseudocoelom and a complete digestive tract

- **Nematodes** or roundworms (phylum Nematoda) are abundant and diverse, with ~500,000 species. Nematodes have
 - bilateral symmetry,
 - three tissue layers = triploblastic
 - a nonliving **cuticle** covering the body that prevents them from drying out,
 - a pseudocoelom body cavity that functions to distribute nutrients and as a hydroskeleton, and
 - a **complete digestive tract** with a mouth and anus.




Nematodes

- Although about 25,000 species of nematodes have been named, estimates of the total number of species range >500,000.
- Humans host at least 50 species of parasitic nematodes.



Invertebrate Phyla



(Figure 18.4)

Phylum Platyhelminthes (Flatworms): the simplest bilateral animals

- The vast majority of animal species belong to the clade Bilateria, consisting of animals with bilateral symmetry.
- **Flatworms** are the simplest bilaterians.
- Live in marine, freshwater, and damp terrestrial habitats.
- Some are parasitic and others are free-living.




Figure 18.7A




There are three major groups of flatworms

1. **Free-living flatworms** (planarians) have

- heads with light-sensitive eyespots,
- flaps to detect chemicals,
- dense clusters of nerve cells that form a simple brain and a pair of nerve cords that runs the length of the body
- a branched gastrovascular cavity with a single opening.





2. **Flukes** are parasitic flatworms with complex life cycles and suckers to attach to their hosts.




3. Tapeworms

are parasitic inhabitants of the digestive tracts of vertebrates, consist of a ribbon-like body with repeated units, have an anterior scolex armed with hooks and suckers that grasp the host, **have no mouth, and simply absorb nutrients across their body surface.** Units at the end of tapeworms are full of ripe eggs that pass out of host.





Phyla Mollusca



Eye candy



Diverse variations on a common body plan

Molluscs (phylum Mollusca) have

- a **muscular foot** that functions in locomotion,
- a **visceral mass** containing most internal organs,
- a **mantle**, which may secrete a shell that encloses the visceral mass, and
- a **true coelom** and a circulatory system that pumps blood throughout the body.
- Many molluscs feed with a rasping **radula**, used to scrape up food.
- The life cycle of many marine molluscs includes a **ciliated trochophore larval stage**.









Figure 18.9A

Visceral mass



Molluscs: Gastropods

The largest group of molluscs and include the snails and slugs.

- found in fresh water, marine, and terrestrial environments,
- the only molluscs that live on land, using the mantle cavity as a lung, and
- often protected by a single, spiral shell.
- Slugs have lost their mantle and shell and some have long colorful projections that function as gills.



Molluscs: Bivalves



Mussels (each about 6 cm long)

A scallop
(about 10 cm
in diameter)

- Include clams, oysters, mussels, and scallops and
- Have shells divided into two halves that are hinged together.
- Most bivalves are sedentary suspension feeders, attached to the substrate by strong threads.

Phyla Mollusca: Cephalopods



A squid (internal shell)



A chambered nautilus (about 21 cm in diameter)



- Are fast, agile predators,
- Have large brains and sophisticated sense organs, including complex image-focusing eyes,
- Have a shell that is large in a nautilus, small and internal in a squid, or missing in an octopus.
- Squid are fast, streamlined predators that use a muscular siphon for jet propulsion.
- Octopuses live on or near the seafloor, where they are active predators.



(direct development in octopuses)



Squid are the largest invertebrates



Phylum Annelida: The segmented worms



Annelids have

- **segmentation**, the subdivision of the body along its length into a series of repeated parts,
- a **true coelom** that functions as a hydrostatic skeleton,
- **triploblastic** tissue organization
- a **nervous system** that includes a simple brain and ventral nerve cord, and
- a **closed circulatory system** in which blood remains enclosed in vessels throughout the body.

Annelids are found in damp soil, the sea, and most freshwater habitats.





Figure 18.10A



© 2012 Pearson Education, Inc.

3 classes: Oligochaetes, Polychaetes and Leeches

Types of annelids

Oligochaetes (earthworms) ingest soil and extract nutrients, aerating soil and improving its texture.



Polychaetes are the largest group of annelids.

- Each polychaete segment has a pair of fleshy appendages with stiff bristles or chaetae.
- Polychaetes search for prey on the seafloor or live in tubes and filter food particles.




Most **leeches** are free-living carnivores; or blood suckers.

- Blood-sucking leeches use razor-like jaws, secrete an anesthetic and an anticoagulant, and suck up to 10 times their own weight in blood.



Invertebrate Phyla




(Figure 18.4)

Phylum: Cnidaria

- Characterized by radial symmetry
- Two tissue layers—epidermis and digestive lining—with a jelly-filled middle region.
- Cnida (nematocysts) – stinging cells (hallmark of the phyla)

Cnidarians exhibit two kinds of body forms.

- The most sedentary **polyp** body is cylindrical with tentacles projecting from one end and/or
- The more mobile **medusa** form is exemplified by a jellyfish.



Medusa form



**A marine jellyfish
(~6 cm in diameter)**

Polyp form

**A hydra
(~2–25
mm tall)**



**A sea anemone
(~6 cm in diameter)**

Phylum Cnidaria: with stinging cells

Cnidarians are carnivores that use their tentacles to capture prey and to push prey into their mouths.

- The mouth leads to the **gastrovascular cavity**, which functions in digestion and circulation and as a hydrostatic skeleton.
 - **Cnidocytes** are unique stinging cells that capture prey and function in defense.
-






*Anthopleura
elegantissima*



Metridium senile

Many have zooxanthellae



Cnidocytes





- Complex intracellular organelles cnidocytes (specialized cells that produce nematocysts)

PURPOSE

- Food capture
- Protection




Cnidocil - triggered chemically or mechanically; operculum pops open, rapid change in pressure discharges thread (~ 3 msec.)





Prey Mimicry



Ex: Portuguese man-of-war (*Physalia*)

passive feeders; lure prey by wriggling individual specialized tentacles





Fig. 2. (A) *Agalma okeni* nematocyst battery resembling a copepod. The terminal filaments (*tf*) appear similar to the antennae of a copepod. (B) Comparison of a fish larva (top) with a nematocyst battery from *Athorybia rosacea*. Two pigmented spots (*ps*) at the enlarged "head" resemble eyes, and two terminal filaments (*tf*) curl back in the position of pectoral fins; *t*, tentillum; *nb*, nematocyst battery. Scale bars, 1.0 mm.



(Purcell 1980)

Box Jellyfish/Cubomedusae (*Chironex fleckeri*)

- Medusa is cuboidal
- Tentacles hang from corners
- Polyps reduced or absent
- Highly toxic nematocysts
- Can eat large fish



death can occur 3-20 min
after sting

Why so deadly?



Tortuous welts left by stings mark the trunk of a woman (below) and a girl's legs.


Both survived, though they were scarred for life.

Box jellyfish are about 95 percent water—practically invisible in turbid coastal

waters. During the summer, says emergency doctor John L. Holmes, "it is quite unsafe to swim in the ocean in tropical northern Australia"—something travel brochures do not stress. Swimmers are urged to use




PETER FENNER (TOP); SURF LIFE SAVING QUEENSLAND (ABOVE)



Lethal lasso snags a banana prawn and draws the meal mouthward. The victim's carapace was instantly punctured by the jelly's stinger capsules, unleashing a multipronged arsenal of toxins that attack breathing and blood cells.






Figure 18.15




Phylum Porifera (sponges)



- No symmetry
- No tissues
- Spicules
- Very primitive/basal



Sponge Anatomy



Water enters through side pores into a central cavity and then flows out through a larger opening.

The inner layer of flagellated **choanocytes** filters food and engulfs it by phagocytosis







Figure 18.15



Phyla Echinodermata

Echinoderms have spiny skin, an endoskeleton, and a water vascular system for movement


Echinoderms: a diverse group including sea stars, sea cucumbers sand dollars, and sea urchins,

- slow-moving or sessile
- all marine
- radially symmetrical, and
- deuterostomes (along with the chordates).

Echinoderms have

- an **endoskeleton** of hard calcareous plates under a thin skin,
 - a **water vascular system** based on a network of water-filled canals that branch into extensions called tube feet, and
 - the ability to regenerate lost arms.
-

Figure 18.13A



© 2012 Pearson Education, Inc.

© 2012 Pearson Education, Inc.






Features of our own phylum, Chordata

Contains vertebrates and their invertebrate ancestors.

The simplest chordates (tunicates and lancelets) lack a backbone.

Most **chordate** embryos and/or adults possess

- a **dorsal, hollow nerve cord**,
 - a flexible, supportive **notochord**
 - **pharyngeal slits**, and
 - a muscular **post-anal tail**.
-



Lancets

