

kedahensis in Kedah and *S. panhai* in southeastern Thailand; all localities being on the Malay Peninsula. Despite the close proximity of all these localities, the difference indicate that some degree of isolation has occurred in the past.

The highlands of Satun, where *S. panhai* occurs, are separated from the Tenasserim range of mountains, while the highlands where *S. ekavibhathai* is found, runs down the eastern side of southern Thailand, continuing into the Main Range of mountains in Peninsular Malaysia. These highlands appear to be the northernmost border of the genus *Stoliczia*, which they are replaced by *Potamon manii*.

Acknowledgements. Thanks are due to Dr. Michael Türkay (SMF), Dr. R. U. Gooding (CMZ), Prof. Phaibul Naiyanetr (CNHM), Dr. Maya Deb (ZSI), Mrs. C. M. Yang (ZRC), Dr. A. Sasekumar (University of Malaya, Kuala Lumpur), Dr. R.P. Lim (formerly University of Malaya, Kuala Lumpur) and Dr. Wong Tat Meng (USM) for their kind help during this study. Prof. Dr. L.B. Holthuis and Dr. Zdravko Stevcic kindly read an early draft of the manuscript. Most of the photographs were taken by Mr. H.K. Yip (National University of Singapore).

References

- Alcock, A. (1910). *Brachyura I. Fasc. II. The Indian Freshwater Crabs - Potamonidae. Catalogue of the Indian Decapod Crustacea in the collection of the Indian Museum. Calcutta*, pp. 1-135, Pls. 1-14.
- Bott, R. (1966). Potamiden aus Asien (*Potamon* Savigny und *Potamiscus* Alcock) (Crustacea, Decapoda). *Senckenbergiana biol., Frankfurt* 47: 469- 509, Pls. 16-21.
- Bott, R. (1970). Die Süßwasserkrabben von Europa, Asien, Australien und ihre Stammesgeschichte. Eine Revision der Potamoidea und Parathelphusoidea (Crustacea, Decapoda). *Abh. Sencken. Naturf. Ges., Frankfurt* 526: 1-338, Pls. 1-58.
- Kemp, S. (1918). Zoological results of a tour in the Far East. Decapod and Stomatopod Crustacea. *Mem. As. Soc. Bengal* 6: 210-297.
- Lanchester, W.F. (1906). Report on the Crustacea. In: *Fasciculi Malayensis* Part 3: 127-134, 1 Pl.
- Man, J. G. De (1892). Decapoden des Indischen Archipels. *Zool. Ergeb. Reise Niederl. Ost-Ind.* 2: 265-527.
- Man, J. G. De (1902). Die von Herrn Professor Kükenthal in Indischen Archipel gesammelten Dekapoden und Stomatopoden. In: W. Kükenthal, Ergebnisse einer Zoologischen Forschungsreise in den Molukken und Borneo. *Abh. Senckenberg. Naturforsch. Gesell.* 25: 467-929, Pls. 19-27.
- Miers, E.J. (1880). On a collection of Crustacea from the Malayan Region. Part II. Telphusoidea, Catometopa and Oxytomata. *Ann. Mag. nat. Hist.* 5: 304-317, Pl. 14.
- Naiyanetr, P. (1978). The geographic distribution of freshwater crabs in Thailand. *Geog. J., Geog. Assn. Thailand, Bangkok* 3 (3): 24-43.
- Naiyanetr, P. (1980a). The use of freshwater shrimps and crabs as the evidence of geographic relationships of Thailand. *J. Natn. Res. Council Thailand* 12(1): 19-38.
- Naiyanetr, P. (1980b). *Crustacean Fauna of Thailand (Decapoda and Stomatopoda)*. Department of Biology, Fac. Sci., Chulalongkorn Univ. Bangkok, 73 pp. (mimeographed).
- Naiyanetr, P. (1983). Crab of San Ka La Kiri. *Geog. J., Geog. Assn. Thailand, Bangkok* 8(2): 43-46.
- Naiyanetr, P. (1988). Freshwater crabs in Thailand. In: *Book published in memory of the Royal Cremation of Associate Professor Dr. Praphun Chitachummong*, Chulalongkorn University, Phaisalsilpa Press, Bangkok, 15pp., 8 colour plates.
- Ng, P.K.L. (1986). Preliminary descriptions of 17 new freshwater crabs of the genera *Geosesarma*, *Parathelphusa*, *Johora* and *Stoliczia* (Crustacea, Decapoda, Brachyura) from South East Asia. *J. Singapore Natn. Acad. Sci.* 15: 36-44.
- Ng, P.K.L. (1987). A revision of the Malayan freshwater crabs of the genus *Johora* Bott, 1966 stat. nov. (Decapoda: Branchyura: Potamidae). *Malay. Nat. J.* 41: 13-44.

- Ng, P.K.L. (1988). *The Freshwater Crabs of Peninsular Malaysia and Singapore*. Department of Zoology, National University of Singapore, Shinglee Press, Singapore, pp. i-viii, 1-156, 4 colour plates.
- Ng, P.K.L. (1989). The identity of the cavernicolous freshwater crab *Potamon (Thelphusa) bidiense* Lanchester, 1900 (Crustacea: Decapoda: Brachyura: Gecarcinucidae) from Sarawak, Borneo, with description of a new genus. *Raffles Bull. Zool., Singapore* 37(1 & 2) : 63-72.
- Ng, P. K. L. & Tan, L.W.H. (1988). The identities of *Heteropilumnus subinteger* (Lanchester, 1900) and *Heteropilumnus hirsutior* (Lanchester, 1990) stat. nov., with description of a new species, *Heteropilumnus holthuisi* sp. nov. (Crustacea, Decapoda, Brachyura, Pilumnidae). *Crustaceana, Leiden* 54(1) : 13-24.
- Rathbun, M. J. (1905). Les crabes d'eau douce. *Nouv. Arch. Mus. Hist. nat., Paris*, (4)7: 159-323, Pls. 13-22.
- Roux, J. (1934). New freshwater decapod crustaceans from the Malay Peninsula. *Bull. Raffles Mus., Singapore*, 9: 28-34.
- Tweedie, M.W.F. (1936). On the crabs of the family Grapsidae in the collection of the Raffles Museum. *Bull. Raffles Mus., Singapore*. 12: 44-70, Pls. 14, 15.
- Wood Mason, J. (1871). Contribution to Indian carcinology. *J. As. Soc. Bengal* 40(2): 189-207, 449-454, Pls. 11-14, 27.

