

A. C. Fariña · J. Freire · E. González-Gurriarán

## Megabenthic decapod crustacean assemblages on the Galician continental shelf and upper slope (north-west Spain)

Received: 21 May 1996 / Accepted: 9 August 1996

**Abstract** The structure of megabenthic decapod crustacean assemblages on the Galician (north-west Spain) continental shelf (100 to 200 depth) and upper slope (200 to 500 m) was analyzed based on surveys carried out in autumn and spring, from 1980 to 1987. Forty species belonging to 19 families were caught. The portunid crab *Polybius henslowii*, a species with pelagic phases, was the most abundant species, but displayed strong spatial and temporal fluctuations. Other dominant species were the Norway lobster *Nephrops norvegicus*, the portunid *Lio-carcinus depurator*, the galatheids *Munida intermedia* and *M. sarsi*, and the shrimps *Solenocera membranacea*, *Plesionika heterocarpus*, *Pasiphaea sivado* and *Dichelopandalus bonnieri*. Total abundance and biomass (average values excluding *Polybius henslowii* = 255 individuals and 2.06 kg/30 min tow) and species richness and diversity,  $H'$  (6.85 species and  $H' = 1.45$  per tow) displayed a significant positive correlation with depth, and strong interannual fluctuations. The factors determining community organization were depth and, to a lesser extent, spatial structure. There was clear evidence of bathymetric zonation, differentiating between species characteristic of the slope (*D. bonnieri* and *Pasiphaea sivado*), shelf-slope edge (*Macropipus tuberculatus*, *Pontophilus spinosus*, *Munida sarsi*, *S. membranacea*, *Processa* spp.) and shelf (*L. depurator*, *Macropodia tenuirostris*, Paguridae and *Chlorotocus crassicornis*). The spatial zonation was related to changes in oceanography and sediment along the continental margin. *Goneplax*

*rhomboides*, *N. norvegicus*, *C. crassicornis* and *Alpheus glaber* are benthic species which generally exhibit burrowing behaviour, and they were found mainly in the southern area where there are fine sediments due to the outwelling from the Rías Baixas. Different benthopelagic shrimps (*Pontophilus spinosus*, *Plesionika heterocarpus*, *Processa* spp. and *Pasiphaea sivado*) were typical of the zone just north of Fisterra, characterized by a convergence of water masses bringing about an increase in productivity due to upwelling. The benthic anomuran and brachyuran crabs *Munida intermedia*, *M. sarsi*, *L. depurator* and *Macropipus tuberculatus* were characteristic of the northwestern zone between Fisterra and Estaca, where the infauna reaches high biomass despite coarser sediments with a lower concentration of organic material than in the southern area. Lastly, both the Paguridae and *Macropodia tenuirostris* were species typically found in the waters in the northern shelf. Based on interannual changes in assemblage structure, two periods could be distinguished: between 1980 and 1984, when *Polybius henslowii*, *D. bonnieri* and *Pasiphaea sivado* had abundance peaks; and another period from 1985 to 1987 when *L. depurator*, *Munida intermedia*, *M. sarsi* and *Macropipus tuberculatus* increased in abundance.

### Introduction

Decapod crustaceans make up one of the dominant groups of megabenthic invertebrates on the Atlantic continental shelf and slope (Lagardère 1973, 1977a; Haedrich et al. 1975, 1980; Wenner and Boesch 1979; Markle et al. 1988; Basford et al. 1989; Olaso 1990; Macpherson 1991; Bianchi 1992a,b; Setubal 1992; García-Castrillo and Olaso 1995), although the structure and organization of their communities is not as well known as that of demersal fishes (see Fariña et al. 1996). However, recent studies in the north-west Mediterranean Sea have made important data available on the

Communicated by A. Rodríguez, Puerto Real

A.C. Fariña  
 Instituto Español de Oceanografía,  
 Apartado 130, E-15080 A Coruña,  
 Spain

J. Freire (✉) · E. González-Gurriarán  
 Departamento de Biología Animal,  
 Biología Vegetal e Ecoloxía,  
 Universidade da Coruña, Campus da Zapateira s/n,  
 E-15071 A Coruña,  
 Spain

distribution and abundance patterns of the crustacean communities, chiefly in deep waters (Abelló et al. 1988; Cartes and Sardá 1992, 1993; Cartes 1993a; Cartes et al. 1994; Sardá et al. 1994a).


The Galician continental margin (shelf and upper slope, 100 to 500 m depth; north-west Spain) comprises a narrow strip (30 to 60 km) (see Fig. 1) characterized by seasonal upwelling (Fraga 1981; Blanton et al. 1984; McClain et al. 1986) and a high primary productivity that determines the structure of the pelagic and benthic food webs (Tenore et al. 1984; Penas 1986; López-Jamar et al. 1992; Varela 1992; Fariña et al. 1997). This area is subject to an intensive demersal fishery targeting different fish species in addition to the Norway lobster (Fernández et al. 1978; Trujillo et al. 1993; Fariña et al. 1994).

Earlier studies have been carried out on the Galician continental margin which examined the distribution patterns of some decapod species (Alonso-Allende 1980, 1982; González-Gurriarán 1987; González-Gurriarán and Olaso 1987; Freire et al. 1991, 1992; González-Gurriarán et al. 1993; Fariña et al. 1994; Fariña and Pereiro 1995). Only González-Gurriarán and Olaso (1987) described the structure of the decapod assemblages on the Galician continental margin, although its relationship to environmental factors and its temporal changes were not analyzed. Olaso (1990) and García-Castrillo and Olaso (1995) studied the megabenthic invertebrates in the Cantabrian Sea, an area adjacent to the Galician coast. Moreover, there have been a number of studies on the structure of crustacean assemblages in the Galician rías (coastal embayments) (González-Gurriarán 1982, 1986; Romero et al. 1982; Iglesias and González-Gurriarán 1984; González-Gurriarán et al. 1991).

This paper analyzes the megabenthic crustacean assemblages on the Galician continental shelf and upper slope based on fishery research-surveys in autumn and spring between 1980 and 1987. The main purposes of the research were to examine the spatial distribution and interannual variation in the assemblages and to determine how they relate to environmental and oceanographic conditions.

## Materials and methods

Sampling methodology has been described in detail in previous papers (González-Gurriarán and Olaso 1987; Sánchez et al. 1991; Fariña et al. 1994, 1997). For the present study, demersal trawl surveys carried out on the Galician continental shelf (100 to 200 m depth) and upper slope (200 to 500 m) in autumn (1980 to 1986) and in spring (1984, 1986 and 1987) were taken into account. A "baca" type trawl gear with a 20 mm codend cover and an estimated opening of 22.0 m horizontally and 1.9 m vertically was used aboard the R.V. "Coornide de Saavedra". The study area was sampled according to a random stratified design. The 200 m isobath separated depth strata, and three geographic areas were pre-determined: Miño-Fisterra (M-F), Fisterra-Estaca de Bares (F-E), and Estaca de Bares-Ribadeo (E-R) (Fig. 1). The area between 200 and 500 m of the E-R area was not sampled because it has a rocky bottom with a steep slope. A total of 293 tows were carried out in



**Fig. 1** Galician coast (north-west Spain). Geographic areas (M-F Miño-Fisterra; F-E Fisterra-Estaca de Bares; E-R Estaca de Bares-Ribadeo) and depth strata (100 to 200 m and 200 to 500 m) sampled between 1980 and 1987

autumn and 168 in spring. Tows were carried out in daylight (08:00 to 19:00 hrs) at a speed of 3 knots. Up to 1984, each tow lasted 60 min; in the survey of spring 1984, 60 and 30 min tows were carried out, and subsequently tows were 30 min long (data from 60 min tows were transformed to 30 min, and density and biomass were defined as the number of individuals/30 min tow or gram wet wt/30 min tow, respectively).

Decapod crustaceans caught in each tow were identified to species, and the number of specimens and total wet weight were recorded for each species. The different species of the family Paguridae (*Pagurus prideauxi*, *P. alatus*, *P. bernhardus*, *P. excavatus* and *Anapagurus laevis*) were identified on some occasions only to genus or family level, and are therefore grouped as Paguridae in this study.

Density, biomass, species richness and diversity ( $H'$  Shannon-Wiener index in  $\log_2$ ) were compared among years, seasons and geographic areas using different models of analysis of covariance (ANCOVA), where depth was included as a covariate. The por-tunid crab *Polybius henslowii* was not included in the data analyses since the catches of this species showed great spatial and temporal fluctuations (mean abundance for each survey ranged between 0 and 7808 individuals/30 min tow, with tows where as many as  $\approx 2 \times 10^5$  individuals were caught). This species has partially pelagic habits, occupying the entire water column and forming dense swarms (Allen 1968; González-Gurriarán 1987) that cannot be sampled adequately with the gear used.

The structure of the decapod crustacean assemblages and its relationship to different environmental factors was analyzed with different methods of correspondence analysis using the software CANOCO v. 3.12 (Ter Braak 1986, 1987) and following the methodology used with demersal fishes in the same study area (Fariña et al. 1997). Sixteen species which appeared in  $\geq 5\%$  of the tows were selected (see "Code" column in Table 1), and log-

**Table 1** Megabenthic decapod crustaceans caught on continental shelf and upper slope off coast of Galicia. Mean density (individuals/30 min tow, SD in parentheses) and biomass (g wet wt/30 min tow, SD in parentheses) and frequency of appearance are given for each species in autumn (1980 to 1987; from 1980 to 1984, the data based on 60 min tows were transformed to 30 min tows) and in spring (1984, 1986 and 1987) [Code code for the species included in multivariate analyses (appearing in  $\geq 5\%$  of tows) and used in Figs. 8, 9 and 10; -absent]

Species	Family	Code	Autumn (N = 293)			Spring (N = 168)		
			ind./tow	g/tow	Freq. (%)	ind./tow	g/tow	Freq. (%)
<i>Polybius henslowii</i>	Portunidae		698.35 (6085.5)	6092.13 (24922.3)	82.3	32.38 (293.4)	636.31 (5990.8)	56.5
<i>Nephrops norvegicus</i>	Nephropidae	<i>Nep nor</i>	13.44 (29.3)	344.68 (898.6)	62.1	7.36 (21.3)	241.90 (798.0)	49.4
<i>Liocarcinus depurator</i>	Portunidae	<i>Lio dep</i>	30.94 (105.6)	408.96 (1591.4)	59.0	48.88 (112.8)	471.50 (1026.7)	62.5
<i>Macropodia tenuirostris</i>	Majidae	<i>Mac ten</i>	3.45 (13.8)	10.46 (39.8)	52.6	0.65 (1.5)	1.70 (3.6)	26.8
<i>Paguridae</i>	Paguridae	<i>Pagur</i>	22.37 (100.5)	295.23 (1252.1)	58.4	15.01 (45.4)	233.90 (739.3)	61.9
<i>Macropipus tuberculatus</i>	Portunidae	<i>Mac tub</i>	4.21 (10.5)	63.00 (152.0)	46.4	2.82 (8.6)	37.90 (109.9)	36.9
<i>Munida intermedia</i>	Galatheidae	<i>Mun int</i>	64.86 (414.9)	582.25 (3515.9)	47.1	26.54 (122.6)	218.30 (1058.2)	45.8
<i>Pontophilus spinosus</i>	Crangonidae	<i>Pon spi</i>	3.53 (11.6)	12.29 (126.2)	34.1	2.02 (4.8)	3.90 (8.0)	36.9
<i>Munida sarsi</i>	Galatheidae	<i>Mun sar</i>	34.14 (164.9)	275.62 (1284.5)	33.8	36.09 (122.1)	225.86 (788.3)	45.2
<i>Plesionika heterocarpus</i>	Pandalidae	<i>Ple het</i>	33.08 (206.1)	68.00 (325.9)	35.5	22.23 (88.4)	74.10 (326.4)	41.1
<i>Solenocera membranacea</i>	Solenoceridae	<i>Sol mem</i>	20.12 (72.7)	54.94 (198.8)	39.2	31.44 (78.4)	80.90 (189.5)	44.6
<i>Processia</i> spp.	Processidae	<i>Pro spp.</i>	3.08 (11.3)	7.71 (30.8)	28.0	3.34 (9.1)	7.70 (18.9)	32.7
<i>Chlorotocus crassicornis</i>	Pandalidae	<i>Chl cra</i>	1.94 (9.3)	4.14 (15.3)	22.5	5.71 (12.2)	13.30 (25.3)	47.6
<i>Goneplax rhomboides</i>	Goneplacidae	<i>Gon rom</i>	0.54 (2.0)	3.26 (14.5)	22.5	0.19 (0.5)	1.66 (5.2)	13.7
<i>Alpheus glaber</i>	Alpheidae	<i>Alp gla</i>	0.77 (6.2)	1.96 (15.2)	11.6	1.36 (5.8)	3.41 (15.5)	20.2
<i>Dichelopandalus bonnieri</i>	Pandalidae	<i>Dic bon</i>	7.68 (66.5)	16.12 (144.0)	6.5	13.07 (71.6)	30.64 (151.9)	15.5
<i>Pasiphaea sivado</i>	Pasiphaeidae	<i>Pas siv</i>	4.73 (36.1)	7.42 (55.7)	6.5	42.81 (212.4)	62.13 (310.8)	12.5
<i>Pontophilus</i> sp.	Crangonidae		-	-	-	0.15 (0.5)	0.72 (2.5)	10.1
<i>Galathea</i> sp.	Galatheidae		2.23 (37.2)	30.60 (511.8)	2.4	0.06 (0.3)	0.24 (1.0)	5.4
<i>Plesionika maritima</i>	Pandalidae		0.02 (0.4)	0.12 (1.8)	0.7	0.26 (2.0)	0.68 (5.0)	3.6
<i>Cancer pagurus</i>	Canceridae		0.03 (0.2)	27.56 (185.3)	2.7	-	-	-
<i>Parapeneus longirrostris</i>	Peneidae		0.01 (0.1)	0.25 (2.7)	1.4	0.15 (1.9)	1.33 (17.0)	1.2
<i>Polycheles typhlops</i>	Polychelidae		0.43 (7.2)	9.18 (153.7)	1.0	0.02 (0.1)	0.45 (3.5)	1.8
<i>Bathynectes maravigna</i>	Portunidae		0.02 (0.2)	0.48 (6.3)	0.7	0.85 (10.8)	0.18 (1.7)	1.8
<i>Monodaeus couchii</i>	Xanthidae		0.002 (0.0)	0.017 (0.3)	0.3	0.018 (0.1)	0.298 (2.3)	1.8
<i>Inachus leptochirus</i>	Majidae		0.03 (0.5)	0.56 (8.5)	1.0	-	-	-
<i>Cancer bellianus</i>	Canceridae		-	-	-	0.02 (0.1)	30.36 (227.3)	1.8
<i>Pasiphaea multidentata</i>	Pasiphaeidae		-	-	-	0.30 (2.8)	1.51 (13.7)	1.8
<i>Homarus gammarus</i>	Nephropidae		-	-	-	0.01 (0.1)	3.21 (30.7)	1.2
<i>Atecyclus rotundatus</i>	Atecyclidae		-	-	-	0.04 (0.5)	0.95 (10.9)	1.2
<i>Inachus dorsettensis</i>	Majidae		-	-	-	0.02 (0.2)	0.03 (0.3)	1.2
<i>Paromola cuvieri</i>	Homolidae		-	-	-	0.01 (0.1)	3.57 (46.4)	0.6
<i>Liocarcinus holsatus</i>	Portunidae		-	-	-	0.01 (0.1)	0.09 (1.2)	0.6
<i>Palaemon</i> sp.	Palaemonidae		-	-	-	0.01 (0.1)	0.06 (0.8)	0.6
<i>Liocarcinus vernalis</i>	Portunidae		-	-	-	0.01 (0.1)	0.03 (0.4)	0.6
<i>Xantho</i> sp.	Xanthidae		0.003 (0.1)	0.010 (0.2)	0.3	-	-	-

transformed density-data for each species per tow were used. The following environmental variables were included: year, season (spring and autumn as categorical variables), depth, and geographic coordinates (surface trend defined by latitude, longitude and the interaction latitude  $\times$  longitude). Correspondence analysis (CA) was used to determine the patterns of variability of the assemblages; subsequently, a canonical correspondence analysis (CCA), whereby the extracted axes constituted linear combinations of the environmental variables, was carried out. Successive partial CCAs were carried out (Ter Braak 1988; Borcard et al. 1992), whereby an environmental variable was analyzed in each case, with the others being introduced as covariates.

## Results

### Species composition

Forty species of decapod crustaceans were caught; 31 in the seven surveys carried out in autumn and 37 during three spring surveys (Table 1). The species belonged to 19 families, the most diverse and abundant of which were Portunidae (6 species), Paguridae (5) and Pandallidae (4). Species belonging to the families Nephropidae, Galatheididae and Solenoceridae were also noteworthy in terms of density or biomass.

*Polybius henslowii* was the dominant species in autumn, making up 74% of the density and 73% of the biomass of the specimens caught. In spring, however, catches of this species were smaller (11.0 and 26.6%, respectively). With the exception of *P. henslowii*, the

dominant species in terms of biomass in autumn (comprising  $> 5\%$  of the catches) were *Munida intermedia*, *Liocarcinus depurator*, *M. sarsi*, *Nephrops norvegicus* and Paguridae; along with *Plesionika heterocarpus* and *Solenocera membranacea* these represented  $> 5\%$  of the catches in number. In autumn, these species combined made up 87% of the density and 91% of the biomass, excluding *Polybius henslowii*. In spring, each of the above-mentioned species constituted  $> 5\%$  of density (except *N. norvegicus*) and biomass (except *Plesionika heterocarpus* and *S. membranacea*). *Dichelopandalus bonnieri* and *Pasiphaea sivado* also made up  $> 5\%$  of the catches in number of individuals. The dominant species in spring corresponded to a total of 93% in density and 92% in biomass.

### Abundance and biomass

The density and biomass of the decapods increased significantly as a function of increasing depth (Figs. 2 and 3); depth was the most important factor explaining the variability in the catches, particularly in spring (in terms of percentage of explained variance in ANCOVA, Table 2). Total density and biomass (excluding *Polybius henslowii*) were very similar in the autumn (mean  $\pm$  SD:  $251 \pm 573$  individuals and  $2.23 \pm 4.98$  kg/30 min tow) and spring ( $261 \pm 401$  individuals and  $1.75 \pm 2.41$  kg/tow) surveys (Fig. 3, Table 3,  $P > 0.1$ ). Geographic





Fig. 2 Density (individuals/30 min tow), biomass (kg wet wt/30 min tow), species richness and diversity ( $H'$  index) of decapod crustaceans of the Galician coast as a function of mean depth of each tow. Correlation coefficient ( $r$ ,  $P < 0.001$ ) and linear regression is shown in all cases

**Fig. 3** Density (individuals/30 min tow) and biomass (kg wet wt/30 min tow) of decapod crustaceans as a function of geographic area and depth strata. Mean values  $\pm$  SE are shown for autumn and spring surveys (Abbreviations as in Fig. 1)



variability was of little importance in the catches. Biomass in F-E was higher than in M-F (more than twice as much at > 200 m depth), even though the differences only reached significant values in autumn, due to the inclusion of the 1980 and 1981 surveys ( $P < 0.05$ ). Density followed the same pattern, especially in the slope area, although no significant differences were found between areas.

Interannual fluctuations in decapod density and biomass were marked (Fig. 4, Table 2,  $P < 0.001$  in all analyses), and constituted the most important factor in the autumn surveys, whereas in spring, depth accounted for a greater percentage of variance. Between 1980 and 1983, when the surveys were made only in autumn, catches were low (with minimum values for the 100 to 200 m stratum of 27 individuals and 0.58 kg/tow in 1980 in M-F and 18 individuals and 0.26 kg/tow in 1983 in F-E). In subsequent years catches increased; however, the trend varied depending on geographic area and depth (significant interaction,  $P < 0.05$ ). In Area M-F, maximum density was attained in the autumn of 1984 (684 individuals/tow at > 200 m) and maximum biomass in the autumn of 1986 (3.52 kg/tow). In the F-E area, maximum values were attained at > 200 m in the autumn of 1986 (1470 individuals and 12.3 kg/tow). In the northern area, E-R, catches reached 387 individuals and 4.26 kg/tow in the spring survey of 1987.

#### Species richness and diversity

The mean number of species caught per tow was  $6.9 \pm 3.3$  and diversity ( $H'$  index) was  $1.45 \pm 0.73$ , with maximum values of 16 and 2.98, respectively. Species richness and diversity increased with increasing depth (Fig. 2) and no significant differences were detected between the autumn and spring surveys (Fig. 5, Table 3;  $P > 0.1$ ). Geographic differences in both parameters were of minor quantitative importance and were largely due to increases in the F-E area.

Interannual differences in species richness and diversity in the autumn surveys were mainly due to the low values recorded in 1980 (mean values of tows carried out at <200 m: 1.9 species/tow and  $H' = 0.28$  in Area M-F

and 3.0 species/tow and  $H' = 0.46$  in Area F-E) (Fig. 6, Table 2,  $P < 0.001$  for the 1980 to 1986 series,  $P > 0.1$  for the 1982 to 1986 series). There were significant differences among the different spring surveys; however, these depended on geographic area and depth stratum. Maximum values were consistently obtained in the F-E area in the autumn cruises (<200 m: 9.3 species/tow in 1986 and  $H' = 1.83$  in 1985; > 200 m: 11.9 species/tow in 1986 and  $H' = 2.10$  in 1981).

#### Spatial and temporal changes of the dominant species

The density of the dominant crustacean species underwent major spatial and temporal changes on the Galician coast (Fig. 7). *Polybius henslowii* appeared almost exclusively from Miño to Estaca, with very high densities throughout the entire depth range in the 1980 to 1983 period, and particularly in 1980 and 1981. Most species had wide bathymetric ranges, although the Paguridae appeared almost exclusively at <200 m, *Macropipus tuberculatus*, *Munida intermedia*, *M. sarsi* and *Solenocera membranacea* were more abundant on the upper slope, and *Dichelopandalus bonnieri* and *Pasiphaea sivado* were only caught on the slope. In the northern zone, Area E-R, catches of *Nephrops norvegicus*, *M. intermedia*, *M. sarsi*, *Plesionika heterocarpus* and *S. membranacea* were rare. (*D. bonnieri* and *Pasiphaea sivado* were not caught in this area as tows were not carried out on the slope.) *Liocarcinus depurator*, *M. intermedia*, *M. sarsi* and *P. sivado* were present in greater abundance in Area F-E, whereas *Plesionika heterocarpus* and *D. bonnieri* were more abundant in the south, in Area M-F.

The only important seasonal differences detected were in the abundance of *Macropipus tuberculatus*, *Munida intermedia* and *M. sarsi*, whose densities were greater in autumn than in spring. In contrast, however, major interannual differences in density affected several species. One group of species increased in density over the years under study (*Liocarcinus depurator*, *Macropipus tuberculatus* between Miño and Estaca, and *Munida intermedia* and *M. sarsi* between Fisterra and Estaca). Another group which underwent strong fluctuations but displayed no clear trend was made up of Paguridae,

**Table 2** Results of analyses of covariance comparing density ( $\log_{10}$  individuals/30 min tow), biomass ( $\log_{10}$  g wet wt/30 min tow), species richness and diversity (index  $H'$ ) of decapod crustaceans between years and geographic areas, with depth as covariate (% *expl* percent explained variance; *area abbreviations* as in Fig. 1)

Source of variation	Cruises G80-G86 (Areas M-F, F-E)			Cruises G82-G86 (Areas M-F, F-E, E-R)			Cruises W84, W86, W87 (Areas M-F, F-E, E-R)		
	( <i>df</i> )	<i>F</i> ( <i>P</i> )	% <i>expl</i>	( <i>df</i> )	<i>F</i> ( <i>P</i> )	% <i>expl</i>	( <i>df</i> )	<i>F</i> ( <i>P</i> )	% <i>expl</i>
Density									
depth	(1)	11.13 (0.001)	4.6	(1)	17.02 (0.000)	7.4	(1)	58.11 (0.000)	37.0
year	(6)	11.02 (0.000)	27.1	(4)	5.93 (0.000)	10.4	(2)	9.95 (0.000)	6.3
area	(1)	1.06 (0.304)	0.4	(2)	1.18 (0.308)	1.0	(2)	1.73 (0.181)	1.1
year $\times$ area	(6)	3.12 (0.006)	7.7	(8)	3.64 (0.001)	12.7	(4)	9.61 (0.000)	6.1
error	(244)			(229)			(157)		
Biomass									
depth	(1)	1.55 (0.215)	0.6	(1)	4.78 (0.030)	2.1	(1)	22.37 (0.000)	14.3
year	(6)	7.26 (0.000)	29.5	(4)	4.42 (0.002)	7.7	(2)	18.90 (0.000)	3.6
area	(1)	4.81 (0.029)	3.3	(2)	1.56 (0.213)	1.4	(2)	2.53 (0.083)	3.2
year $\times$ area	(6)	2.93 (0.006)	11.9	(8)	3.36 (0.001)	11.7	(4)	6.13 (0.000)	15.6
error	(244)			(229)			(157)		
Species richness									
depth	(1)	19.40 (0.000)	8.0	(1)	20.20 (0.000)	8.8	(1)	19.29 (0.000)	12.3
year	(6)	13.08 (0.000)	35.8	(4)	1.95 (0.103)	3.4	(2)	3.99 (0.020)	5.1
area	(1)	4.86 (0.028)	2.2	(2)	9.18 (0.000)	8.0	(2)	1.00 (0.370)	1.3
year $\times$ area	(6)	2.87 (0.010)	7.9	(8)	4.12 (0.000)	14.4	(4)	2.93 (0.023)	7.5
error	(244)			(229)			(157)		
Diversity ( $H'$ )									
depth	(1)	3.12 (0.079)	1.3	(1)	2.85 (0.093)	1.3	(1)	0.35 (0.554)	0.2
year	(6)	5.93 (0.000)	15.1	(4)	1.03 (0.395)	1.8	(2)	3.50 (0.033)	4.2
area	(1)	2.11 (0.148)	0.9	(2)	1.31 (0.273)	1.2	(2)	2.81 (0.063)	3.4
year $\times$ area	(6)	0.38 (0.892)	1.0	(8)	1.01 (0.427)	3.6	(4)	1.42 (0.231)	3.4
error	(235)			(223)			(155)		

**Table 3** Results of analyses of covariance comparing density ( $\log_{10}$  individuals/30 min tow), biomass ( $\log_{10}$  g wet wt/30 min tow), species richness and diversity (index  $H'$ ) of decapod crustaceans between seasons and geographic areas, with depth as covariate (% *expl* percent explained variance; *area abbreviations* as in Fig. 1)


Source of variation	Areas M-F, F-E, E-R (seasons autumn / spring)		
	( <i>df</i> )	<i>F</i> ( <i>P</i> )	% <i>expl</i>
Density			
depth	(1)	54.64 (0.000)	12.1
area	(2)	4.58 (0.033)	1.0
season	(1)	0.95 (0.389)	0.4
area $\times$ season	(2)	1.05 (0.352)	0.5
error	(453)		
Biomass			
depth	(1)	17.12 (0.000)	3.8
area	(2)	2.65 (0.072)	1.2
season	(1)	1.08 (0.299)	0.2
area $\times$ season	(2)	0.66 (0.519)	0.3
error	(453)		
Species richness			
depth	(1)	47.06 (0.000)	10.4
area	(2)	7.25 (0.001)	3.2
season	(1)	1.67 (0.197)	0.4
area $\times$ season	(2)	0.73 (0.482)	0.3
error	(453)		
Diversity ( $H'$ )			
depth	(1)	6.83 (0.009)	1.6
area	(2)	3.54 (0.030)	1.6
season	(1)	0.04 (0.847)	0.0
area $\times$ season	(2)	0.92 (0.400)	0.4
error	(439)		

*Plesionika heterocarpus*, *Dichelopnadalus bonnieri* and *Pasiphaea sivado*. *Polybius henslowii* was the only species that displayed a negative density trend.

#### Assemblage structure and faunal zonation

Indirect gradient analysis (CA) indicated that depth was the factor determining the assemblage structure (Table 4, Fig. 8), showing a high correlation with the first axis extracted ( $r = +0.51$ ,  $P < 0.001$ ). Although the second axis also correlated with depth ( $r = -0.22$ ,  $P < 0.001$ ), for the most part it reflected geographic changes, differentiating Area E-R tows from tows in the other areas. We observed high interannual variability within each geographic area and depth stratum, although no clear trend could be determined. There were no major seasonal differences, as indicated by the fact that the centroids of the autumn and spring surveys corresponding to the same years are located close together on the plane formed by Axes 1 and 2 (Fig. 8b). The results of the direct gradient analysis (CCA) were similar, with Axis 1 (which represents 41% of the species–environment correlation) being linked to depth, while the interannual changes, correlated mostly with Axis 2, were more important than the geographic ones which were associated with Axis 3 (32% compared to 15% of the species–environment correlation).


**Fig. 4** Time-series of density (individuals/30 min tow) and biomass (kg wet wt/30 min tow) of decapod crustaceans for each geographic area and depth strata. Mean values  $\pm$  SE are shown for autumn and spring surveys in each year




Partial CCAs were used to obtain the ordination of the species along the gradients defined by a sole environmental variable, previously eliminating the effect of other variables (Table 4, Fig. 9). All factors tested had a significant effect on assemblage structure (Monte-Carlo randomization tests with 99 permutations,  $P = 0.01$ ).

However, the magnitude of the canonical eigenvalues indicate that the importance of depth, year and geographic location was similar and greater than the effect of season. All species had wide temporal ranges, but the outstanding periods were from 1980 to 1984 (associated with *Macropodia tenuirostris*, *Pontophilus spinosus*, *Di-*

**Fig. 5** Species richness and diversity ( $H'$ ) of decapod crustaceans as a function of geographic area and depth strata. Mean values per tow  $\pm$  SE are shown for autumn and spring surveys (Abbreviations as in Fig. 1)



**Fig. 6** Time-series of species richness and diversity ( $H'$ ) of decapod crustaceans as a function of geographic area and depth strata. Mean values per tow  $\pm$  SE are shown for autumn and spring surveys in each year



*chelopandalus bonnieri* and *Pasiphaea sivado*) and from 1985 to 1987 (associated with *Liocarcinus depurator*, *Macropipus tuberculatus*, *Munida intermedia*, *M. sarsi* and *Goneplax rhomboides*). Seasonal differences were not important and did not affect the dominant species (*Chlorotocus crassicornis*, *D. bonnieri*, and *Pasiphaea sivado* were more abundant in spring, while *Macropodia tenuirostris*, *Macropipus tuberculatus* and *G. rhomboides* were more abundant in autumn).

The ordination along the bathymetric gradient (Fig. 9) revealed a clear faunal zonation between species typical of the slope (*Dichelopandalus bonnieri* and *Pasiphaea sivado*), outer shelf and upper slope (*Macropipus tuberculatus*, *Pontophilus spinosus*, *Munida sarsi*, *Solenocera membranacea* and *Processa* spp.) and of the shelf (*Liocarcinus depurator*, *Macropodia tenuirostris*, Paguridae and *Chlorotocus crassicornis*). A major geographic zonation is also evident (Fig. 10), with four large areas: the southern zone (associated with *Nephrops norvegicus*,

*C. crassicornis*, *Goneplax rhomboides* and *Alpheus glaber*); the zone to the north of Fisterra, where shrimps reached their greatest abundance with narrow distribution ranges (*Pontophilus spinosus*, *Plesionika heterocarpus*, *Solenocera membranacea*, *Processa* spp. and *Pasiphaea sivado*); the central zone of Area F-E (where the anomuran and brachyuran crabs were mainly found: *Macropipus tuberculatus*, *L. depurator*, *Munida intermedia* and *M. sarsi*); and Area E-R (*Macropodia tenuirostris* and Paguridae were species characteristic of this area, but displayed a broad geographic distribution).

## Discussion

The structure of decapod crustacean assemblages on the continental margins in different geographic areas is







Fig. 7

(continued overleaf)



**Fig. 7** Time-series of density (individuals/30 min tow) of dominant species of decapod crustaceans in each geographic area and depth strata (continuous lines joining data points 100 to 200 m; dotted lines joining data points 200 to 500 m). Mean values  $\pm$  SE are shown for surveys carried out in autumn (lines joining vertical bars 1980 to 1986) and in spring (lines joining data points 1984, 1986 and 1987)

largely determined by spatial differences in environmental and oceanographic conditions, and in particular, by depth, bottom type and characteristics of the water masses (Lagardère 1973, 1977a; Haedrich et al. 1975, 1980; Wenner and Boesch 1979; González-Gurriarán and Olaso 1987; Abelló et al. 1988; Markle et al. 1988;

Basford et al. 1989; Olaso 1990; Macpherson 1991; Bianchi 1992a,b; Setubal 1992; Cartes 1993a; Cartes and Sardá 1993; Cartes et al. 1994; Sardá et al. 1994a; García-Castrillo and Olaso 1995).

Off the Galician coast, there is a marked faunal zonation along the bathymetric gradient that is especially


**Table 4** Results of correspondence analyses (CA) analyzing structure of decapod crustacean assemblage. Data are eigenvalues of first four axes extracted with (*in parentheses*) percentage of variance explained of the species data [*E total* sum of all eigenvalues; *Ec* sum of all canonical eigenvalues (in both cases after fitting covariates)]; correlation species–environment, *r* (*in parentheses* percentage of variance explained of relationship species–environment); and correlation between the different environmental variables and each axis (for canonical CA) [*Y* year; *S* season (autumn/spring); *D* depth; *G* geographical location (latitude, longitude, latitude × longitude)]

Analysis environmental variables	Covariates	Axis:				E total	Ec
		1	2	3	4		
<b>Non-canonical CA</b>							
–	–	0.376 (13.9)	0.347 (12.8)	0.260 (9.4)	0.206 (7.7)	2.705	–
<b>Canonical CA</b>							
Y, S, D, G	–	0.224 (8.1)	0.175 (6.3)	0.080 (2.9)	0.038 (1.3)	2.705	0.549
<i>r</i>		0.813 (40.8)	0.734 (31.9)	0.572 (14.5)	0.462 (7.0)		
<b>Correlation axis–environmental variable</b>							
year							
season (autumn)		–0.280	–0.948	–0.064	–0.114		
depth		–0.056	–0.553	0.496	–0.649		
latitude		0.834	–0.454	0.287	–0.105		
longitude		0.074	–0.328	0.671	0.160		
latitude × longitude		0.289	0.139	–0.676	–0.565		
		0.351	0.076	–0.608	–0.611		
<b>Partial canonical CA</b>							
Y	S, D, G	0.144 (6.1)	0.303 (12.8)	0.230 (9.7)	0.201 (8.5)	2.369	0.144
<i>r</i>		0.705 (100)	–	–	–		
S	Y, D, G	0.058 (2.5)	0.303 (13.3)	0.230 (10.1)	0.201 (8.8)	2.283	0.058
<i>r</i>		0.553 (100)	–	–	–		
D	Y, S, G	0.152 (6.4)	0.303 (12.7)	0.230 (9.7)	0.201 (8.5)	2.377	0.152
<i>r</i>		0.760 (100)	–	–	–		
G	Y, S, D	0.075 (3.2)	0.022 (0.9)	0.014 (0.6)	0.303 (13.0)	2.335	0.110
<i>r</i>		0.523 (67.3)	0.404 (19.8)	0.339 (12.6)	–		

evident around the shelf break ( $\approx 200$  m). This coincides to a great extent with the zonation reported in the north-west Mediterranean (Abelló et al. 1988) and in other areas of the North Atlantic (Lagardère 1973, 1977a; Haedrich et al. 1975, 1980; Markle et al. 1988; Olaso 1990; García-Castrillo and Olaso 1995). The distribution and abundance patterns of the various species on the coast of Galicia coincide with those described in earlier studies (González-Gurriarán 1987; González-Gurriarán and Olaso 1987; Freire et al. 1991, 1992; González-Gurriarán et al. 1993; Fariña et al. 1994; Fariña and Pereiro 1995) and with other studies done in the north-east Atlantic (Lagardère 1973, 1977a; Olaso 1990; García-Castrillo and Olaso 1995). Our results point to an increase in abundance and diversity of decapod crustaceans with depth. However, other studies report that on the middle and lower slope, these parameters decrease again (Haedrich et al. 1975, 1980), with the exception of

the submarine canyons located on the slope where biomass and density undergo a substantial increase (Sardá et al. 1994a).


In addition to being associated with depth, the spatial structure of the shelf and upper slope crustacean assemblages is related to sediment granulometry that particularly affects benthic species, and to water masses and circulation patterns that affect benthopelagic species (Lagardère 1973, 1977a; Abelló et al. 1988; Basford et al. 1989; Macpherson 1991; Bianchi 1992a,b; Setubal 1992). In deep zones where physical factors are more stable, food availability plays an essential role in zonation (Lagardère 1977a,b; Cartes and Sardá 1993). In our study area, the faunal zonation may be attributable more to oceanographic and sediment changes than to biogeographic factors, since the area is relatively small and the faunal composition resembles other zones located to the north and even in the Mediterranean (La-



**Fig. 8** Correspondence analysis of surveys carried out in autumn (1980 to 1986) and spring (1984, 1986 and 1987). **a** Scores of species of decapod crustaceans on Axes 1 and 2 (*species codes* as in Table 1); **b** centroids of tows for each survey, season (*continuous line* autumn, *dotted line* spring), geographic area, and depth stratum (for simplicity, temporal evolution on plane formed by Axes 1 and 2 of each area and stratum are presented on separate graphs)

gardère 1973; Abelló et al. 1988; Olaso 1990; Cartes 1993a). The faunal zonation observed in this study can be related to the spatial heterogeneity of the physical characteristics of the study area. The species that inhabit the south of the Galician continental margin are benthic, and in many cases exhibit burrowing behavior, depending on the existence of fine sediment (Lagardère

1973, 1977a), which is common in this area due to the outwelling from the Rías Baixas (López-Jamar et al. 1992). The zone directly to the north of Fisterra, where benthopelagic shrimps are abundant, is characterized by a convergence of water masses and upwelling processes with high productivity (Fraga 1981; McClain et al. 1986). Benthic crabs are typical of the area between



**Fig. 9** Ordination of species of decapod crustaceans obtained from partial canonical correspondence analyses analyzing effect of year, season and depth on assemblage structure. Score  $\pm$  tolerance (root mean-square deviation) of each species and centroid of each survey (a) and season (b), and ordination of the bathymetric gradient (obtained by linear regression of depth in respect to score of the tow) (c) along the first axis are given (*Species codes* as in Table 1)


Fisterra and Estaca, although they have a broad spatial distribution. This area has a high level of infaunal biomass as a result of the upwelling processes, even though the sediment is coarser and the organic content lower than to the south of the shelf (López-Jamar and González 1987; López-Jamar et al. 1992).

Despite the fact that about half the species on the continental shelf and slope of Galicia have also been reported in the rías (between 5 and 50 m depth) (González-Gurriarán and Méndez 1985; González-Gurriarán and Olaso 1987), the community structure in the two zones is completely different, so that only *Liocarcinus depurator* and occasionally *Polybius henslowii* are dominant species common to both zones (González-Gurriarán 1982, 1986; Romero et al. 1982; Iglesias and González-Gurriarán 1984; González-Gurriarán and Olaso 1987; González-Gurriarán et al. 1991).

The spatial organization of megabenthic crustacean assemblages and of demersal fish assemblages shows very different patterns in our study area as well as in other zones. The spatial and temporal changes in the demersal fishes of the Galician coast are mainly associated with oceanographic phenomena displaying great seasonal and interannual variability (Fariña et al. 1997), whereas in decapods the more stable sediment and oceanographic conditions are of greater importance. In the case of fishes, which are more mobile and in many

cases carry out ontogenetic migrations through the bathymetric gradient, faunal zonation and spatial structure of the assemblage is of little importance (Markle et al. 1988; Haedrich and Merrett 1990; Macpherson and Duarte 1991; Fariña et al. 1997). Migrations of decapod crustaceans are less important, although the benthopelagic species carry out seasonal and diel vertical movements along the bathymetric gradient (Lagardère 1977a,b; Cartes et al. 1993; Sardá et al. 1994b). On the other hand, some of the dominant species of demersal fishes on the Galician shelf make use of the rías either seasonally or during some life-history stages (Fariña et al. 1997), indicating the importance of mobility and migrations in assemblage structure. In crustaceans, as mentioned earlier, there is very little interchange between the assemblages in the rías and those on the shelf.

The abundance and diversity of fishes on the Galician continental margin decreased with depth, displaying opposite patterns to crustaceans (Fariña et al. 1997; and present paper); this agrees with the inverse relationship between fishes and invertebrates proposed by Haedrich et al. (1975, 1980). Similarly, in the Mediterranean, the dominant species of demersal fishes in the shelf are replaced by benthopelagic crustaceans on the upper slope (Sardá et al. 1994a). These distribution patterns may be caused in part by the trophic interactions between the



**Fig. 10** Geographic distribution of scores of tows and species obtained from partial canonical correspondence analysis analyzing spatial structure of assemblage of decapod crustaceans (variables = latitude, longitude, latitude  $\times$  longitude; covariates = year, season and depth). Isolines of tow scores on first axis are shown (mapping based on regression: score =  $b_1 + b_2 \times \text{latitude} + b_3 \times \text{longitude} + b_4 \times \text{latitude} \times \text{longitude}$ ), together with score  $\pm$  tolerance (root mean-square deviation) of each species (*Species codes* as in Table 1)

two groups. In fact the juvenile stages and small-sized species of demersal fishes, which prey mostly on invertebrates (Macpherson 1981; González et al. 1985; Olaso 1990; Olaso and Rodríguez-Marín 1995), are more abundant in shallow waters (Macpherson and Duarte 1991; Fariña et al. 1997).

Decapod crustaceans play an important ecological role within the megabenthic assemblages on the continental margins, owing to their abundance and trophic relationships. Species such as *Nephrops norvegicus*, *Liocarcinus depurator*, *Macropipus tuberculatus*, *Solenocera membranacea* or Paguridae prey on epibenthic and infaunal organisms (Thomas and Davidson 1962; Lagardère 1977a,b; Abelló and Cartes 1987; Froggia and Gramitto 1987; Abelló 1989; Baden et al. 1990; Freire 1996). Various benthopelagic shrimps such as *Dichelopandalus bonnierii*, *Pasiphaea* spp., *Plesionika* spp., and others prey on benthic, suprabenthic and mesopelagic organisms (Lagardère 1977a,b; Cartes 1991, 1993b,c). *Polybius henslowii* also has a diet composed of benthic

organisms and pelagic fishes and cephalopods (E. Poza and E. González-Gurriarán, unpublished data). It has been suggested that megabenthic decapods have an important scavenging role, feeding particularly on fishes. On the other hand, several decapod species constitute a large part of the diet of demersal fishes that are dominant on the coasts of Galicia (Macpherson 1981; González et al. 1985; Olaso 1990; Olaso and Rodríguez-Marín 1995). These data along with the results of this study suggest that the dominant decapod species on the Galician continental margin constitute an intermediate trophic level, and are a key factor in energy flow to higher levels of the food web and, in the case of the benthopelagic species, in the process of benthopelagic coupling.

**Acknowledgements** We would like to thank all those involved in the demersal trawl surveys on the Galician coast carried out by the Instituto Español de Oceanografía between 1980 and 1987 for their valuable help. C.P. Teed assisted in the preparation of the English version of the manuscript.

## References

- Abelló P (1989) Feeding habits of *Macropipus tuberculatus* (Brachyura, Portunidae) off the Catalan coast (NW Mediterranean). *Miscellània zool* 13: 45–50
- Abelló P, Cartes J (1987) Observaciones sobre la alimentación de *Liocarcinus depurator* (L.) (Brachyura: Portunidae) en el Mar Catalán. *Investigación pesq* 51(Supl 1): 413–419
- Abelló P, Valladares FJ, Castellón A (1988) Analysis of the structure of decapod crustacean assemblages off the Catalan coast (North-West Mediterranean). *Mar Biol* 98: 39–49
- Allen JA (1968) The surface swarming of *Polybius henslowii* [Brachyura: Portunidae]. *J mar biol Ass UK* 48: 107–111
- Alonso-Allende JM (1980) Distribución de la cigala (*Nephrops norvegicus*) en las costas de Galicia. *Investigación pesq* 44: 347–360
- Alonso-Allende JM (1982) Abundancia de la cigala (*Nephrops norvegicus*) (Decapoda: Reptantia) en las costas de Galicia. *Investigación pesq* 46: 63–82
- Baden SP, Pihl L, Rosenberg R (1990) Effects of oxygen depletion on the ecology, blood physiology and fishery of the Norway lobster *Nephrops norvegicus*. *Mar Ecol Prog Ser* 67: 141–155
- Basford DJ, Eleftheriou A, Raffaelli D (1989) The epifauna of the northern North Sea (56°–61°N). *J mar biol Ass UK* 69: 387–407
- Bianchi G (1992a) Demersal assemblages of the continental shelf and upper slope of Angola. *Mar Ecol Prog Ser* 81: 101–120
- Bianchi G (1992b) Study of the demersal assemblages of the continental shelf and upper slope off Congo and Gabon, based on the trawl surveys of the RV 'Dr Fridtjof Nansen'. *Mar Ecol Prog Ser* 85: 9–23
- Blanton JO, Atkinson LP, Fernández de Castillejo F, Lavín A (1984) Coastal upwelling off the Rías Bajas, Galicia, Northwest Spain. I. Hydrographic studies. *Rapp P-v Réun Cons perm int Explor Mer* 183: 79–90
- Borcard D, Legendre P, Dapreau P (1992) Partialling out the spatial component of ecological variation. *Ecology* 73: 1045–1055
- Cartes JE (1991) Análisis de las comunidades y estructura trófica de los crustáceos decápodos batiales del Mar Catalán. Ph D thesis. Universidad Politécnica de Catalunya, Barcelona, Spain
- Cartes JE (1993a) Deep-sea decapod fauna of the Western Mediterranean: bathymetric distribution and biogeographic aspects. *Crustaceana* 65: 29–40

- Cartes JE (1993b) Feeding habits of pasiphaeid shrimps close to the bottom of the Western Mediterranean slope. *Mar Biol* 117: 459–468
- Cartes JE (1993c) Diets of deep-water pandalid shrimps on the Western Mediterranean slope. *Mar Ecol Prog Ser* 96: 49–61
- Cartes JE, Company JB, Maynou F (1994) Deep-water decapod crustacean communities in the Northwestern Mediterranean: influence of submarine canyons and season. *Mar Biol* 120: 221–229
- Cartes JE, Sardá F (1992) Abundance and diversity of decapod crustaceans in the deep-Catalan Sea (Western Mediterranean). *J nat Hist* 26: 1305–1323
- Cartes JE, Sardá F (1993) Zonation of deep-sea decapod fauna in the Catalan Sea (Western Mediterranean). *Mar Ecol Prog Ser* 94: 27–34
- Cartes JE, Sardá F, Company JB, Leonart J (1993) Day–night migrations by deep-sea decapod crustaceans in experimental samplings in the Western Mediterranean Sea. *J exp mar Biol Ecol* 171: 63–73
- Fariña AC, Freire J, González-Gurriarán E (1994) *Nephrops norvegicus* in the Galician continental shelf (NW Spain): abundance and distribution. *Fish Res* 19: 333–347
- Fariña AC, Freire J, González-Gurriarán E (1997) Demersal fish assemblages in the Galician continental shelf and upper slope (NW Spain): spatial structure and long-term changes. *Estuar, cstl Shelf Sci* (in press)
- Fariña AC, Pereiro FJ (1995) Distribution and abundance of molluscs and decapod crustaceans in trawl samples from the Galician Shelf (NW Spain). *ICES mar Sci Symp (Act Symp)* 199: 189–199
- Fernández A, Pereiro FX, Iglesias S, Porteiro C, Pallarés P (1978) La pesquería demersal gallega. Estrategias de pesca para su regulación racional en base a la merluza. *Boln Inst esp Oceanogr* 4: 67–109
- Fraga F (1981) Upwelling off the Galician coast, Northwest Spain. In: Richards FA (ed) *Coastal upwelling*. American Geophysical Union, Washington DC, 176–182
- Freire J (1996) Feeding ecology of *Liocarcinus depurator* (Decapoda: Portunidae) in the Ría de Arousa (Galicia, north-west Spain): effects of habitat, season, and life history. *Mar Biol* 126: 297–311
- Freire J, González-Gurriarán E, Olaso I (1992) Spatial distribution of *Munida intermedia* and *M. sarsi* (Crustacea: Anomura) on the Galician continental shelf (NW Spain): application of geostatistical analysis. *Estuar, cstl Shelf Sci* 35: 637–648
- Freire J, González-Gurriarán E, Olaso I, Fernández L (1991) Análisis geoestadístico de la distribución espacial de *Plesionika heterocarpus* y *Solenocera membranacea* (Crustacea: Decapoda) en la plataforma continental de Galicia (NW España). *Boln Inst esp Oceanogr* 7: 79–88
- Frogliá C, Gramitto ME (1987) Notes on growth and biology of *Solenocera membranacea* (Risso, 1816) in the Central Adriatic Sea (Decapoda: Solenoceridae). *Investigación pesq* 51(Supl 1): 189–199
- García-Castrillo G, Olaso I (1995) Composition and structure of the invertebrate megabenthos on the shelf of the Cantabrian Sea. *ICES mar Sci Symp (Act Symp)* 199: 151–156
- González R, Olaso I, Pereda P (1985) Contribución al conocimiento de la alimentación de la merluza (*Merluccius merluccius* L.) en la plataforma continental de Galicia y del Cantábrico. *Boln Inst esp Oceanogr* 2: 49–60
- González-Gurriarán E (1982) Estudio de la comunidad de crustáceos decápodos (Brachyura) en la Ría de Arousa (Galicia NW España), y su relación con el cultivo de mejillón en batea. *Boln Inst esp Oceanogr* 7: 223–254
- González-Gurriarán E (1986) Seasonal changes of benthic megafauna in the Ría de Muros e Noia (Galicia, North-West Spain). II. Decapod crustaceans (Brachyura). *Mar Biol* 92: 201–210
- González-Gurriarán E (1987) El patexo *Polybius henslowii* Leach (Decapoda–Brachyura), en las costas de Galicia (NW de España): I. Distribución espacial y cambios temporales. *Investigación pesq* 51(Supl 1): 361–374
- González-Gurriarán E, Fernández L, Freire J, Muíño R, Rodríguez Solórzano M (1991) Estructura de la comunidad megabentónica (crustáceos decápodos – Brachyura – y peces demersales) de la Ría de Ferrol (Galicia, NW España). *Boln Inst esp Oceanogr* 7: 89–99
- González-Gurriarán E, Freire J, Fernández L (1993) Geostatistical analysis of spatial distribution of *Liocarcinus depurator*, *Macropipus tuberculatus* and *Polybius henslowii* (Crustacea: Brachyura) over the Galician continental shelf (NW Spain). *Mar Biol* 115: 453–461
- González-Gurriarán E, Méndez M (1985) Crustáceos decápodos das costas de Galicia. I. Brachyura. *Cuad Área Ciencias biol, Semin Estud Galegos* 2: 1–242
- González-Gurriarán E, Olaso I (1987) Cambios espaciales y temporales de los crustáceos decápodos de la plataforma continental de Galicia (NW de España). *Investigación pesq* 51(Supl 1): 323–341
- Haedrich RL, Merrett NR (1990) Little evidence of faunal zonation or communities in deep sea demersal fish fauna. *Prog Oceanogr* 24: 239–250
- Haedrich RL, Rowe GT, Polloni PT (1975) Zonation and faunal composition of epibenthic populations on the continental slope south of New England. *J mar Res* 33: 191–212
- Haedrich RL, Rowe GT, Polloni PT (1980) The megabenthic fauna in the deep sea south of New England, USA. *Mar Biol* 57: 165–179
- Iglesias J, González-Gurriarán E (1984) Primeros datos sobre la megafauna bentónica de la ría de Pontevedra: peces demersales y crustáceos decápodos (Brachyura). *Cuad Área Ciencias mar, Semin Estud Galegos* 1: 303–319
- Lagardère JP (1973) Distribution des décapodes dans le sud du Golfe de Gascogne. *Revue Trav Inst (scient tech) Pêch marit* 37: 77–95
- Lagardère JP (1977a) Recherches sur la distribution verticale et sur l'alimentation des crustacés décapodes benthiques de la pente continentale du Golfe de Gascogne. *Analyse des groupements carcinologiques*. *Bull Cent Étud Rech scient Biarritz* 11: 367–440
- Lagardère JP (1977b) Recherches sur le régime alimentaire et le comportement prédateur des décapodes benthiques de la pente continentale de l'Atlantique Nord ridental (Golfe de Gascogne et Maroc). In: Keegan BF, Ceidigh PO, Boaden PSJ (eds) *Biology of benthic organisms*. Pergamon Press, New York, pp 397–408
- López-Jamar E, Cal RM, González G, Hanson RB, Rey J, Santiago G, Tenore KR (1992) Upwelling and outwelling effects on the benthic regime of the continental shelf off Galicia, NW Spain. *J mar Res* 50: 465–488
- López-Jamar E, González G (1987) Infaunal macrobenthos of the Galician continental shelf off La Coruña Bay, Northwest Spain. *Biol Oceanogr* 4: 165–192
- Macpherson E (1981) Resource partitioning in a Mediterranean demersal fish community. *Mar Ecol Prog Ser* 4: 183–193
- Macpherson E (1991) Biogeography and community structure of the decapod crustacean fauna off Namibia (southeast Atlantic). *J Crustacean Biol* 11: 401–415
- Macpherson E, Duarte CM (1991) Bathymetric trends in demersal fish size: is there a general relationship?. *Mar Ecol Prog Ser* 71: 103–112
- Markle DF, Dadswell MJ, Halliday RG (1988) Demersal fish and decapod crustacean fauna of the upper continental slope off Nova Scotia from LaHavre to St. Pierre Banks. *Can J Zool* 66: 1952–1960
- McClain CR, Chao SY, Atkinson LP, Blanton JO, Fernández de Castillejo F (1986) Wind-driven upwelling in the vicinity of Cape Finisterre, Spain. *J geophys Res* 91: 8470–8486
- Olaso I (1990) Distribución y abundancia del megabentos invertebrado en fondos de la plataforma cantábrica. *Publnes esp Inst españ Oceanogr, Madrid* 5: 1–128
- Olaso I, Rodríguez-Marín E (1995) Decapod crustaceans in the diets of demersal fish in the Cantabrian Sea. *ICES mar Sci Symp (Act Symp)* 199: 209–221
- Penas E (1986) Modelo preliminar del ecosistema de la plataforma continental de Galicia. *Boln Inst esp Oceanogr* 3: 43–56

- Romero P, González-Gurriarán E, Penas E (1982) Influence of mussel rafts on spatial and seasonal abundance of crabs in the Ría de Arousa, North-West Spain. *Mar Biol* 72: 201–210
- Sánchez F, Pereiro FJ, Rodríguez-Marín E (1991) Abundance and distribution of the main commercial fish on the Northern coast of Spain (ICES divisions VIIIc and IXa) from bottom trawl surveys. *Int Counc Explor Sea Comm Meet (Demersal Fish Comm)* 1991/G: 53: 1–30
- Sardá F, Cartes JE, Company JB (1994a) Spatio-temporal variations in megabenthos abundance in three different habitats of the Catalan deep-sea (Western Mediterranean). *Mar Biol* 120: 211–219
- Sardá F, Cartes JE, Norbis W (1994b) Spatio-temporal structure of the deep-water shrimp *Aristeus antennatus* (Decapoda: Aristeidae) population in the western Mediterranean. *Fish Bull US* 92: 599–607
- Setubal AM (1992) Structure and dynamics of benthic megafauna on the continental shelf offshore of Ubatuba, southeastern Brazil. *Mar Ecol Prog Ser* 86: 63–76
- Tenore KR, Cal RM, Hanson RB, López-Jamar E, Santiago G, Tietjen JH (1984) Coastal upwelling off the Rías Bajas, Galicia, Northwest Spain. II. Benthic studies. *Rapp P-v Réun Cons perm int Explor Mer* 183: 91–100
- Ter Braak CJF (1986) Canonical correspondence analysis: a new eigenvector technique for multivariate direct gradient analysis. *Ecology* 67: 1167–1179
- Ter Braak CJF (1987) CANOCO - A FORTRAN program for canonical community ordination by [partial] [detrended] [canonical] correspondence analysis, principal component analysis and redundancy analysis. TNO Institute of Applied Computer Science, Wageningen, The Netherlands
- Ter Braak CJF (1988) Partial canonical correspondence analysis. In: Bock HH (ed) *Classification and related methods of data analysis*. Elsevier, North Holland, pp 551–558
- Thomas HJ, Davidson C (1962) The food of the Norway lobster. *Mar Res* 3: 1–15
- Trujillo V, Meixide M, Porteiro C, Pérez N, Pereiro FJ (1993) Cambios de malla y esfuerzo en las pesquerías multiespecíficas en aguas atlánticas de la Península Ibérica. *Boln Inst esp Oceanogr* 9: 23–40
- Varela M (1992) Upwelling and phytoplankton ecology in Galician (NW Spain) rías and shelf waters. *Boln Inst esp Oceanogr* 8: 57–74
- Wenner EL, Boesch DF (1979) Distribution patterns of epibenthic decapod Crustacea along the shelf-slope coenocline, Middle Atlantic Bight, USA. *Bull biol Soc Wash* 3: 106–133