

- _____, 1983, *Turritella* zonation across the Cretaceous-Tertiary boundary, California: University of California Publications in Geological Sciences, v. 125, 165 p.
- _____, 1986, Pacific Coast Cretaceous molluscan faunas: Time and aspect of changes, in Abbott, P.L., ed., Cretaceous stratigraphy, western North America: Society of Economic Paleontologists and Mineralogists, Pacific Section, Guidebook, p. 131-136.
- _____, 1989, California Late Cretaceous donaciform bivalves: *The Veliger*, v. 32, p. 188-208.
- _____, and Popenoe, W.P., 1962, *Meekia*, Enigmatic Cretaceous pelecypod genus: University of California Publications in Geological Sciences, v. 40, p. 289-344.
- _____, and _____, 1992, Pacific slope Cretaceous bivalves of the genus *Calva*: Natural History Museum of Los Angeles County, Contributions in Science, no. 433, p. 1-68.
- Sohl, N.F., 1971, North American Cretaceous biotic provinces delineated by gastropods: North American Paleontological Convention, September 1969, Proceedings, Part L, p. 1610-1638, 13 figs.
- Verosub, K.L., Haggart, J.W., and Ward, P.D., 1989, Magnetostratigraphy of Upper Cretaceous strata of the Sacramento Valley, California: Geological Society of America Bulletin, v. 101, p. 521-533.
- Walker, R.G., 1978, Deep water sandstone facies and ancient submarine fans -- Models for exploration for stratigraphic traps: American Association of Petroleum Geologists Bulletin, v. 62, p. 932-966.
- _____, 1984, Turbidites and associated coarse clastic deposits, in Walker, R.G., ed., Facies models, second edition: Geoscience Canada, Reprint Series 1, p. 171-188.
- Ward, P.D., 1978, Revisions of the stratigraphy and biochronology of the Upper Cretaceous Nanaimo Group, British Columbia and Washington State: Canadian Journal of Earth Sciences, v. 15, p. 405-423.
- Whidden, K.J., Lund, S.P., Bottjer, D.J., Champion, D.E., and Howell, D.G., 1991, Paleomagnetic evidence from Upper Cretaceous strata for autochthoneity of the central block of Salinia: Geological Society of America, Abstracts with Program, v. 23, p. A109.



View of Pigeon Point from the north with inner-fan channel conglomerate in the foreground.

Plate 1. Mollusks from the Pigeon Point Formation. Fossil localities (Loc.) are shown on Figure 2. All figures natural size unless otherwise indicated.

1. *Inoceramus subundatus* Meek, 1861, left valve, x0.4. USNM 468567. Loc. 5, USGS Mesozoic locality M8748.
2. *Christitys* cf. *C. martini* Popenoe and Saul, 1987. USNM 468568. Loc. 11, USGS Mesozoic locality M8611.
3. *Glycymeris* (*Glycymerita*) *anae* Smith, 1945, left valve. USNM 468569. Loc. 7, USGS Mesozoic locality M8599.
4. *Glycymeris* (*Glycymerita*) *apletos* Dailey and Popenoe, 1966, left valve. USNM 468570. Loc. 8, USGS Mesozoic locality M8601.
5. *Didymoceras* sp. Broken portion of retroversal hook. UCMP 398603. Loc. 2, UCB locality A7601.
6. *Inoceramus* (*Endocostea*) *balticus* Böhm, 1907, right valve, x0.8. UCMP 398604. Loc. 2, UCB locality B6358.
- 7-8. *Indogrammatodon?* cf. *I. vancouverensis* (Meek, 1858). 7) Left valve. USNM 468571. 8) Right valve. USNM 468572. Loc. 12, USGS Mesozoic locality 9593.
- 9-10. *Ampullina packardi* Popenoe, 1937. USNM 468573. Loc. 10, USGS Mesozoic locality M8825.
11. *Meekia* (*Mygallia*) *bella* Saul and Popenoe, 1962, right valve. USNM 468574. Loc. 7, USGS Mesozoic locality M8600.
- 12-13. *Willimaetra* (*Petromactra*) *truncata* (Gabb, 1864), right valve. LACMIP 12125. Loc. 4, UCLA locality 4347.
- 14-15. *Gyrodes canadensis* Whiteaves, 1903. USNM 468575. Loc. 7, USGS Mesozoic locality M8600.
16. *Turritella chicoensis pescaderoensis* Arnold, 1908. USNM 468576. Loc. 3, USGS Mesozoic locality M8752.
- 17-18. *Calva* (*Egelicalva*) *bowersiana* (Cooper, 1894), left valve. LACMIP 8196. Loc. 6, UCLA locality 4348.