

North and South America throughout the Cenozoic (Johnson, 1987). However, rock-inhabiting organisms are rarely preserved (Carter, 1972), because the habitat is destructive to the shells (Woodring, 1931; Kotaka and Ogasawara, 1974; Parsons and Brett, 1991).

Many tropical abalone species occur in reef environments and tend to hide in cracks or underneath boulders. Upon death, the shells are likely to remain in the cryptic habitat and will eventually be incorporated into the biogenic limestone of the reef. Tectonic uplift or eustatic sea level change may expose these reefs above sea level, making them accessible [e.g., the Pleistocene record of *H. pustulata cruenta* Reeve, 1846, from the Red Sea (Hall and Standen, 1907)].

A major constituent of abalone shells is aragonitic platelets in the nacreous layer. This type of microstructure are very prone to disintegration, dissolution, and diagenetic change (Dodd, 1966). Disintegration accounts for the general scarcity of complete abalone specimens in the fossil record. The most frequent mode of abalone preservation is as internal and/or external molds (Fig. 1.1–1.3), occasionally with some attached shelly remains. Preserved shell fragments are all calcitic, indicating some diagenetic effects (cf. Dodd, 1966), because the inner nacreous layer of the shell is always composed of aragonite. In addition, Durham (1979b) observed recrystallization in *H. lomaensis*. The effects of diagenesis make the wide application of shell mineralogy in the taxonomy of fossil abalone doubtful, although results on the mineralogy of shells from Recent abalone are promising (see also previous section on phylogeny).

PALEOENVIRONMENTS

The habitat of abalone, with their large aperture, is the rocky shores, as is the case for other morphologically similar groups such as limpets. Not surprisingly fossil Haliotidae inhabited comparable habitats. Below we present supporting evidence.

Reef paleoenvironments.—Both fossil and Recent abalone are found in similar microhabitats within tropical reefs. *Haliotis antillesensis* has been found in a rudist-framework bioherm in the Upper Cretaceous of Puerto Rico (Sohl, 1992), part of the shallow tropical Tethys Sea (Kauffman and Sohl, 1974; Sohl, 1987). Associated species include abundant neritid gastropods, indicators for an intertidal to shallow subtidal hard substrate in the tropics. *Haliotis pustulata cruenta* has been found in a Pleistocene raised coral reef of the Red Sea area [Newton, 1900; Hall and Standen, 1907 (as *H. unilateralis* Lamarck, 1822); Brauwer et al. cited in Yaron, 1983] in association with the typical, molluscan reef fauna found in the Red Sea. The reconstructed paleoenvironment is fully congruent with the modern habitat of this species (Yaron, 1983; Geiger, 1996). For European sites, Davidaschvili (1937) and Krach (1981) interpreted a nearshore shallow-water reef environment for their *H. volhynica* based on a community analysis.

Shallow-water rocky paleoenvironments.—Fossil abalone also are reported from deposits associated with rocky shorelines, a common habitat for Recent members of the family. In the Cretaceous of California, *H. lomaensis* was found with an oyster, a serpulid worm, and encrusting calcareous algae, which together indicate a shallow-water, rocky environment (Durham, 1979b). In the Miocene of California, *H. palaea* Woodring, 1931, was discovered with other molluscan genera indicative of rocky substrate, i.e., *Tegula*, *Arca*, *Crassadoma* [as *Hinnites*], *Lima*, and *Mytilus* (Woodring, 1931). Only the limids may be freely mobile as adults and associated with sandy substrates. *Haliotis elsmersensis* Vokes, 1935, from the Pliocene of California, however, was found admixed with infaunal bivalves (*Chione*) as well as with rocky shore bivalves (*Arca*). This mixed fauna indicates some transport (Vokes, 1935). Similarly, *H. cracherodii* Leach,

1814, from the Pleistocene of northwestern Baja California (Ad-dicott and Emerson, 1959). *Haliotis rufescens* Swainson, 1822, has been found in a Pliocene shallow-water channel deposit with rock-dwelling species, as well as infaunal species (Groves, 1991). Additional, indirect evidence of California abalone paleohabitat stems from the type specimen of the Late Pleistocene vermetid gastropod *Petalococonchus anellum* (Mörch, 1861), which is attached to a shell of an unspecified abalone (Grant and Gale, 1931). Vermetids are common in shallow, rocky habitats.

In Japan, *H. kurosakiensis* Kotaka and Ogasawara, 1974 (upper Miocene); *H. fujiokai* Hatai, Kotaka, and Noda, 1937 (middle Miocene); and *H. kochibei* Hatai and Nisiyama, 1952 (Pliocene), have been found in strata containing typical rock-inhabiting molluscan genera such as *Ostrea*, *Arca*, and *Acmaea* admixed with sand-dwellers, such as *Macoma* and *Lucinoma* (Hatai et al., 1970; Kotaka and Ogasawara, 1974; Noda et al., 1995). As with *H. elsmersensis*, some transport of the original fauna must be assumed, but a rocky shore habitat for abalone is indicated.

In summary, available evidence indicates that the ecology of the fossil abalone is comparable to that of Recent species (see e.g., Crofts, 1929; Cox, 1962; Shepherd, 1973). Particularly, no change in the depth distribution (=on-shore/off-shore pattern) can be detected, which has been documented during the geological history of a number of invertebrate taxa (Hickman, 1984; Bottjer and Jablonski, 1988). It may seem farfetched to consider an on-shore/off-shore pattern in a largely herbivorous organism, however, one abalone species (*H. cracherodii*) is known to feed on bacterial mats as adults (Stein, 1984). Juvenile abalone feed on diatoms and the associated biofilms (Kawamura et al., 1998, for review). Accordingly, the potential to exploit deep water habitats directly or by neoteny is present in abalone.

TIME RANGE

Fossil abalone have been found from the Late Cretaceous through the late Pleistocene, with a conspicuous hiatus in the early Paleogene (see Appendix for details). A controversy has long existed regarding the true identity and systematic affinities of some of the earliest records (Late Cretaceous: Maastrichtian) in this family: *H. antiqua* Binkhorst, 1861, from the Netherlands; *H. cretacea* Lundgren, 1894, from Sweden, and *H. lomaensis* from San Diego County, California (Binkhorst, 1861; Lundgren, 1894; Anderson, 1902). Davies and Eames (1971) considered none of these records to represent abalone. Woodring (1931) doubted that *H. lomaensis* is an abalone, because of poor preservation; the only diagnostic characteristic preserved was the flat shell. In addition, Knight et al. (1960: 221) cited the Cretaceous record of *Haliotis* with a query. Vokes (1935) considered *H. lomaensis* to be an abalone, but most likely a young specimen. Equally, Sohl (1992) discussed these three Cretaceous taxa and concluded that only *H. lomaensis* is an abalone, as did Durham (1979b) and Lindberg (1992). More recently, Sohl (1992) described a second Late Cretaceous abalone from the Maastrichtian of Puerto Rico and Jamaica, *H. antillesensis*. We consider only *H. lomaensis* and *H. antillesensis* to be Cretaceous abalone, both from Maastrichtian rocks.

Within the Tertiary, Delhaes (1909) and Beu and Maxwell (1990) reported abalone from the Eocene. However, the Standard European Stages in Delhaes (1909) are Miocene (Lindberg, 1992), and consequently only one record, from New Zealand, is of Eocene age. Three factors may be considered to explain the hiatus in the fossil record between the Maastrichtian and the Miocene. First, Paleogene rocky-shore habitats are hardly preserved (Lindberg and Squires, 1990). Second, in general there is a taphonomic bias against rock-clinging mollusks (Sohl, 1992). Third,




FIGURE 3—Localities of fossil abalone listed in Appendix. The two inserts provide details for California, and Honshu, Japan; the location of each area on the larger map is shaded. Overlapping symbols indicate different geological times being represented at the same locality.

by comparison to ecologically similar taxa such certain limpets (Acmaeidae), which are known from the Triassic through the Recent (Knight et al., 1960), Cretaceous abalone can be inferred as rare with just two localities. K-T extinction may have reduced the abundance of abalone to such low densities that they do not show up in the fossil record of the Paleocene. However, complete extinction at the K-T boundary with convergent evolution of the same shell morphology by another gastropod group in the Eocene-Oligocene is unlikely. Cretaceous and Tertiary/Recent specimens share many similarities, such as the row of tremata, the flattening of the shell, and the high translation value of the spire. This combination of structural elements is not known from any gastropod, living or extinct. These similarities are best explained by common descent, rather than two separate radiations.

GEOGRAPHIC DISTRIBUTION

In general, one might expect the Neogene distribution of taxa to be fairly congruent with its present-day distribution. However, the very few recorded fossil abalone specimens from the central Indo-Pacific (Fig. 3; Appendix) is surprising, because it houses the highest diversity of Recent Haliotidae (Delhaes, 1909; Lindberg, 1992). This discrepancy may be associated with the greater paleontological effort being carried out in the United States and Europe as compared to the Indo-Pacific. In the few, more extensive studies in tropical settings [Hall and Standen, 1907 (Red Sea); Ladd and Hofmeister, 1945; Ladd, 1966 (Indo-Pacific)] abalone specimens have been recovered. Therefore, more extensive searches for fossil abalone in the Indo-Pacific, as well as in the Australian region, might close both this noticeable geographical gap for fossil abalone, as well as narrow the Cretaceous to late Eocene hiatus in the abalone record.

Geiger is currently preparing a phylogenetic analysis of the

Recent species of the family, and the biogeographical investigation will be able to evaluate the proposed models of radiation, summarized below:

1) Living abalone are most diverse in the Indo-Pacific, which may indicate that this is their center of radiation. However, this is certainly not a safe conclusion, as discussed by Lindberg (1992). It is unclear whether several small radiations or a single large-scale radiation took place within this family.

2) Talmadge (1963) proposed a "Pacific Rim" model, in which he postulated that abalone were originally found on an island arc from Japan to northern Australia and radiated from this area to California, southern Australia, and the Indo-Pacific. This proposal is similar to one of the two models in Lee and Vacquier (1995).

3) A third scenario is based on chromosomal data, which have been documented for 10 species (Table 2). Because basal gastropods have a very low diploid number of chromosomes (18–20: Patterson, 1967; Haszprunar, 1988), a model of progressive increase in chromosome number can be postulated, from a low of diploid number of 28 in *H. tuberculata* from the Mediterranean, to 32 in Indo-Pacific species, to 36 in North Pacific species. This would suggest that *H. tuberculata* is a relict species from the ancient Tethys Sea, and that abalone dispersed eastward, which is in agreement with the eastward dispersal pattern in the Pacific (Briggs, 1995). Note the marked contrast to the westward dispersal of other mollusks (Squires, 1987). In westward dispersing families molluscan families such as Cypraeidae, Volutidae, and Ranellidae, however, larvae are teleplanktic. Abalone, on the other hand, have a rather limited dispersal capability, the precise extent of which is currently debated (Tegner and Butler, 1985; Prince et al., 1987; McShane et al.,

TABLE 2—Chromosome number in Recent *Haliotis* spp. as indicated in the respective source. H, Haploid number; D, Diploid number; O, Geographic occurrence; EM, European-Mediterranean; IP, Indo-Pacific; NP, North Pacific.

Taxon	H	D	O	Source
<i>Haliotis tuberculata</i>	14	28	EM	Colombera and Tagliaferri, 1983
		28	EM	Arai and Wilkins, 1986
<i>H. lamellosa</i> ¹	14		EM	Colombera and Tagliaferri, 1983
<i>H. aquatilis</i> ³	17	34	IP	Nakamura, 1985
<i>H. diversicolor aquatilis</i> ²	16	32	IP	Nakamura, 1985, 1986
<i>H. diversicolor</i>		32	IP	Arai et al., 1988; Yang et al., 1998
<i>H. exigua</i> ⁴		32	IP	Arai et al., 1988
<i>H. planata</i> ⁵		32	IP	Arai et al., 1988
<i>H. asinina</i>		32	IP	Jarayanbhand et al., 1998
<i>H. ovina</i>		32	IP	Jarayanbhand et al., 1998
<i>H. varia</i>		32	IP	Jarayanbhand et al., 1998
<i>H. varia</i>	16	32	IP	Nakamura, 1986
<i>H. cracherodii</i>		36	NP	Minkler, 1977
<i>H. discus discus</i>		36	NP	Arai et al., 1982
<i>H. discus hannai</i>		36	NP	Arai et al., 1982
<i>H. madaka</i> ⁶		36	NP	Nakamura, 1986

¹ *Haliotis lamellosa* has been shown to be a synonym/ecomoroph of *Haliotis tuberculata* (Lee and Vacquier, 1995).

² As *Haliotis japonica*.

³ As *Haliotis diversicolor aquatilis* in Nakamura (1985), but as *H. aquatilis* in Nakamura (1986).

⁴ As *Haliotis planata*.

⁵ As *Haliotis varia*.

⁶ As *Haliotis gigantea*.

1988; Wells and Keesing, 1990; Shepherd et al., 1992). Under the Tethyan origin model, the Upper Cretaceous abalone from California and the Caribbean would not be early representatives of the family and the origin of the family would have to be much earlier. A similar model has been proposed by Lee and Vacquier (1995) and is preferred by Eagle (1996).

None of these three biogeographic models has more than anecdotal evidence. The phylogenetic trees published to date (Brown and Murray, 1992; Brown, 1993; Lee and Vacquier, 1995) are all unrooted, and species in a basal position may not be truly basal.

It is possible that in the Pleistocene, species of *Haliotis* migrated between Asia and North America (Grant and Gale, 1931, p. 96). Biochemical data support this hypothesis. In a tree based on the cDNA sequences of the fertilization protein lysin (Vacquier and Lee, 1993, for review) of 22 abalone species, all California species cluster closely together with the west Pacific species *H. discus* Reeve, 1846, and *H. gigantea* Gmelin, 1791, whereas another species of the latter region (*H. diversicolor* Reeve, 1846) is far removed from the western Pacific species group (Lee and Vacquier, 1992, 1995). This pattern has been corroborated with allozyme frequency data from 17 abalone taxa (Brown, 1993). The California species have the most derived character states, which suggests an east-to-west progression. All three of the above models imply favorable conditions for dispersal of abalone between the east and west coasts of the north Pacific, with subsequent reproductive isolation of the amphipacific populations. Glaciation events could facilitate dispersal across the Pacific as sea level dropped, because the distance between the two shores at any latitude would be somewhat shortened (see also Lee and Vacquier, 1995). On the other hand, the distance across the Pacific is extensive, hence, dispersal across the Pacific must be termed a rare chance event. Accordingly, one may debate the possible contribution of glaciation to dispersal. Migration at northern latitudes along the Bering Bridge seems unlikely due to the absence of any Recent species generally beyond 50° north or south (60°N for *H. k. kamtschaticana*).

ACKNOWLEDGMENTS

D. Bottjer [University of Southern California (USC)], J. H. McLean (Natural History Museum of Los Angeles County), R.

Squires (California State University Northridge), the reviewers D. R. Lindberg and D. Dockery, and technical editor L. C. Anderson critically reviewed various drafts of the manuscript. D. Lindberg [University of California Berkeley (UCMP)], P. Mikkelsen (American Museum of Natural History, New York), G. Rosenberg and N. Gilmore [Academy of Natural Sciences Philadelphia (ANSP)], P. Lozouet and P. Maestrati (Muséum National d'Histoire Naturelle, Paris), and J. Todd (The Natural History Museum, London) gave access to some of the specimens mentioned. A. Matsukuma (Kyushu University), and staff at Allan Hancock Foundation Library (USC) and the Research Library of the Natural History Museum of Los Angeles County kindly helped with some of the obscure references. D. Stramski (USC) and A. Warén (Swedish Museum of Natural History) translated a Polish and a Swedish paper, respectively, for us. T. Rodica (Muzeul de Istorie Naturala "Grigore Antipa", Bucharest) drew our attention to a Romanian publication and translated the relevant parts. Additional information was provided by C. Hedegaard (UCMP), H.-D. Laatsch and M. Jakubeit (Wolfsburg, Germany), and P. Jarayanbhand (Thailand). The ANSP visit by DLG was made possible by a Jessup Fellowship. P. Irvine (California Division of Mines and Geology) collected and donated the figured specimen of *Haliotis* sp. from the Topanga Canyon Formation to LACMIP.

REFERENCES

- ABBOTT, R. T., AND S. P. DANCE. 1983. *Compendium of Seashells*. E. P. Dutton, Inc., New York, 411 p.
- ADDICOTT, W. O., AND W. K. EMERSON. 1959. Late Pleistocene invertebrates from Punta Cabras, Baja California, Mexico. *American Museum Novitates*, 1925:1-33.
- AGUAYO, C. G., AND M. L. JAUME. 1947. *Gastropoda—Haliotidae*. *Catálogo Molluscos de Cuba*, No. 140: 1 p.
- ANDERSON, F. M. 1902. Cretaceous deposits of the Pacific coast. *Proceedings of the California Academy of Sciences*, 3rd Series, Geology, 2:1-154.
- ARAI, K., AND N. P. WILKINS. 1986. Chromosomes of *Haliotis tuberculata*. *Aquaculture*, 58:305-308.
- , E. FUJINO, AND K. MASAHIRO. 1988. Karyotype and zymogram differences among three species of abalone *Haliotis planata*, *H. varia*, and *H. diversicolor*. *Bulletin of the Japanese Society of Scientific Fisheries*, 54:2055-2064.
- , H. TSUBAKI, Y. ISHITANI, AND K. FUJINO. 1982. Chromosomes

- of *Haliotis discus hannai* Ino and *H. discus* Reeve. Bulletin of the Japanese Society of Scientific Fisheries, 48:1689–1691.
- ARNOLD, R. 1903. The paleontology and stratigraphy of the marine Pliocene and Pleistocene of San Pedro, California. Memoirs of the California Academy of Sciences, 3:9–420.
- BANDEL, K. 1998. Scissurellidae als Modell für die Variationsbreite einer natürlichen Einheit der Schlitzbandschnecken (Mollusca, Archaeogastropoda). Mitteilungen des Geologisch-Paläontologischen Instituts der Universität Hamburg, 81:1–120.
- BARNARD, K. H. 1963. Contributions to the knowledge of South African marine Mollusca. Part IV. Gastropoda: Prosobranchia: Rhipidoglossa, Docoglossa. Tectibranchiata. Polyplacophora. Solenogastres. Scaphopoda. Annals of the South African Museum, 47:201–360.
- BARTSCH, P. 1940. The west American *Haliotis*. Proceedings of the United States National Museum, 89:49–57.
- BEU, A. G., AND P. A. MAXWELL. 1990. Cenozoic Mollusca of New Zealand. New Zealand Geological Survey Paleontological Bulletin, 58:1–518.
- BINKHORST, J.-T. 1861. Monographie des Gastéropodes et des Céphalopodes de la Craie Supérieure du Limbourg. C. Muquardt, Bruxelles, 83 p. [gastropods] + 44 p. [cephalopods].
- BOTTJER, D. J., AND D. JABLONSKI. 1988. Paleoenvironmental patterns in the evolution of post-Paleozoic benthic marine invertebrates. Palaios, 3:540–560.
- BRIGGS, J. C. 1995. Global Biogeography. Elsevier, Amsterdam, 452 p.
- BROWN, L. D. 1993. Biochemical genetics and species relationships within the genus *Haliotis* (Gastropoda: Haliotidae). Journal of Molluscan Studies, 59:429–444.
- , AND N. D. MURRAY. 1992. Genetic relationship within the genus *Haliotis*, p. 19–23. In S. A. Shepherd, M. J. Tegner, and S. A. Guzman del Prío (eds.), Abalone of the World: Biology Fisheries and Culture. Fishing News Books, Oxford.
- CARTER, R. M. 1972. Wanganui Strata of Komako District, Pohangina Valley, Ruahine Range, Manawatu. Journal of the Royal Society of New Zealand, 2:293–324.
- CHACE, E. P., AND E. M. CHACE. 1919. An unreported exposure of the San Pedro Pleistocene. Lorquinia, 2:1–3. [Should read 41–43].
- COATES, A. G., AND J. A. OBANDO. 1996. The geologic evolution of the Central American Isthmus, p. 21–56. In J. B. C. Jackson, A. F. Budd, and A. G. Coates (eds.), Evolution and Environment in Tropical America. University of Chicago Press, Chicago.
- COLOMERA, D., AND F. TAGLIAFERRI. 1983. Chromosomes from male gonads of *Haliotis tuberculata* and *Haliotis lamellosa* (Haliotidae, Archaeogastropoda, Mollusca). Caryologia, 36:231–234.
- COSSMANN, M. 1918. Essais de Paléoconchologie Comparée. Privately published, Paris, 388 p.
- COTTON, B. C. 1952. The Mollusca of the Adelaidean Stage. Australian Department of Mines Bulletin, 27:1–11, Appendix IV.
- COX, K. W. 1962. California abalones, family Haliotidae. California Division of Fish and Game, Fish Bulletin, 118:1–131.
- CROFTS, D. R. 1929. *Haliotis*. Liverpool Marine Biology Committee Memoirs, 29:1–174.
- DAUPHIN, Y., AND A. DENIS. 1995. Implications of statistical analyses based on localized chemical contents on some species of *Haliotis* shells (Mollusca, Archaeogastropoda). Annales des Sciences Naturelles—Zoologie et Biologie Animale, 16:7–19.
- , J. P. CUIF, H. MUTVEI, AND A. DENIS. 1989. Mineralogy, chemistry and ultrastructure of the external shell-layer in ten species of *Haliotis* with reference to *H. tuberculata* (Mollusca: Archaeogastropoda). Bulletin of the Geological Institution of the University of Uppsala New Series, 15:7–38.
- DAVIDASCHVILI, L. S. 1937. On the ecology of animals of the middle Miocene reefs of Ukrainian SSR. Problems of Paleontology, 2–3: 537–563.
- DAVIES, A. M., AND F. M. EAMES. 1971. Tertiary Faunas. Volume 1. The Composition of Tertiary faunas. American Elsevier Publishing Company, Inc., New York, 571 p.
- DELHAES, W. 1909. Beiträge zur Morphologie und Phylogenie von *Haliotis* Linné. Inaugural Dissertation, Bonn, Germany, 55 p.
- DODD, J. R. 1966. Process of conversion of aragonite to calcite with examples from the Cretaceous of Texas. Journal of Sedimentary Petrology, 36:733–741.
- DURHAM, J. W. 1979a. A fossil *Haliotis* from the Galápagos Islands. The Veliger 21:369–372.
- , 1979b. California Cretaceous *Haliotis*. The Veliger, 21:373–374.
- EAGLE, M. K. 1996. A new species of *Haliotis* (Mollusca) from the early Miocene basal Waitemata Beds, Rodney District, North Auckland. Records of the Auckland Institute and Museum, 33:15–171.
- EMERSON, W. K. 1980. Invertebrate faunules of late Pleistocene age, with zoogeographic implications from Turtle Bay, Baja California Sur, Mexico. The Nautilus, 94:67–89.
- , G. L. KENNEDY, J. F. WEHMILLER, AND E. KEENAN. 1981. Age relations and zoogeographic implications of late Pleistocene marine invertebrate faunas from Turtle Bay, Baja California Sur, Mexico. The Nautilus, 95:105–116.
- FINET, Y. 1993. Marine molluscs of the Galápagos Islands. Vita Marina, 42:33–39.
- FISCHER, W. 1993. Beiträge zur Kenntnis der rezenten und fossilen marinen Molluskenfauna Zyperns (1): Die Mollusken der Khrysokhou Bucht, sowie ein kleiner Beitrag zu den rezenten Foraminiferen. Club Conchylia Informationen, 25:37–46.
- FLEMING, A. E., AND P. W. HONE. 1996. Abalone culture (Special issue). Aquaculture, 140:1–195.
- FLEMING, C. A. 1952. Notes on the genus *Haliotis* (Mollusca). A new subgenus from New Zealand and a new species from the late Cenozoic of Ohope, Bay of Plenty. Transactions of the Royal Society of New Zealand, 80:229–232.
- FOSTER, R. W. 1946. The family Haliotidae in the Western Atlantic. Johnsonia, 2:36–40.
- GEIGER, D. L. 1991. Einige Aspekte über die Biologie von *Haliotis*. Club Conchylia Informationen, 23:87–108.
- , 1996. Haliotids in the Red Sea, with neotype designation for *Haliotis unilateralis* Lamarck, 1822 (Gastropoda: Prosobranchia). Revue Suisse de Zoologie, 103:339–354.
- , 1998a. Recent genera and species of the family Haliotidae Rafinesque, 1815 (Gastropoda: Vetigastropoda). The Nautilus 111:85–116.
- , 1998b. Note on the identity of *Haliotis clathrata* Lichtenstein, 1794 (not Reeve, 1846). Molluscan Research, 19:157–159.
- , AND K. A. STEWART. 1998. Case 3036. Suppression of *Haliotis clathrata* Lichtenstein, 1794, to preserve *H. clathrata* Reeve, 1846, and *H. elegans* Philippi, 1844 (Mollusca, Gastropoda). Bulletin of Zoological Nomenclature, 55:209–211.
- GHISOTTI, F. 1964. Genus *Haliotis* Linneo, 1758. Pubblicazioni mensile edita della unione malacologica Italiana, 4 p.
- GLIBERT, M. 1962. Les Archaeogastropoda fossiles du Cénozoïque étranger des collection de l'Institut Royal des Sciences Naturelles de Belgique. Mémoires de l'Institut Royal des Sciences Naturelles de Belgique, Série 2, 68:3–131.
- GRANT, U. S., IV, AND H. R. GALE. 1931. Catalogue of the marine Pliocene and Pleistocene Mollusca of California and adjacent regions. Memoirs of the San Diego Society of Natural History, 1:1–1036.
- GROVES, L. T. 1991. Paleontology and biostratigraphy of the Plio-Pleistocene lower Saugus Formation, Santa Susana Mountains, Southern California. M.Sc. thesis, California State University, Northridge, 383 p.
- HALL, W. J., AND R. STANDEN. 1907. On the Mollusca of a raised coral reef on the Red Sea coast. Journal of Conchology, 12:65–68.
- HANZAWA, S., A. KIYOSHI, AND F. TAKAI. 1961. Catalogue of Type-Specimens of Fossils in Japan. Palaeontological Society of Japan, twenty-fifth Anniversary Volume, 422 p.
- HARRIS, G. F. 1897. Catalogue of Tertiary Mollusca of the Department of Geology British Museum (Natural History), Pt. I. The Australasian Tertiary Mollusca. British Museum of Natural History, London, 407 p.
- HARRY, H. W. 1966. *Haliotis pourtalesii* Dall, 1881 from Yucatan. The Veliger, 8:207–208.
- HASZPRUNAR, G. 1988. On the origin and evolution of major gastropod groups, with special reference to the Streptoneura. Journal of Molluscan Studies, 54:367–441.
- HATAI, K., AND S. NISIYAMA. 1952. Check list of the Japanese Tertiary marine Mollusca. The Science Reports of the Tôhoku University, Second Series (Geology) Special Volume, 3: 464 p.
- , T. KOTAKA, AND H. NODA. 1970. Some marine fossils from southwest of Shiroishi City, Miyagi Prefecture, northeast Honshu, Japan. Saito Ho-on Kai Museum Research Bulletin, 39:13–23.

- HEMPHILL, H. 1907. Notes on the genus *Haliotis*. Transactions of the San Diego Society of Natural History, 1:56-60.
- HENDERSON, J. B. 1915. Rediscovery of Pourtales' *Haliotis*. Proceedings of the United States National Museum, 48:659-661.
- HEDEGAARD, C., AND H.-R. WENK. 1998. Microstructure and texture pattern of mollusc shells. Journal of Molluscan Studies, 64:133-136.
- HERTLEIN, L. G. 1934. Pleistocene mollusks from the Tres Marias Islands, Cedros Island, and San Ignacio Lagoon, Mexico. Bulletin of the Southern California Academy of Sciences, 22:59-73.
- . 1937. *Haliotis koticki*, a new species from the lower Miocene of California. Bulletin of the Southern California Academy of Sciences, 36:93-97.
- , AND U. S. GRANT, IV. Unpublished manuscript. The Geology and Paleontology of the marine Pliocene of San Diego, California (Paleontology: Gastropoda), Pt. 2C. [Available in LACM-Malacology/Invertebrate Paleontology reprint collection]
- HICKMAN, C. S. 1984. Composition, structure, ecology, and evolution of six Cenozoic deep-water mollusk communities. Journal of Paleontology, 58:1215-1234.
- , AND J. H. LIPPS. 1985. Geologic youth of Galápagos Islands confirmed by marine stratigraphy and paleontology. Science, 227:1578-1580.
- , AND J. H. MCLEAN. 1990. Systematic revision and suprageneric classification of trochacean gastropods. Natural History Museum of Los Angeles County Science Series, 35:1-169.
- HÖRNES, M. 1856. Die fossilen Mollusken des Tertiär-Beckens von Wien. Volume I. Univalven. K. K. Hof- und Staatsdruckerei, Wien, 736 p.
- INO, T. 1952. Biological studies on the propagation of Japanese abalone (Genus *Haliotis*). Bulletin of the Tokai Regional Fisheries Research Laboratory, 5:1-102.
- ITOIGAWA, J., AND S. TOMIDA. 1982. *Miohaliotis amabilis*, a new haliotid fossil from the Miocene Mizunami group, with special reference to fossil haliotid fauna in Neogene and Quaternary of Japan. Bulletin of the Mizunami Fossil Museum, 9:1-14.
- JARAYANBHAND, P., R. YOM-LA, AND A. POPONGVIWAT. 1998. Karyotypes of marine molluscs in the family Haliotidae found in Thailand. Journal of Shellfish Research, 17:761-764.
- JOHNSON, M. E. 1987. Why are rocky shores so uncommon? Journal of Geology, 96:469-480.
- JORDAN, E. K., AND L. G. HERTLEIN. 1926. Contribution to the geology and paleontology of the Tertiary of Cedros Island and adjacent parts of Lower California. Proceedings of the California Academy of Sciences, 4th series, 15:409-464.
- , AND —. 1936. The Pleistocene fauna of Magdalena Bay, Lower California. Contributions from the Department of Geology of Stanford University, 13:107-173.
- JUNG, P. 1968. Fossil *Pleurotomaria* and *Haliotis* from Barbados and Carriacou, West Indies. Eclogae geologicae Helvetiae, 61:593-605.
- KAICHER, D. 1981. Card Catalogue of World Shells. Pack 28, *Haliotis*. Privately published. St. Petersburg, Florida.
- KANAKOFF, G. P., AND W. K. EMERSON. 1959. Late Pleistocene invertebrates of the Newport Bay area, California. Los Angeles County Museum of Natural History Contributions in Science, 31:1-47.
- KAUFFMAN, E. G., AND N. F. SOHL. 1974. Structure and evolution of Antillean Cretaceous rudist frameworks. Verhandlungen der Naturforschenden Gesellschaft in Basel, 84:399-467.
- KAWAMURA, T., R. D. ROBERTS, AND H. TAKAMI. 1998. A review of the feeding and growth of postlarval abalone. Journal of Shellfish Research, 17:615-625.
- KENNETT, J. P., AND S. SRINIVASAN. 1983. Neogene Planktonic Foraminifera. Hutchison Ross Publishing Company, Stroudsburg, Pennsylvania, 265 p.
- KENSLEY, B. 1972. Pliocene marine invertebrates from Langebaanweg, Cape Province. Annals of the South African Museum, 60:173-190.
- KERN, J. P. 1973. Early Pliocene marine climate and environment of the eastern Ventura Basin, southern California. University of California Publications in Geological Sciences, 96:1-117.
- KEEN, A. M., AND H. BENTSON. 1944. Check list of California Tertiary marine Mollusca. Special Papers of the Geological Society of America, 56:1-280.
- KLAPPENBACH, M. A. 1968. Notas malacológicas I. Comunicaciones zoológicas del Museo de historia natural de Montevideo, 9:1-7.
- KNIGHT, J. B., L. R. COX, A. M. KEEN, R. L. BATTEN, E. L. YOCH-ELSON, AND R. ROBERTSON. 1960. Systematic descriptions [Archaeogastropoda], p. 1169-1310. In R. C. Moore (ed.), Treatise on Invertebrate Paleontology, I, Mollusca 1. Geological Society of America, and University of Kansas Press, Kansas.
- KOTAKA, T., AND K. OGASAWARA. 1974. A new abalone form the Miocene of Aomori Prefecture, northeast Honshu, Japan. Venus, 33B:117-128.
- KRACH, W. 1981. The Baden reef formations in Roztocze Lubelskie. Prace Geologiczne, 121:5-91.
- LADD, H. D. 1966. Chitons and gastropods (Haliotidae through Adeirobidae) from the western Pacific islands. United States Geological Survey Professional Paper, 531:1-98.
- , AND J. E. HOFFMEISTER. 1945. Geology of Lau, Fiji. Bernice P. Bishop Museum Bulletin, 181:3-399.
- LECOINTRE, G., K. J. TINKLER, AND H. G. RICHARDS. 1967. The marine Quaternary of the Canary Islands. Proceedings of the Academy of Natural Sciences of Philadelphia, 119:325-344, fig. 1, pls. 1-3.
- LEE, D. E., R. M. CARTER, R. P. KING, AND A. F. COOPER. 1983. An Oligocene rocky shore community from Mt. Luxmore, Fiordland. New Zealand Journal of Geology and Geophysics, 26:123-126.
- LEE, Y.-H., AND V. D. VACQUIER. 1992. The divergence of species specific abalone sperm lysin is promoted by positive Darwinian selection. Biological Bulletin, 182:97-104.
- , AND —. 1995. Evolution and systematics in Haliotidae (Mollusca: Gastropoda): inferences from DNA sequences of sperm lysin. Marine Biology, 124:267-278.
- LINDBERG, D. R. 1992. Evolution, distribution and systematics of Haliotidae, p. 3-18. In S. A. Shepherd, M. J. Tegner, and S. A. Guzman del Próo (eds.), Abalone of the World: Biology Fisheries and Culture. Fishing News Books, Oxford.
- , B. ROTH, M. G. KELLOGG, AND C. L. HUBBS. 1980. Invertebrate megafossils of Pleistocene (Sangamon Interglacial) age from Isla de Guadalupe, Baja California, Mexico, p. 41-62. In D. M. Power (ed.), The California Islands: Proceedings of a Multidisciplinary Symposium. Santa Barbara Museum of Natural History, Santa Barbara.
- , AND R. L. SQUIRES. 1990. Patellogastropods (Mollusca) from the Eocene Tejon Formation of southern California. Journal of Paleontology, 64:578-587.
- LIPPS, J. H. 1967. Age and environment of a marine terrace fauna, San Clemente Island, California. The Veliger, 9:388-398.
- LOZOUET, P. 1986. Les Gastéropodes prosobranches de l'Oligocène supérieur du bassin de l'Adour (Système, Paléoenvironnements, Paléoclimatologie, Paléobiogéographie). Diploma Thesis L'École Pratique des Hautes Etudes, Dijon, 475 p.
- LUDBROOK, N. H. 1954. The molluscan fauna of the Pliocene strata underlying the Adelaide Plains, Pt. I. Transactions of the Royal Society of Southern Australia, 77:42-64.
- . 1956. The molluscan fauna of the Pliocene strata underlying the Adelaide Plains, Pt. III, Scaphopoda, Polyplacophora, Gastropoda (Haliotidae to Tornidae). Transactions of the Royal Society of Southern Australia, 79:1-36.
- LUNDGREN, B. 1894. Jämförelse mellan Molluskfaunan i Mammillatus och Mucronata Zonera i Nordöstra Skåne (Kruistianstadsområdet). Kongl Svenska Vetenskaps-Akademiens Handlingar, 26:3-58.
- MACNEIL, F. S. 1960. Tertiary and Quaternary Gastropoda of Okinawa. United States Geological Survey Professional Paper, 339: 148 p.
- MAKIYAMA, J. 1927. Preliminary report on the Tertiary fossils from Kamiminochigun, Shinano. Chikyu, 8:181-188.
- MALATESTA, A. 1960. Malacofauna Pleistocenica di Grammichelle (Sicilia). Memorie per servire alla descrizione della carta geologica d'Italia, 12:9-392.
- MARINCOVICH, L. Jr. 1976. Late Pleistocene molluscan faunas from upper terraces of the Palos Verdes Hills, California. Los Angeles County Museum of Natural History Contributions in Science, 281:1-28.
- MARTINEZ, R. E., AND L. B. RUIZ. 1994. Nota acerca de la presencia del gastropodo *Haliotis (Padollus) pourtalesii* Dall, 1881 (Archaeogastropoda, Pleurotomariacea) en aguas del Caribe Venezolano. Acta Biologica Venezuelica, 15:63-64.
- MASUDA, K. 1966. Molluscan fauna of the Higashi-Innai Formation of Noto Peninsula, Japan—II; Remarks on molluscan assemblage and description of species. Transactions and Proceedings of the Palaeontological Society of Japan, New Series, 64:317-337.

- MCCOY, F. 1876. Prodomus of the Palaeontology of Victoria. Decade III. George Skinner, Melbourne, 40 p.
- MCLEAN, J. H. 1970. New species of tropical eastern Pacific Gastropoda. *Malacological Review*, 2:115–130.
- . 1984. A case for derivation of the Fissurellidae from the Bellerophonacea. *Malacologia*, 25:3–20.
- , AND D. L. GEIGER. 1998. New genera and species having the *Fissurisepta* shell form, with a generic level phylogenetic analysis (Gastropoda: Fissurellidae). Los Angeles County Museum of Natural History Science Series, 475:32 p.
- MCSHANE, P. E., K. P. BLACK, AND M. G. SMITH. 1988. Recruitment Processes in *Haliotis rubra* (Mollusca: Gastropoda) and regional hydrodynamics in southeastern Australia imply localized dispersal of larvae. *Journal of Experimental Marine Biology and Ecology*, 124: 175–203.
- , D. R. SCHIEL, S. F. MERCER, AND T. MURRAY. 1994. Morphometric variation in *Haliotis iris* (Mollusca: Gastropoda): analysis of 61 populations. *New Zealand Journal of Marine and Freshwater Research*, 28:357–364.
- MINKLER, J. 1977. Chromosomes of the black abalone (*Haliotis cracherodii*). *Experientia*, 33:1143.
- MOISESCU, G. 1955. Stratigraphy and Molluscs' Fauna from the Tortonian and Sarmatian deposits from the Buituri locality—Romania. Bucharest. [Citation personal commun. T. Rodica].
- MULLINER, D. K. 1984. Notes on *Haliotis*. *Opisthobranch*, 16:20–22.
- MUTVEI, H., Y. DAUPHIN, AND J.-P. CUIF. 1985. Observations sur l'organisation de la couche externe du test des *Haliotis* (Gastropoda): un cas exceptionnel de variabilité minéralogique et microstructurale. *Bulletin du Muséum d'Histoire Naturelle*. Paris, 4^eème Serie 7 Section A, No. 1:73–91.
- NAKAMURA, H. K. 1985. The chromosomes of *Haliotis diversicolor aquatilis* (Archaeogastropoda: Haliotidae). *Malacological Review*, 18: 113–114.
- . 1986. Chromosomes of Archaeogastropoda (Mollusca: Prosobranchia), with some remarks on their cytotaxonomy and phylogeny. *Publications of the Seto Marine Biological Laboratory*, 31:191–267.
- NEWTON, R. B. 1900. Pleistocene shells from the raised beach deposits of the Red Sea. *Geological Magazine*, London, 7:500–514, 544–560.
- NIJSEN-MEYER, J. 1969. On the occurrence of *Haliotis pourtalesii* Dall, 1881, off Suriname (South America). *Zoologische Mededelingen*, 43: 203–206.
- NODA, H., R. WATANABE, AND Y. KIKUCHI. 1995. Pliocene marine molluscan fauna from the Hitachi Formation in the northeastern part of Ibaraki Prefecture, Japan. *Scientific Reports of the Institute of Geosciences, University of Tsukuba, Section B*, 16:39–93.
- NOMURA, S., AND H. NIINO. 1932. Fossil Mollusca from Izu and Hakone. *The Scientific Reports of the Tôhoku Imperial University, Sendai, Japan, Second Series (Geology)*, 15:1–188.
- ODÉ, H. 1986. Distribution and records of the marine Mollusca in the northwest Gulf of Mexico (A continuing monograph). *Texas Conchologist*, 22:69–73.
- ORR, P. C. 1960. Late Pleistocene marine terraces on Santa Rosa Island, California. *Bulletin of the Geological Society of America*, 71:1113–1120.
- OSTERGAARD, J. M. 1935. Recent and fossil Mollusca of Tongatabu. *Bernice P. Bishop Museum Bulletin*, 131:3–59.
- OZAWA, T., T. NAKAGAWA, AND K.-I. TAKEYAMA. 1986. Middle Miocene molluscan fauna of the Uchiura group, Wakasa Province, southwest Japan. *Palaeontological Society of Japan, Special Papers*, 29: 135–148.
- PARSONS, K. M., AND C. E. BRETT. 1991. Taphonomic processes and biases in modern marine environments: An actualistic perspective on fossil assemblage preservation, p. 22–65. *In* S. K. Donovan (ed.), *The Processes of Fossilization*, Belhaven Press, London.
- PATTERSON, C. M. 1967. Chromosome number and systematics in streptoneuran snails. *Malacologia*, 5:111–125.
- PAULUS, M. 1949. *Malacologie Marine et Saumatre de la Camargue (Anciens Cordons Littoraux, Littoral, Estuaires et Étangs)*. Imprimerie Delevaud, Saintes, 162 p.
- PHILIPPI, R. A. 1844. *Enumeratio Molluscorum Siciliae, Volume II. Sumptibus Eduardi Anton., Halis Saxonum*, 303 p.
- PICKERY, R. 1995. Recent Stomatellidae. *Gloria Maris*, 33:104–112.
- POWELL, A. W. B. 1938. Tertiary molluscan faunules from the Waitemata Beds. *Transactions and Proceedings of the Royal Society of New Zealand*, 68:362–379.
- PRINCE, J. D., T. L. SELLER, T. W. FORD, AND S. R. TALBOT. 1987. Experimental evidence for limited dispersal of haliotid larvae (Genus *Haliotis*; Mollusca: Gastropoda). *Journal of Experimental Marine Biology and Ecology*, 106:243–263.
- ROWLAND, R. W. 1968. Paleontology and Paleocology of the San Diego Formation in Northwestern Baja California. M.Sc. thesis, University of California, Davis, 61 p.
- . 1972. Paleontology and paleocology of the San Diego Formation in northwestern Baja California. *Transactions of the San Diego Society of Natural History*, 17:25–32.
- RUGGIERI, G. 1990. Una *Haliotis* del Miocene superiore (Saheliano) della Sicilia. *Bolletino Malacologico*, 25:349–354.
- SARSUA, H. 1968. Primer hallazgo del género *Haliotis* (Mollusca: Gastropoda) en Aguas Cubanas. *Poeyana*, A55:1–8.
- SCHREMP, L. A. 1981. Archaeogastropoda from the Pliocene Imperial Formation of California. *Journal of Paleontology*, 55:1123–1136.
- SHEPHERD, S. A. 1973. Studies on Southern Australian abalone (Genus *Haliotis*) I. Ecology of five sympatric species. *Australian Journal of Marine and Freshwater Research*, 24:217–257.
- , M. AVALOS-BORJA, AND M. O. QUINTANILLA. 1995b. Towards a chronology of *Haliotis fulgens*, with a review of abalone shell microstructure. *Marine and Freshwater Research*, 46:607–615.
- , R. W. DAY, AND A. J. BUTLER. 1995a. Progress in abalone fisheries research. *Marine and Freshwater Research*, 46:519–695.
- , D. LOWE, AND D. PARTINGTON. 1992. Studies on southern Australian abalone (genus *Haliotis*) XIII: larval dispersal and recruitment. *Journal of Experimental Marine Biology and Ecology*, 164:247–260.
- , M. J. TEGNER, AND S. A. GUZMÁN DEL PRÓO. 1992. *Abalone of the World: Biology, Fisheries and Culture*. Fishing News Press, Oxford, 608 p.
- SHIBATA, M. 1957. Some molluscan fossils from the eastern part of the Tanzawa Mountainland. *Transactions and Proceedings of the Paleontological Society of Japan, New Series*, 25:21–25.
- SIMONE, L. R. C. 1998. Morphology of the western Atlantic Haliotidae (Gastropoda, Vetigastropoda), with description of a new species from Brazil. *Malacologia*, 39:59–75.
- SINCLAIR, M. 1963. Studies on the Paua, *Haliotis iris* Martyn, in the Wellington district 1945–46. *Zoological Publications Victoria University, Wellington*, No. 35:1–16.
- SMITH, A. 1977. Fossil Abalone. *Of Sea and Shore*, 8:249.
- SMITH, A. B. 1994. *Systematics and the Fossil Record: Documenting Evolutionary Patterns*. Blackwell Scientific Publications, London, 223 p.
- SOHL, N. F. 1987. Cretaceous gastropods: contrast between Tethys and the temperate provinces. *Journal of Paleontology*, 61:1085–1111.
- . 1992. Upper Cretaceous gastropods (Fissurellidae, Haliotidae, Scissurellidae) from Puerto Rico and Jamaica. *Journal of Paleontology*, 66:414–434.
- SQUIRES, R. L. 1987. Eocene molluscan paleontology of the Whitaker Peak area, Los Angeles and Ventura Counties, California. *Los Angeles County Museum of Natural History Contributions in Science*, 338, 93 p.
- STEIN, J. L. 1984. Subtidal gastropods consume sulfur-oxidizing bacteria: Evidence from coastal hydrothermal vents. *Science*, 223:696–698.
- STEWART, K. A., AND D. L. GEIGER. 1999. Designation of lectotype for *Haliotis crebrisculpta* Sowerby, 1914, with a discussion of *H. clathrata* Reeve, 1846 (*non* Lichtenstein, 1794). *The Veliger*, 42:85–96.
- STRAUSZ, L. 1966. *Die Miozän-Mediterranen Gastropoden Ungarns*. Akadémia Kiadó, Budapest, 692 p.
- SUSUKI, T., AND C. J. STADUM. 1978. A Neogene section of northeastern San Clemente Island, California. *Los Angeles County Museum of Natural History Contributions in Science*, 299:24 p.
- SUTER, H. 1913. *Manual of the New Zealand Mollusca*, with [1915] Atlas of Plates. Government Printer, Wellington, 1120 p.
- TALAVERA, F. G., S. J. KARDAS, JR., AND H. G. RICHARDS. 1978. Quaternary marine mollusks from Tenerife, Canary Islands. *The Nautilus*, 92:97–102.
- TALMADGE, R. R. 1960. Variations in the Australian haliotid, *Haliotis cocciradiata* Reeve, 1846. *Journal of the Malacological Society of South Australia*, 4:60–63.

- . 1963. Insular haliotids in the western Pacific. *The Veliger*, 5:129–139.
- . 1964. The races of *Haliotis fulgens* Philippi (Mollusca: Gastropoda). *Transactions of the San Diego Society of Natural History*, 13: 369–376.
- . 1974. Some notes on the distribution and taxonomy of *Haliotis*. *The Veliger*, 16:411–412.
- TEGNER, M. J., AND R. A. BUTLER. 1985. Drift-tube study of the dispersal potential of green abalone (*Haliotis fulgens*) larvae in the Southern California Bight: implications for recovery of depleted populations. *Marine Ecology—Progress Series*, 26:73–84.
- TITGEN, R. H., AND T. J. BRIGHT. 1985. Notes on the distribution and ecology of the western Atlantic abalone, *Haliotis pourtalesii* Dall, 1881 (Mollusca: Gastropoda). *Northeast Gulf Sciences*, 7:147–152.
- TOMIDA, S. 1996. Late Neogene tropical and subtropical molluscan faunas from the south Fossa-Magna region, central Japan. *Bulletin of the Mizunami Fossil Museum*, 23:89–140.
- TRECHMANN, C. T. 1937. The base and top of the coral-rock in Barbados. *Geological Magazine*, London, 74:337–359.
- VACQUIER, V. D., AND Y.-H. LEE. 1993. Abalone sperm lysin: unusual mode of evolution of a gamete recognition protein. *Zygote*, 1:181–196.
- VALENTINE, J. W. 1957. Late Pleistocene faunas from the northwestern coast of Baja California, Mexico. *Transactions of the San Diego Society of Natural History*, 16:289–308.
- . 1958. Late Pleistocene megafauna of Cayucos, California and its zoogeographic significance. *Journal of Paleontology*, 32:687–696.
- . 1962. Pleistocene molluscan notes. 4. Older terrace faunas from Palos Verdes Hills, California. *Journal of Geology*, 70:92–101.
- VEDDER, J. G., AND R. M. NORRIS. 1963. Geology of San Nicolas Island, California. *United States Geological Survey Professional Paper*, 369:1–65.
- VOKES, H. E. 1935. A new species of *Haliotis* from the Pliocene of southern California. *Journal of Paleontology*, 9:251–252.
- . 1978. On fossil abalones. *Of Sea and Shore*, 9:46.
- WEBB, R. W. 1937. Paleontology of the Pleistocene of Point Loma, San Diego County, California. *Transactions of the San Diego Society of Natural History*, 8:337–348.
- WEINKAUFF, H. C. 1868. *Die Conchylien des Mittelmeeres, ihre geographische und geologische Verbreitung*. Band II *Mollusca cephalo*. Verlag Theodor Fischer, Cassel, 512 p.
- WELLS, F. E., AND J. K. KEESING. 1990. Population characteristics of the abalone *Haliotis roei* on intertidal platforms in the Perth metropolitan area. *Journal of the Malacological Society of Australia*, 11: 65–71.
- WILLETT, G. 1937. An upper Pleistocene fauna from the Baldwin Hills, Los Angeles County, California. *Transactions of the San Diego Society of Natural History*, 8:381–404.
- WOODRING, W. P. 1931. A Miocene *Haliotis* from southern California. *Journal of Paleontology*, 5:34–39.
- . 1932. A Miocene mollusk of the genus *Haliotis* from the Temblor Range, California. *Proceedings of the United States National Museum*, 81:1–4.
- , M. N. BRAMLETTE, AND W. S. W. KEW. 1946. Geology and paleontology of Palos Verdes Hills, California. *United States Geological Survey Professional Paper*, 207, 145 p.
- YABE, H., AND K. HATAI. 1941. Additional fossils from the Simaziri Beds of Okinawa-zima, Ryūkyū Islands, Japan. *Japanese Journal of Geology and Geography*, 18:71–78, pl. 7.
- YANG, H.-S., H.-C. CHEN, AND Y.-Y. TING. 1998. Induction of polyploidy and embryonic development of the abalone, *Haliotis diversicolor*, with temperature treatment. *American Malacological Bulletin*, 14:139–147.
- YARON, I. 1983. The Haliotidae of the Red Sea. *Levantina*, 42:486–493.

ACCEPTED 26 APRIL 1999

APPENDIX

List of fossil abalone taxa, as indicated in the original source, by first occurrence in the fossil record and locality compiled from the literature and museum specimens. General locations such as "California" have been omitted if more precise information is available for the same age. Taxa preceded by an asterisk (*) are living species. u, upper; m, middle; l, lower; BC, Baja California, Mexico; CA, State of California, USA; NZ, New Zealand; Pref., Prefecture; AMNH, American Museum of Natural History, New York; ANSP, Academy of Natural Sciences Philadelphia; BMNH, The Natural History Museum, London; LACMIP, Natural History Museum of Los Angeles County, Invertebrate Paleontology; MNHN, Muséum National d'Histoire Naturelle, Paris. Circumpacific geologic time intervals have been correlated with European ones according to Kennett and Srinivasan (1983).

Taxon	Geologic time interval	Locality	Literature source and museum specimens
CRETACEOUS			
<i>antillesensis</i>	Maastrichian	Puerto Rico and Jamaica	Sohl, 1992
<i>lomaensis</i>	Maastrichian	San Diego County, CA	Anderson, 1902; Woodring, 1931, 1932; Bartsch, 1940; Cox, 1962; Durham, 1979b
TERTIARY: Eocene			
sp.	Eocene, l	NZ	Beu and Maxwell, 1990; Eagle, 1996
Oligocene			
sp. aff. <i>waitemataensis</i>	Oligocene/Miocene	Cookson Volcanics, NZ	Beu and Maxwell, 1990 ¹
<i>benoisti</i>	Oligocene, u; Chattien Adour	St.-Paul-Lès-Dax, France	Delhaes, 1909; Lozouet, 1986
	Oligocene, u; Chattien Adour	Belus "Marcon", Landes, France	MNHN no #
	Miocene, l	St.-Martin-de-Hinx, France	MNHN no #
	Miocene, l	Cabanes, St.-Paul-Lès-Dax, France	MNHN no #
	Miocene, l	Meilhan, Landes, France	MNHN no #
	Miocene, l	Larrey, Saucats, Gironde, France	MNHN no #
	Miocene, l	Merignac, France	Cossmann, 1918
	Miocene	Gironde, France	Delhaes, 1909
sp.	Miocene, m	Mirebeau, Indre and Loire, France	MNHN no #
sp.	Miocene, u	Beugnon, France	MNHN no #
Miocene			
<i>amabilis</i>	Miocene, m	Gifu Pref., Japan	Itoigawa and Tomida, 1982
<i>kurosakiensis</i>	Miocene, u	Aomori Pref., Japan	Kotaka and Ogasawara, 1974; Itoigawa and Tomida, 1982; Lindberg, 1992

APPENDIX

Continued

Taxon	Geologic time interval	Locality	Literature source and museum specimens
<i>kochibei</i>	Miocene, l	Kanagawa Pref., Japan	Hatai and Nisiyama, 1952; Hatai et al., 1970
	Miocene, m	Ibaraki Pref., Japan	Kotaka and Ogasawara, 1974; Itoigawa and Tomida, 1982
<i>koikei</i>	Miocene, m-u	Ibaraki Pref., Japan	Hatai and Nisiyama, 1952
	Pliocene	Ishikawa Pref., Japan	Lindberg, 1992
	Pliocene	Ibaraki Pref., Japan	Shibata, 1957; Noda et al., 1995
<i>glabrosa</i>	Miocene, l	Kanagawa Pref., Japan	Hanzawa et al., 1961; Hatai et al., 1970; Itoigawa and Tomida, 1982
	Pliocene	Kanagawa Pref., Japan	Kotaka and Ogasawara, 1974
	Pliocene, l	South Fossa-Magna, Japan	Tomida, 1996
<i>koyamai</i>	Miocene, l-m	Shizuoka Pref., Japan	Nomura and Niino, 1932; Hanzawa et al., 1961; Hatai et al., 1970; Kotaka and Ogasawara, 1974; Itoigawa and Tomida, 1982
	Pliocene, l	Shizuoka Pref., Japan	Itoigawa and Tomida, 1982
	Miocene; Pliocene, u ²	South Fossa-Magna, Japan Nagano Pref., Japan	Tomida, 1996 Makiyama, 1927; Hatai and Nisiyama, 1952; Hanzawa et al., 1961; Hatai et al., 1970; Kotaka and Ogasawara, 1974; Itoigawa and Tomida, 1982
<i>notoensis</i>	Miocene, m	Ishikawa Pref., Japan	Masuda, 1966; Kotaka and Ogasawara, 1974; Itoigawa and Tomida, 1982
<i>moniwaensis</i>	Miocene, m	Fukui Pref., Japan	Ozawa et al., 1986
	Miocene	Miyagi Pref., Japan	Kotaka and Ogasawara, 1974; Itoigawa and Tomida, 1982
<i>fujiokai</i>	Miocene, m	Miyagi Pref., Japan	Hatai et al., 1970; Kotaka and Ogasawara, 1974; Itoigawa and Tomida, 1982
* <i>japonica koticki</i>	Miocene, l	Miyagi Pref., Japan	Hatai and Nisiyama, 1952; Lindberg, 1992
	Miocene, l	Santa Barbara County, CA	Hertlein, 1937; Bartsch, 1940; Keen and Benton, 1944
<i>lasia</i>	Miocene, u	San Luis Obispo County, CA	Woodring, 1932; Bartsch, 1940; Keen and Benton, 1944
<i>palaea</i>	Miocene	Los Angeles County, CA	Woodring, 1931; Bartsch, 1940; Keen and Benton, 1944
<i>flemingi</i>	Miocene, l	Kawau Isl, NZ	Powell, 1938; Eagle, 1996
* <i>iris</i>	Pleistocene	Whakatane, NZ	Beu and Maxwell, 1990; Lindberg, 1992
	Miocene	Cape Rodney, NZ	Harris, 1897
	Miocene	NZ	Suter, 1913 ¹⁰
<i>mathesonensis powelli</i>	Pliocene, u-Recent	NZ	Beu and Maxwell, 1990
	Pleistocene	Lyttleton, NZ	BMNH 98048
	Miocene, l	Leigh, Rodney District, NZ	Eagle, 1996
	Miocene, l	Mt. Luxmore, Fiordland, NZ	Fleming, 1952; Lee, et al., 1983 ^{3,4}
<i>waitematensis moorabolensis</i>	Pliocene	Komako, NZ	Carter, 1972 ⁴ ; Beu and Maxwell, 1990; Eagle, 1996
	Pleistocene	Bay of Plenty, NZ	Fleming, 1952
	Miocene, l	Kawau Isl., NZ	Powell, 1938 ³ ; Eagle, 1996
<i>naevosoides</i>	Miocene	Moorabool River, Australia	McCoy, 1876 ⁵
	Miocene	Victoria, Australia	Harris, 1897
	Miocene, u	Flemington, Australia	BMNH G1934
	Miocene	Flemington, Victoria, Australia	Lindberg, 1992 ⁵
	Miocene, u	Flemington, Australia	BMNH G1935
	Pliocene	Adelaide, Australia	Cotton, 1952; Ludbrook, 1954
<i>ovinoides</i> sp.	Pliocene	Adelaide and Melbourne, Australia	Ludbrook, 1956
	Pliocene	Royal Park, Melbourne, Australia	BMNH GG2373
	Pliocene	Flemington, Australia	McCoy, 1876; Harris, 1897
	Miocene, u	Victoria, Australia	McCoy, 1876
	Miocene, l	New Caledonia	Eagle, 1996
	Miocene, l: f	Fiji	Ladd and Hoffmeister, 1945; Ladd, 1966
* <i>clathrata</i> Reeve, 1846	Miocene, l: f	Fiji	Ladd and Hoffmeister, 1945 ⁵ ; Ladd, 1966
	Miocene, u: g	Tinian	Ladd, 1966 ⁷
<i>neuvillei monilifera</i>	Plio-, Pleistocene	Guam	Ladd, 1966 ⁷
	Miocene, lowest: Aquitanian	Gironde, France	Delhaes, 1909
	Miocene, l: Elveziano	Turin, Italy	Delhaes, 1909
<i>anomiaeformis tuberculata</i> var.	Miocene	Turin, Italy	Glibert, 1962
	Miocene, l: Elveziano	Turin, Italy	Delhaes, 1909
	Miocene, l: Elveziano	Turin, Italy	Delhaes, 1909
<i>tuberculata lamellosoides</i>	Miocene	Miodoborow, W. Ukraine	Davidaschvili, 1937; Krach, 1981
	Pliocene	Cserhát Mountains, Hungary	<i>vide</i> Strausz, 1966
<i>ovata</i> sp.	Pliocene	Piemont, Italy	Delhaes, 1909
	Miocene	Italy	Delhaes, 1909; Ruggieri, 1990
<i>tuberculata volhynica</i>	Miocene	Tétényer Plateau, Hungary	Strausz, 1966
	Miocene	Vienna Bassin, Austria	H. D. Laatsch, personal commun.
	Miocene	Roztocze Lubelskie, Poland	Krach, 1981
	Miocene	Poland and Italy	Ghisotti, 1964; Ruggieri, 1990
	Miocene	Austria, Romania, Bulgaria, West Ukraine	Krach, 1981

APPENDIX

Continued

Taxon	Geologic time interval	Locality	Literature source and museum specimens
	Tortonian/Sarmatian	Buituri, Romania	Moiescu, 1955
	Miocene, l	Austria and Hungary	Delhaes, 1909
	Miocene	Cserhát Mountains, Hungary	Strausz, 1966
	Tertiary	Gauderndorf, Austria	Hörnes, 1856
	Miocene	Malta	BMNH 27501
	Miocene, u	Phoukasa, Cyprus	BMNH GH3252-5
	Tertiary	Lebanon	BMNH no #
	?	Asti and Turin, Italy	Hörnes, 1856
<i>torrei</i>	Miocene, u	Sicily, Italy	Fuggieri, 1990
sp.	Miocene	Mallorca, Balearic Islands, Spain	H. D. Laatsch, personal commun.
sp.	Miocene	Cyprus	Hertlein, 1937
sp.	Miocene, u (?)	Cyprus	H. D. Laatsch, personal commun.
sp.	Miocene	Turkey (Asia minor)	Hertlein, 1937
		Pliocene	
<i>*gigantea</i>	Pliocene	Ishikawa Pref., Japan	Hatai et al., 1970
	Pleistocene, l	Ishikawa Pref., Japan	Itoigawa and Tomida, 1982
<i>gigantoides</i>	Pliocene	Okinawa Pref., Japan	Yabe and Hatai, 1941; Hanzawa et al., 1961; Hatai et al., 1970; Kotaka and Ogasawara, 1974
<i>*discus</i>	Pliocene, u	Ishikawa Pref., Japan	Itoigawa and Tomida, 1982
<i>*diversicolor</i>	Pliocene, u	Okinawa Pref., Japan	Itoigawa and Tomida, 1982
	Plio-, Pleistocene	Okinawa Pref., Japan	MacNeil, 1960
<i>*sieboldii</i>	Pliocene, u	Japan	Itoigawa and Tomida, 1982
<i>*pourtalesi</i>	Pliocene, early	Riverside County, CA	Schremp, 1981
<i>elsmerensis</i>	Pliocene, early	Los Angeles County, CA	Vokes, 1935; Bartsch, 1940; Keen and Bentson, 1944
<i>*fulgens</i>	Pliocene, l	Los Angeles County, CA	Woodring, 1931; Grant and Gale, 1931; Vokes, 1935
	Pliocene	NW BC, Mexico	Rowland, 1968, 1972
	Pleistocene	Los Angeles County, CA	Arnold, 1903; Grant and Gale, 1931; Woodring, 1931; Woodring et al., 1946
	Pleistocene	San Diego County, CA	Arnold, 1903; Grant and Gale, 1931; Woodring, 1931; AMNH 12409; AMNH 12411
	Pleistocene	San Clemente Island, CA	Lipps, 1967
	Pleistocene	Orange County, CA	Kanakoff and Emerson, 1959
	Pleistocene	BC, Mexico	Hertlein, 1937
	Pleistocene	BC Sur, Mexico	Jordan and Hertlein, 1926; Emerson, 1980; Emerson et al., 1981
<i>*cracherodii</i>	Pliocene	CA	Lindberg, 1992
	Pliocene/Pleistocene	Punta China, BC, Mexico	ANSP 31487
	Pleistocene	San Diego County, CA	Webb, 1937; Glibert, 1962; this study: LACMIP loc. 12015; AMNH 12409
	Pleistocene	Santa Barbara Island, CA	this study; LACMIP locs. 326, 329, 5068
	Pleistocene	Santa Rosa Island, CA	Orr, 1960
	Pleistocene	San Nicolas Island, CA	Vedder and Norris, 1963
	Pleistocene, u	San Clemente Island, CA	Susuki and Stadum, 1978
	Pleistocene	Los Angeles County, CA	Chace and Chace, 1919; Willett, 1937; Woodring et al., 1946; Marincovich, 1976; this study: LACMIP loc. 10439
	Pleistocene	Orange County, CA	Kanakoff and Emerson, 1959
	Pleistocene	San Luis Obispo County, CA	Valentine, 1962
	Pleistocene	NW BC, Mexico	Addicott and Emerson, 1959; Hertline, 1934; Valentine, 1957; this study: LACMIP locs. 23161, 22715-22718, 8220-8224; 10131, 23162
<i>*rufescens</i>	Pliocene	Humboldt County, CA	Woodring, 1931
	Pliocene	Ventura County, CA	Woodring, 1931; Grant and Gale, 1931; Groves, 1991
	Pliocene	Los Angeles County, CA	Vokes, 1978
	Pleistocene	San Quintin, BC, Mexico	AMNH 12410
	Pleistocene	Monterey County, CA	Glibert, 1962
	Pleistocene	Santa Barbara Island, CA	this study: LACMIP locs. 5066-5067
	Pleistocene	Santa Rosa Island, CA	Orr, 1960
	Pleistocene	Santa Clemente Island, CA	Lipps, 1967
	Pleistocene	San Nicolas Island, CA	Vedder and Norris, 1963
	Pleistocene	Anacapa Island, CA	this study: LACMIP locs. 24381, 24383
	Pleistocene	San Luis Obispo County, CA	Valentine, 1958
	Pleistocene	Los Angeles County, CA	Chace and Chace, 1919; Grant and Gale, 1931; Woodring et al., 1946; Valentine, 1962; Marincovich, 1976
	Pleistocene	Orange County, CA	Kanakoff and Emerson, 1959
	Pleistocene	San Diego County, CA	Webb, 1937; this study: LACMIP loc. 11701
	Pleistocene	NW BC, Mexico	Valentine, 1957
	Pleistocene	BC, Mexico	this study: LACMIP loc. 22723, 22717

APPENDIX

Continued

Taxon	Geologic time interval	Locality	Literature source and museum specimens
*cf. <i>rufescens</i>	Pliocene Pleistocene	Isla Cedros, BC, Mexico Isla Cedros, BC, Mexico	Jordan and Hertlein, 1926; Woodring, 1931 Jordan and Hertlein, 1926; Grant and Gale, 1931
* <i>corrugata</i>	Pleistocene Pliocene Pliocene, m Pleistocene, m Pleistocene, m	Los Angeles County, CA Ventura County, CA Ventura County, CA Orange County, CA San Diego County, CA	Woodring, 1931 Woodring, 1931 Grant and Gale, 1931 Kanakoff and Emerson, 1959 Webb, 1937
* <i>assimilis</i>	Pliocene	CA	Lindberg, 1992
* <i>walallensis</i>	Pliocene	San Diego County, CA	Hertlein and Grant, MS
sp.	Pliocene	Los Angeles County, CA	Woodring, 1931; Kern, 1973
sp.	Pliocene	Ventura County, CA	Woodring, 1931
sp.	Pliocene	San Diego County, CA	Woodring, 1931
<i>santacruzensis</i>	Pliocene, u	Santa Cruz, Isla Galápagos	Durham, 1979a ⁸
* <i>ovina</i>	Pliocene, Pleistocene	Guam	Ladd, 1966
	Pliocene	—	Talmdage, 1974
	Pleistocene	Tongatapu, Tonga	Ostergaard, 1935
*cf. <i>varia</i>	Pliocene	Kankasanturai, Sri Lanka	BMNH G51083
* <i>tuberculata</i>	Pliocene	Piemont, Italy	Delhaes, 1909; BMNH G32151
	Pliocene	Palermo, Italy	BMNH 31287
	Pleistocene	Cyprus	Fischer, 1993
	Pleistocene	Sicily, Italy	Glibert, 1962
	Pleistocene	Ravagnee, Calabria, Italy	BMNH G31514
	Pleistocene	Panchia di Livorno, Italy	BMNH G11529
	Pleistocene	Malta	BMNH no #
	?	Favignana IIs. and Taranto, Italy	Philippi, 1844
	?	Sicily, Italy	Philippi, 1844; Weinkauff, 1868
*cf. <i>tuberculata</i>	Miocene, m	Loire Bassin, France	Glibert, 1962
sp.	1 Pliocene/u Miocene	Milazzo, Sicily, Italy	H. D. Laatsch, personal commun.
<i>saldanhae</i>	Pliocene	Langebaanweg, South Africa	Kensley, 1972
		Pleistocene	
* <i>australis</i>	Pleistocene, u-Recent	NZ	Beu and Maxwell, 1990
	?	Gisborne, North Island, NZ	Smith, 1977
* <i>virginea</i>	Pleistocene, u-Recent	NZ	Beu and Maxwell, 1990
* <i>cyclobates</i>	Pleistocene	Fleurieu Peninsula, Australia	Lindberg, 1992
* <i>emmae</i>	Pleistocene	Victoria, Australia	Lindberg, 1992
* <i>laevigata</i>	Pleistocene	Victoria, Australia	Lindberg, 1992
* <i>rubra</i>	Pleistocene	Victoria, Australia	Lindberg, 1992
* <i>kantschatkana</i>	Pleistocene	San Nicolas Island, CA	Vedder and Norris, 1963
*cf. <i>sorenseni</i>	Pleistocene	San Nicolas Island, CA	Vedder and Norris, 1963
*cf. <i>cracherodii</i>	Pleistocene	San Luis Obispo County, CA	Valentine, 1962
sp.	Pleistocene	Ventura County, CA	Woodring, 1931
	Pleistocene	Santa Barbara County, CA	this study: LACMIP loc. 12477
	Pleistocene	NW BC, Mexico	Valentine, 1957
* <i>fulgens guadalupensis</i>	Pleistocene	Isla Guadalupe, Mexico	Lindberg et al., 1980
<i>barbadensis</i>	Pleistocene	Whitehaven, Rugged Point and Spring Hall, Barbados	Trechmann, 1937; Jung, 1968; Durham, 1979a; BMNH GG4018-26; BMNH GG1913; BMNH GG4001-6; BMNH GG3977; BMNH GG9012-16 ⁹
<i>lamellosa</i>	Pleistocene	Camargue, France	Paulus, 1949
	Pleistocene	Monte Pellegrino, Italy	Ghisotti, 1964
	Pleistocene	Sicily, Italy	Malatesta, 1960; Glibert, 1962
sp.	Pleistocene/subfossil	Lanarca, Cyprus	H. D. Laatsch, personal commun.
* <i>tuberculata</i>	Pleistocene	Lanzarote, Canary Isl.	Lecointre et al., 1967
* <i>tuberculata coccinea</i>	Pleistocene	Tenerife, Canary Isl.	Talavera et al., 1978
* <i>midiae</i>	Pleistocene	Algoa Bay, South Africa	Barnard, 1963
* <i>unilateralis</i>	Pleistocene and Recent	Port Sudan, Sudan	Hall and Standen, 1907
	Pleistocene	Zanzibar, Tanzania	BMNH G41536
* <i>pustulata cruenta</i>	Pleistocene	Gulf of Suez, Egypt	Newton, 1900
	Pleistocene	Gharib Lighthouse, Red Sea	BMNH G24386
	Pleistocene	Mugarsim Isl., Sudan	BMNH G35528
	Pleistocene	Zanzibar, Tanzania	BMNH G41537

¹ *Haliotis* sp. is similar to *H. waitemataensis* (Beu and Maxwell, 1990) and the illustration strongly resembles *H. emmae*.

² The Shigarami Formation is interpreted as two different ages by the four publications cited.

³ Correlation of stages according to Beu and Maxwell (1990:8-9).

⁴ Originally reported as *Haliotis* sp. but corresponds exactly to the type of *H. powelli* illustrated in Beu and Maxwell (1990).

⁵ Delhaes (1909) interpreted these finds as Eocene.

⁶ *Haliotis tuvuthaensis* has been synonymized with *H. clathrata* Reeve, 1846 (Stewart and Geiger, 1999).

⁷ Plate 2, figure 5 illustrates a fairly round specimen superficially rather looking like *Haliotis ovina* than *H. clathrata* Reeve, 1846. However, this specimen is only 15.9 mm long. Small specimens of *H. clathrata* Reeve, 1846, are much more round than adult ones, which makes the identification of Ladd (1966) valid.

⁸ The Miocene of Durham (1979a) has been revised to late Pliocene by Hickman and Lipps (1985).

⁹ Durham re-dated the coral rock, in which the abalone was found from Jung's Pliocene to Pleistocene.

¹⁰ The identification of the specimen by Suter (1913) as *Haliotis iris* is rejected by Eagle (1996).