

TABLE 4.
Cancer amphioetus, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width														
				Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
1	F	L	45.0	10.7	8.1	16.3	22.3	6.1	4.3	11.5	2.6	4.3	7.2	2.4	11.2	11.7	AHF 16
2	M	L	26.4	5.6	4.3	8.6	11.9	3.1	2.2	6.1	1.4	2.1	3.9	1.3	5.8	6.2	AHF 17
3																	
4																	
5																	
6																	
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.757	.767														.0071	.928
T/H m	.570	.570														.0000	100.0
Hf/H m	.402	.375														.0191	4.91
H/L mf	.374	.360														.0099	2.69
H/L f	.597	.538														.0417	7.34
T/H f	.558	.620														.0438	7.45
H/L c	.958	.935														.0162	1.71

TABLE 5.
Cancer antennarius, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width															
				Manus					Movable finger			Fixed finger			Carpus		Locality	
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L		
1	?	R	?	18.5	14.1	27.0	35.0	11.0	8.1	18.7	5.5	6.0	10.9	4.6	17.2	21.3	UCMP (locality?)	
2	?	L	?	27.8	19.0	39.1	54.7	16.3	11.8	30.0	8.1	9.8	17.8	7.5	24.5	31.1	" "	
3	?	R	?	30.7	23.0	43.0	55.9	18.0	12.8	30.4	8.6	9.4	16.6	7.5	25.6	31.1	" "	
4	M	R	58.7	15.8	11.9	22.2	30.5	9.3	6.9	16.0	4.3	5.1	9.7	4.0	15.4	17.4	AHF i	
5	F	R	91.5	24.0	17.4	34.0	46.8	14.0	9.9	25.4	6.5	7.2	15.2	5.5	22.2	26.1	AHF 2	
6	F	L	91.5	24.5	18.2	35.3	48.5	14.4	10.0	26.4	6.9	7.3	15.6	6.1	22.2	26.7	AHF 2	
7	M	R	102.9	34.4	23.5	47.1	63.6	20.4	13.9	36.2	10.1	10.7	20.7	8.9	28.6	33.9	AHF 3	
8	M	L	102.9	34.6	23.6	47.0	63.6	20.7	14.3	35.8	10.3	10.7	20.6	9.4	29.4	34.6	AHF 3	
9	M	R	95.4	32.3	23.5	44.4	56.7	20.0	13.7	30.2	9.6	9.7	16.3	8.5	27.0	33.5	AHF 4	
10	F	L	74.8	19.0	13.7	26.8	36.7	10.8	8.3	19.9	5.6	6.5	12.6	5.2	18.0	21.1	AHF 4	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.762	.683	.750	.754	.726	.743	.683	.682	.727	.721	.738	.725	.756	.748		.0277	3.80
T/H m	.595	.587	.587	.588	.583	.588	.593	.598	.600	.568	.593	.598	.603	.600		.0090	1.53
Hf/H m	.324	.353	.306	.323	.300	.298	.315	.314	.301	.342	.335	.309	.321	.322		.0163	5.13
H/L mf	.433	.394	.422	.432	.390	.379	.384	.395	.453	.417	.419	.407	.400	.385		.0220	5.39
H/L f	.551	.551	.566	.526	.473	.468	.524	.527	.595	.516	.566	.542	.512	.528		.0346	6.50
T/H f	.767	.765	.798	.785	.763	.836	.820	.873	.877	.800	.827	.868	.786	.784		.0403	4.98
H/L c	.807	.788	.823	.885	.851	.832	.843	.850	.806	.853	.883	.853	.835	.853		.0266	3.17

TABLE 5.
Cancer antennarius, statistical data (measurements in millimeters). (continued)

specimen no.	sex	left or right	carapace width														
				Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	L1	T	H	L	T	H	L	T	H	L	
11	F	R	74.8	20.6	15.2	29.8	39.5	12.2	8.6	20.5	5.9	6.9	12.2	5.7	19.6	22.2	AHF 4
12	F	L	56.6	14.9	10.8	21.2	28.1	8.9	5.9	14.5	4.1	4.6	8.5	4.0	14.0	16.4	AHF 4
13	M	R	51.4	13.1	9.9	19.3	26.0	7.9	5.4	13.5	3.7	4.2	8.2	3.3	12.6	15.1	AHF 1
14	F	L	47.4	11.5	8.6	17.0	22.8	6.9	4.5	11.7	3.1	3.7	7.0	2.9	11.0	12.9	AHF 5
15																	
16																	
17																	
18																	
19																	
20																	
21																	
22																	
23																	
24																	
25																	
26																	
27																	
28																	

TABLE 6.
Cancer anthonyi, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
				1	F	L	63.7	15.3	12.9	23.8	32.0	9.1	6.3	16.4	4.0	4.8	
2	F	R	63.7	15.3	12.8	23.8	32.1	9.1	6.3	16.3	4.0	4.8	9.7	3.5	14.6	16.7	AHF 6
3	M	L	83.5	23.9	20.3	36.6	47.6	14.2	10.2	24.8	6.3	7.7	15.2	6.3	20.9	24.1	AHF 3
4	M	R	83.5	21.2	18.2	34.5	44.6	12.4	9.0	23.5	5.7	7.5	13.5	5.2	19.6	24.1	AHF 3
5	F	R	98.4	24.2	20.6	37.6	49.5	14.4	10.5	25.5	6.8	7.6	15.1	6.3	22.9	26.6	AHF 3
6	F	L	98.4	24.3	20.5	37.7	50.0	14.5	10.4	25.2	6.6	7.8	15.1	6.3	22.7	26.4	AHF 3
7	M	R	51.5	12.7	11.0	20.2	27.2	7.5	5.5	14.0	3.4	4.3	8.4	2.9	12.3	14.5	AHF 3
8	M	L	51.5	11.8	10.4	19.0	25.3	6.9	4.9	12.8	3.1	3.7	7.8	2.8	11.6	13.3	AHF 6
9	M	R	54.1	12.8	10.8	20.2	27.1	7.5	5.5	14.1	3.3	4.3	8.4	3.0	12.4	14.4	AHF 6
10	M	L	40.8	9.5	8.0	15.0	19.8	5.6	3.9	10.0	2.5	3.1	5.9	2.4	9.3	10.7	AHF 6

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.843	.847	.850	.858	.851	.843	.866	.882	.844	.842	.850	.750	.862	.860		.0299	3.53
T/H m	.595	.595	.594	.584	.595	.597	.590	.585	.586	.589	.588	.589	.603	.592		.0053	0.895
lf/H m	.314	.314	.322	.283	.314	.321	.339	.314	.336	.327	.315	.308	.328	.322		.0131	4.12
H/L mf	.397	.387	.412	.383	.412	.413	.393	.383	.390	.390	.354	.388	.391	.380		.0156	3.99
H/L f	.485	.495	.507	.556	.490	.517	.512	.474	.513	.526	.479	.480	.500	.500		.0219	4.37
T/H f	.728	.730	.818	.693	.852	.807	.683	.757	.697	.773	.783	.833	.763	.733		.0534	7.02
H/L c	.865	.874	.868	.813	.861	.860	.848	.873	.862	.869	.850	.873	.877	.864		.0163	1.89

TABLE 6.
Cancer anthonyi, statistical data (measurements in millimeters). (continued)

specimen no.	sex	Left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
				11	M	R	34.1	7.3	6.2	11.8	15.7	4.3	2.9	8.2	1.9	2.3	
12	M	L	34.1	7.8	6.6	12.3	16.3	4.6	3.3	8.5	2.0	2.4	5.0	2.0	7.6	8.7	AHF 6
13	M	L	49.6	11.6	10.0	18.7	25.0	7.0	5.0	12.8	2.9	3.8	7.6	2.9	11.4	13.0	AHF 6
14	F	L	39.5	9.3	8.0	14.7	19.6	5.5	3.8	10.3	2.5	3.0	6.0	2.2	8.9	10.3	AHF 6
15																	
16																	
17																	
18																	
19																	
20																	
21																	
22																	
23																	
24																	
25																	
26																	
27																	
28																	

TABLE 7.
Cancer branneri, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
				1	F	L	27.5	5.8	4.9	9.4	13.9	3.3	2.4	7.0	1.4	2.0	
2	F	R	33.3	7.0	6.0	11.6	16.7	4.0	3.0	8.4	1.9	2.4	5.4	1.7	7.9	7.7	AHF 8
3	M	R	24.1	5.1	4.3	8.3	11.7	3.0	2.1	5.9	1.3	1.8	3.7	1.1	5.6	5.6	AHF 9
4																	
5																	
6																	
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.844	.857	.843													.0081	0.957
T/H m	.568	.572	.588													.0106	1.84
Hf/H m	.345	.343	.360													.009	0.848
H/L mf	.343	.357	.356													.007	2.22
H/L f	.435	.443	.487													.028	6.15
T/H f	.650	.708	.612													.048	7.37
H/L c	.923	.975	1.000													.039	4.18

TABLE 8.
Cancer gracilis, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width														
				Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
1	F	R		12.6	10.0	19.5	26.7	6.9	5.4	13.0	3.2	4.3	8.5	2.9	11.5	12.2	AHF 7
2	M	R	84.3	23.4	18.0	47.6	48.0	12.4	9.4	22.7	5.7	3.4	6.6	5.0	21.1	20.5	AHF 2
3	M	L	62.5	16.8	13.6	25.8	36.0	9.3	7.1	17.7	4.4	5.8	12.3	3.8	15.8	15.9	AHF 10
4	F	L	53.4	12.9	10.2	19.7	27.7	6.8	5.3	13.9	3.1	4.4	9.4	2.9	12.1	13.4	AHF 10
5	F	R	55.0	13.9	11.0	20.8	29.3	7.2	5.7	14.8	3.2	4.9	9.8	3.0	13.0	13.1	AHF 10
6	F	R	55.5	13.3	11.0	21.0	29.0	7.1	5.6	14.8	3.2	5.0	9.8	3.0	13.5	13.4	AHF 10
7	F	L	67.7	16.1	12.5	24.6	33.5	8.8	6.6	16.2	4.0	5.0	11.0	3.6	15.0	15.2	AHF 11
8	F	L	57.6	13.2	10.4	20.5	29.0	7.3	5.4	14.5	3.3	4.5	10.0	3.1	13.0	12.7	AHF 11
9	F	R	43.2	10.1	8.0	15.6	22.3	5.5	4.3	11.0	2.5	3.5	7.4	2.3	9.9	9.8	AHF 11
10	F.	L	46.1	10.7	8.8	16.7	23.4	6.0	4.3	11.7	2.6	3.8	8.0	2.3	10.6	10.4	AHF 11

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
T _{lu} /H m	.793	.770	.810	.792	.792	.827	.777	.788	.792	.822						.0183	2.30
T/H m	.548	.530	.553	.527	.518	.533	.547	.553	.545	.561						.0141	2.60
H _f /H m	.333	.316	.345	.341	.353	.376	.329	.341	.347	.355						.0163	4.75
H/L m f	.408	.414	.402	.381	.385	.378	.407	.373	.391	.368						.0162	4.16
H/L f	.493	.515	.472	.468	.500	.510	.482	.450	.473	.475						.0229	4.74
T/H f	.691	.676	.655	.659	.612	.600	.679	.690	.657	.605						.0348	5.33
H/L c	.943	.952	.993	.903	1.010	.993	.987	.977	.990	.981						.0316	3.25

TABLE 9.
Cancer jordani, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width														Locality
				Manus					Movable finger			Fixed finger			Carpus		
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
1	M	L	27.4	7.3	5.6	10.5	14.8	4.3	2.8	7.3	2.0	2.3	4.5	1.7	6.9	7.4	Loc. unknown
2	M	R	38.2	11.8	9.6	17.0	22.9	7.3	4.3	11.4	3.2	3.7	6.9	2.7	10.5	11.6	AHF 12
3	M	L	26.1	7.0	5.5	10.2	14.3	4.0	2.7	7.1	1.9	2.2	4.6	1.6	6.4	7.1	AHF 13
4	M	R	27.9	7.6	5.6	11.0	15.0	4.4	3.0	7.4	2.0	2.4	4.6	1.9	7.2	7.8	AHF 14
5																	
6																	
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.767	.813	.786	.737												.0321	4.14
T/H m	.589	.618	.572	.578												.0204	3.46
Hf/H m	.315	.314	.314	.316												.0010	0.319
H/L m f	.384	.377	.380	.405												.0127	3.29
H/L f	.418	.537	.478	.522												.0534	10.9
T/H f	.739	.730	.727	.792												.030	4.07
H/L c	.933	.907	.903	.923												.0140	1.53

TABLE 10.
Cancer magister, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
				1	F	R	88.0	15.9	17.0	28.4	38.5	9.0	7.0	19.9	4.9	5.6	
2	F	L	101.3	18.7	18.5	32.1	44.4	10.4	8.1	23.0	5.6	7.0	15.8	5.3	20.3	21.9	AHF 18
3	F	R	101.3	18.7	18.3	32.5	44.7	10.5	8.3	23.0	5.6	6.8	15.7	5.3	19.9	21.8	AHF 18
4	M	R	83.2	14.9	15.5	25.9	34.6	8.8	6.4	18.1	4.4	5.4	12.6	4.1	16.5	17.7	AHF 18
5	M	L	83.2	15.0	15.4	26.4	35.0	8.6	6.4	18.2	4.5	5.6	12.7	4.1	16.6	17.7	AHF 18
6	M	L	75.1	12.9	13.5	23.0	31.7	7.3	5.7	16.3	3.9	4.6	11.4	3.4	14.6	15.8	AHF 18
7	M	R	76.4	13.4	13.7	23.4	31.8	7.7	5.6	16.3	4.0	5.0	11.2	3.3	14.6	15.7	AHF 18
8	M	L	76.4	13.3	13.8	23.8	32.2	7.7	5.7	16.5	4.0	5.0	11.1	3.3	14.6	16.0	AHF 18
9	M	R	110.0	19.7	20.6	34.8	48.3	11.7	8.7	24.6	6.0	7.4	17.2	5.3	21.7	22.9	AHF 19
10	M	L	110.0	19.6	20.9	34.8	48.2	11.8	8.8	24.6	6.0	7.5	17.2	5.7	21.7	23.2	AHF 19
11	F	R	75.4	13.3	13.5	23.3	32.1	7.3	5.6	16.5	3.7	4.8	11.3	3.4	14.6	16.0	AHF 20
12	F	L	75.4	13.4	13.3	23.5	32.3	7.3	5.9	16.7	3.9	5.0	11.4	3.6	14.6	16.4	AHF 20
13	F	R	41.7	7.5	7.6	12.4	17.4	4.5	3.3	9.0	2.2	2.7	6.4	1.9	8.5	8.9	AHF 20
14	F	L	41.7	7.4	7.5	12.7	17.8	4.5	3.3	9.0	2.2	2.6	6.4	1.8	8.5	8.9	AHF 20

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	1.070	.990	.978	1.040	1.025	1.046	1.022	1.037	1.045	1.065	1.015	.993	1.012	1.012		.027	2.63
T/H m	.567	.557	.562	.590	.573	.566	.574	.579	.594	.602	.548	.545	.600	.608		.0202	3.51
Hf/H m	.353	.374	.364	.362	.374	.357	.373	.376	.375	.383	.361	.373	.360	.351		.0097	2.66
H/L mf	.351	.352	.361	.354	.352	.350	.344	.346	.354	.358	.340	.353	.367	.367		.0078	2.34
H/L f	.415	.443	.433	.428	.441	.403	.446	.450	.431	.437	.466	.481	.423	.406		.0215	4.95
T/H f	.768	.757	.779	.758	.732	.739	.660	.660	.717	.760	.708	.720	.703	.693		.0379	5.23
H/L c	.931	.927	.913	.932	.938	.923	.930	.913	.956	.936	.913	.890	.978	.978		.0245	2.63

TABLE II.
Cancer oregonensis, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	L1	T	H	L	T	H	L	T	H	L	
				1	F	R	24.8	8.0	4.9	10.0	12.9	4.6	2.8	7.3	2.2	2.6	
2	F	L	33.2	10.9	6.8	13.5	16.4	6.1	4.0	8.8	3.1	3.5	4.5	2.4	9.1	11.8	AHF 15
3	F	R	33.2	10.8	6.8	13.5	16.2	6.1	3.9	8.5	3.0	3.3	4.5	2.6	9.1	11.9	AHF 15
4	M	L	21.7	8.3	5.1	10.0	12.8	4.8	3.0	7.3	2.4	2.9	3.9	1.9	7.3	8.8	AHF 15
5	M	R	21.7	8.5	5.1	10.1	13.0	4.7	3.0	7.2	2.3	3.0	3.9	2.0	7.3	8.7	AHF 15
6	F	L	30.1	10.1	6.1	12.1	16.1	5.7	3.7	9.3	2.8	3.3	5.3	2.4	8.7	10.7	AHF 15
7	F	R	24.6	8.0	4.9	10.0	13.2	4.6	2.7	7.3	2.3	2.7	4.2	2.0	6.9	8.7	AHF 15
8	F	L	33.2	11.5	7.0	13.9	13.1	6.4	4.0	9.5	3.3	3.7	5.2	2.8	9.6	11.9	AHF 15
9	F	R	27.7	9.4	5.3	11.3	14.8	5.4	3.2	8.1	2.3	3.1	4.6	2.0	8.1	10.0	AHF 15
10	F	L	25.6	8.5	5.3	10.8	14.1	4.9	2.9	7.5	2.3	3.0	4.3	2.2	7.3	9.4	AHF 15

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.613	.623	.629	.615	.600	.603	.613	.608	.653	.612						.0179	2.94
T/H m	.575	.559	.564	.578	.553	.565	.575	.557	.574	.577						.0092	1.63
Hf/H m	.325	.321	.305	.349	.353	.327	.338	.322	.330	.353						.0158	4.77
H/L mf	.384	.454	.458	.412	.417	.398	.370	.422	.395	.387						.0290	7.09
H/L f	.650	.777	.733	.743	.769	.623	.643	.712	.674	.698						.0538	7.67
T/H f	.769	.686	.788	.655	.770	.728	.740	.756	.645	.735						.0494	6.79
H/L c	.828	.771	.765	.829	.839	.813	.793	.807	.810	.777						.0258	3.22

TABLE 12.
Cancer productus, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
				1	?	R		21.8	18.4	32.7	42.5	12.2	9.4	21.5	5.5	7.0	
2	?	R		31.5	28.4	48.4	62.2	18.5	13.8	31.2	8.6	10.5	17.7	7.9	28.4	31.1	Loc. unknown
3	?	L		38.9	34.3	57.4	72.5	21.7	17.1	38.2	11.0	13.0	21.1	10.1	34.3	37.5	Loc. unknown
4	M	R	69.1	15.5	14.8	25.0	32.0	8.5	6.8	15.9	3.9	5.1	8.9	3.6	14.9	17.0	AHF 18
5	M	L	88.8	19.7	18.9	32.4	41.0	11.4	8.5	20.5	4.9	6.2	11.5	4.8	19.2	21.9	AHF 18
6	M	R	88.8	19.6	18.8	32.2	41.3	11.4	8.6	20.4	4.9	6.4	11.4	4.7	19.2	21.7	AHF 18
7	F	L	66.2	13.3	12.7	21.8	28.4	7.5	5.5	14.5	3.3	4.3	8.3	3.2	12.7	15.2	AHF 18
8	F	R	66.2	13.3	12.7	22.1	28.9	7.8	5.6	14.2	3.2	4.3	8.4	3.2	13.2	15.2	AHF 18
9	F	L	43.7	9.4	8.8	15.1	19.1	5.2	4.0	9.6	2.3	3.1	5.2	2.2	9.2	10.4	AHF 18
10	F	R	43.7	9.3	8.8	14.9	19.0	5.1	4.1	9.6	2.3	3.0	5.2	2.2	9.2	10.5	AHF 18
11	M	L	50.9	11.3	10.1	18.0	23.2	6.5	4.8	11.6	2.8	3.7	6.6	2.7	11.1	12.6	AHF 18
12	M	R	59.0	13.4	12.0	20.9	26.7	7.5	5.7	13.5	3.3	4.3	7.5	3.1	12.7	14.3	AHF 18
13	F	L	82.0	17.9	16.5	28.3	36.5	9.7	7.6	18.3	4.6	5.9	10.1	4.1	16.8	19.1	AHF 18
14	F	R	38.9	8.1	7.3	13.2	16.7	4.7	3.4	8.2	2.0	2.6	4.6	1.9	7.9	9.3	AHF 18
15	F	L	38.9	5.5	4.9	8.7	12.2	3.3	2.3	6.5	1.6	2.0	3.8	1.6	5.4	6.7	AHF 18

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.844	.902	.882	.955	.960	.960	.954	.954	.936	.947	.893	.896	.922	.902	.892	.0356	3.87
T/H m	.560	.587	.558	.548	.578	.582	.563	.587	.553	.548	.575	.559	.542	.580	.600	.0174	3.06
Hf/H m	.321	.334	.334	.329	.315	.326	.323	.323	.330	.323	.328	.321	.330	.321	.364	.0113	3.40
H/L mf	.437	.446	.460	.428	.414	.422	.380	.394	.417	.427	.413	.423	.415	.414	.354	.0256	6.18
H/L f	.583	.593	.616	.580	.538	.562	.518	.512	.596	.577	.562	.573	.583	.565	.527	.030	5.32
T/H f	.728	.753	.787	.707	.773	.773	.744	.744	.710	.733	.730	.721	.695	.731	.800	.0291	3.94
H/L c	.882	.913	.915	.876	.877	.885	.835	.866	.885	.876	.877	.888	.880	.850	.806	.0274	3.14

TABLE 13.
Cancer davidi, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
				1		L	47.0	11.7	8.8	17.2		6.4	5.1		2.9		
2																	
3												3.2	6.9	2.2			LACMIP 463
4									5.0	10.5							USGS M1611
5																	
6																	
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.752																
T/H m	.547																
Hf/H m																	
H/L mf				.476													
H/L f			.463														
T/H f			.687														
H/L c	.833																

TABLE 14.
Cancer fissus, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
				1	F	L	71.9	14.2	13.7	23.2	27.5	8.5				5.6	
2	F	R	71.9			24.1		8.3				5.6		3.4	13.6		UCMP D-362
3																	
4																	
5																	
6																	
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.965																
T/H m	.598																
Hf/H m	.394																
H/L mf																	
H/L f	.651																
T/H f	.582	.607															
H/L c	.773																

TABLE 15.
Cancer granti, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width	Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
				1			56.0	16.3	13.2		31.0		6.2	17.0	4.7	5.7	
2																	
3																	
4																	
5																	
6																	
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.810																
T/H m																	
Hf/H m	.350																
H/L mf	.365																
H/L f	.570																
T/H f	.614																
H/L c	.933																

TABLE 16.
Cancer allisoni, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width														
				Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
1		L						8.9	20.0	5.0	6.5	11.4	4.8			LACMIP 499	
2		L						5.3	13.2	3.4						LACMIP 456	
3		R						3.3	7.8	2.1						LACMIP 456	
4		R						3.5	8.6	2.2						LACMIP 463	
5		R						2.5	5.9	1.6						LACMIP 463	
6		R						3.4	8.7	2.3						USNM 18284	
7		L						2.9	7.2	2.0						USNM 18284	
8		L									4.0	6.7	2.6			LACMIP 462	
9		L									4.0	6.7	2.8			LACMIP 466	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m																	
T/H m																	
Hf/H m																	
H/L mf	.445	.402	.423	.407	.423	.391	.403										
H/L f	.570							.597	.597								
T/H f	.728							.650	.700								
H/L c																	

TABLE 17.
Cancer marri, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width													Locality	
				Manus					Movable finger			Fixed finger			Carpus		
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H		L
1				9.6	8.1	15.7		5.3	4.4		2.4	3.8		2.3	10.0	9.6	UCMP D-3402
2				12.2	10.3			6.7				4.7	7.7	2.9			UCMP D-3402
3															22.6	23.5	UCMP D-3402
4		L							11.3	27.4	6.5						UCMP D-2506
5		R							7.8	17.7	3.7						UCMP D-3389
6				14.5	12.8			6.5									UCMP D-3403
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.843	.843				.623											
T/H m	.552	.548				.448											
Hf/H m	.396	.385															
H/L mf				.412	.441												
H/L f		.610															
T/H f	.606	.617															
H/L c	.960		.962														

TABLE 18.
Cancer yanceyi, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width														Locality
				Manus					Movable finger			Fixed finger			Carpus		
				H	Lu	Lm	L1	T	H	L	T	H	L	T	H	L	
1		L		27.7	20.0			13.3				7.2		6.3			UCMP B-7727
2		R		22.6	15.6	30.2		11.9				6.8		4.5			UCMP B-7727
3																	
4																	
5																	
6																	
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	.723	.690															
T/H m	.481	.527															
Hf/H m	.260	.301															
H/L mf																	
H/L f																	
T/H f	.875	.662															
H/L c																	

TABLE 19.
Cancer durhami, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width														
				Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
1		R		21.3	21.4	39.1	54.6	10.4	9.3	27.1	5.3	8.0	17.8	5.0	22.8	26.9	UCMP A-9733
2																	
3																	
4																	
5																	
6																	
7																	
8																	
9																	
10																	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m	1.005																
T/H m	.488																
Hf/H m	.376																
H/L mF	.343																
H/L f	.448																
T/H f	.625																
H/L c	.848																

TABLE 20.
Cancer coosensis, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width															
				Manus					Movable finger			Fixed finger			Carpus		Locality	
				H	Lu	Lm	L1	T	H	L	T	H	L	T	H	L		
1		R							11.1	33.2	7.0							USNM 18284
2		R		18.1	16.5	25.9		6.6					7.0		3.8			USNM 18284
3		L							10.5	31.7	6.9							USNM 18284
4		R		27.9	22.5	32.9							9.6		5.1			USNM 18284
5																		
6																		
7																		
8																		
9																		
10																		

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m		.913		.822													
T/H m																	
Hf/H m		.386		.344													
H/L mf	.334		.331														
H/L f																	
T/H f		.543		.531													
H/L c																	

TABLE 21.
Cancer chaneyi, statistical data (measurements in millimeters).

specimen no.	sex	left or right	carapace width														
				Manus					Movable finger			Fixed finger			Carpus		Locality
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
1		R						6.4	16.9	3.9						USNM 496181	
2		R						5.5	14.1	3.2						USNM 496181	
3		L	22.0					8.7	23.2	5.0	7.3	13.8	4.9			LACMIP 130-7	
4		L						2.0	5.6	1.0						LACMIP 130-7	
5		R						5.6	14.0	3.4						LACMIP 130-7	
6		R						4.3	10.4	2.3						LACMIP 130-7	
7		R						3.1	8.0	1.6						LACMIP 130-7	
8								2.7	7.2	1.7						LACMIP 130-7	
9		R									6.0	11.5	4.4			LACMIP 130-7	
10		R									6.3	12.6	4.5			LACMIP 130-7	

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m																	
T/H m																	
Hf/H m																	
H/L mf	.378	.390	.375	.357	.400	.413	.388	.375									
H/L f			.529						.522	.500							
T/H f			.671						.733	.714							
H/L c																	

TABLE 21.
Cancer chaneyi, statistical data (measurements in millimeters). (continued)

specimen no.	sex	left or right	carapace width															
				Manus					Movable finger			Fixed finger			Carpus		Locality	
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L		
1		L										2.5	5.1	1.6			LACMIP 466	
2		L										4.5	8.6	3.3			UCLA L-2716	
3		R										5.9	11.0	3.7			LACMIP 466	
4		R										3.7	7.3	2.4			LACMIP 466	
5		L										6.0	11.1	4.1			LACMIP 466	
6		R		19.5	18.1	30.1	39.5	10.8				6.3	11.7	4.5	17.1	18.6	USNM 353394	
7		L										6.4	12.8	3.9			USNM 324300	
8																		
9																		
10																		

RATIOS

spec. no.	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	S.	V.
Lu/H m						.928											
T/H m						.553											
Hf/H m						.323											
H/L mf																	
H/L f	.490	1.523	.537	.507	.540	.538	.500										
T/H f	.640	.733	.627	.648	.683	.714	.608										
H/L c						.919											

TABLE 21.
Cancer chaneyi, statistical data (measurements in millimeters) (continued)

specimen no.	sex	left or right	carapace width														Locality
				Manus					Movable finger			Fixed finger			Carpus		
				H	Lu	Lm	Ll	T	H	L	T	H	L	T	H	L	
1		R										2.2	4.1	1.6			LACMIP 130-7
2		R								2.1	5.0	1.0					LACMIP 130-4
3		R										5.3	10.1	3.3			LACMIP 130-4
4		R										2.8	5.4	1.8			LACMIP 130-4
5												4.5	8.7	3.4			LACMIP 130-4
6		R										6.0	11.5	4.3			LACMIP 463
7		R								2.4	6.2	1.4					LACMIP 466
8		R								2.5	7.0	1.4					LACMIP 466
9		R								3.0	6.7	1.4					LACMIP 466
10		L										2.4	4.4	1.5			LACMIP 466

RATIOS

spec. no.	11	12	13	14	15	16	17	18	19	20							
Lu/H m																	
T/H m																	
Hf/H m																	
H/L mf		.420					.387	.357	.448								
H/L f	.537		.525	.518	.517	.522				.545							
T/H f	.727		.623	.643	.755	.717				.625							
H/L c																	

LITERATURE CITED

- ARNOLD, RALPH. 1903. The paleontology and stratigraphy of the marine Pliocene and Pleistocene of San Pedro, California. Calif. Acad. Sci., Mem. 3:1-420, 37 pl.
- . 1909. Paleontology of the Coalinga District, Fresno and Kings Counties, California. U.S. Geol. Survey, Bull. 396:1-173, 30 pls.
- BALSS, H. 1922. Ostasiatische Decapoden. IV. Die Brachyrhynchen (Cancridae). Archiv für Naturgeschichte, Abt. A, Heft. 11:95-166.
- BELL, THOMAS. 1853. A history of the British Stalk-eyed Crustacea. Van Voorst, London, 386 pp.
- BENNETT, E. W. 1964. The marine fauna of New Zealand: Crustacea Brachyura. New Zealand Oceanographic Inst. Mem. 22:1-120.
- BUTLER, T. H. 1967. A bibliography of the Dungeness crab, *Cancer magister* Dana. Fisheries Research Board of Canada, Tech. Rept. 1.
- CAIN, STANLEY A. 1944. Foundations of Plant Geography. Harper and Bros., New York.
- CALIFORNIA, STATE OF. 1958. Geologic Map of California, Santa Cruz Sheet, Dept. Nat. Res., Div. of Mines, 1:250,000.
- CALMAN, W. T. 1911. The life of Crustacea, 289 p., 32 pls., 85 text-figs., Methuen Co., Ltd., London.
- CHILTON, C., AND E. W. BENNETT. 1929. Contributions for a revision of the Crustacea Brachyura of New Zealand. Proc. New Zealand Inst., Trans. 59(4):731-778.
- CHRISTIANSEN, MARIT E. 1969. Marine invertebrates of Scandinavia. No. 2, Decapoda Brachyura. Universitets Forlaget, Oslo, pp 1-143.
- CUMMINGS, F. C., R. M. TOURING AND E. E. BRABB. 1962. Geology of the northern Santa Cruz Mountains, California. Calif. Div. Mines and Geology, Bull. 181:179-220, 5 pls.
- DURHAM, J. W. 1950. Cenozoic marine climates of the Pacific Coast. Geol. Soc. America, Bull. 61:1243-1264, figs. 1-3.
- DURHAM, J. W., R. H. JAHNS AND D. E. SAVAGE. 1954. Marine-nonmarine relationships in the Cenozoic section of California. In: Geology of southern California, Calif. Div. Mines, Bull. 170(1):59-71.
- EKMANN, S. 1953. Zoogeography of the sea. Sidgwick and Jackson, Ltd., London, 417 pp.
- GABB, W. M. 1869. Cretaceous and Tertiary fossils. Calif. Geol. Survey, Paleontology 2:1-299.
- GARTH, J. S. 1957. Reports of the Lund University Chile Exped. 1948-49, No. 29. The Crustacea Decapoda Brachyura of Chile. Lunds Universitets Arsskrift, n.f., Avd. 2, Bd. 53, Nr. 7:1-131.
- . 1960 (1961). Distribution and affinities of the Brachyuran Crustacea. In: The Biogeography of Baja California and Adjacent Seas (A Symposium). Systematic Zoology 9:105-123.
- GISTL, J. VON N.F.X. 1848. Naturgeschichte des Thierreichs für Höhere Schulen. Hoffman'sche Verlags-Buchhandlung, Stuttgart, 216 pp.
- GLAESSNER, M. F. 1929. Fossilium Catalogus. 1: Animalia, 41: Crustacea Decapoda, 101-109 pp.
- . 1960. The fossil decapod Crustacea of New Zealand and the evolution of the Order Decapoda. New Zealand Geol. Survey, Paleont. Bull. 31:1-79.
- . 1969. Decapoda. In: Moore 1969. Treatise on Invertebrate Paleontology, Part R, Arthropoda 4, 2:651.
- GLEN, WILLIAM. 1959. Pliocene and Lower Pleistocene of the Western part of the San Francisco Peninsula, Univ. of Calif., Publ. in Geol. Sci. 36(2):147-198, pl. 15-17, 5 text-figs.
- GRANT, U. S., IV, AND LEO G. HERTLEIN. 1938. The West American Cenozoic Echinoidea. Univ. California Los Angeles, Pubs. in Math. and Phys. Sci. 2:1-225, 30 pls.
- HELLER, C. 1863. Die Crustaceen des sublichen Europa, Crustacea. Podothalmia. Wilhelm Braumüller, Vienna, 336 pp.
- HOLLAND, R. D., JR., AND A. M. CVANCARA. 1958. Crabs from the Cannonball Formation (Paleocene) of North Dakota. Jour. Paleont. 32(3):495-505, 3 text-figs., pl. 74.
- HOOTS, H. W. 1930. Geology of the eastern part of the Santa Monica Mountains, Los Angeles County, California. U.S. Geol. Survey, Prof. Paper 165-C:1-134, 19 pls., 2 figs.
- HOLTHUIS, L. B. 1949. Fossil decapod Crustacea from Miocene and younger deposits of the Netherlands. Med. Geol. stichting, n.s., 3:55-68.
- IMAIZUMI, R. 1962. Miocene *Cancer* of Japan. Sci. Repts. Tohoku Univ., Ser. 2 (Geology), 5:233-245.
- JOHNSON, J. Y. 1861. Description of a new species of *Cancer* obtained at Madeira. Proc. Zool. Soc. London, 16:240-242.
- JOHNSON, M. E., AND H. J. SNOOK. 1927. Seashore animals of the Pacific Coast. MacMillan, New York, 659 p., 498 color pls., 700 figs.
- KEROHER, G. C. 1966. Lexicon of geologic names of the United States for 1936-1960. U.S. Geol. Survey, Bull. 1200.
- LAWSON, A. C. 1894. The geomorphogeny of the Coast of Northern California. Univ. of Calif., Dept. Geology, Bull. 1(8):241-272.
- LINNAEUS, CAROLUS. 1758. Systema naturae. Tenth edition, 1, 824 pp.
- LUTZ, G. C. 1951. The Sobrante Sandstone. Univ. Calif. Pubs., Dept. Geol. Sci., Bull. 28(13):367-406.
- MAC KAY, D. C. G. 1943. Temperature and the world distribution of crabs of the genus *Cancer*. Ecology 24(1):113-115.
- . 1943. Relative growth of the European edible crab, *Cancer pagurus*. III. Growth of the sternum and appendages. Growth 7(4):401-412, 3 figs., 2 tables.
- MENZIES, R. J. 1948. A revision of the brachyuran genus *Lophonanopeus*. Allen Hancock Foundation, Occ. Pap. 4:1-45.
- . 1951. Pleistocene Brachyura from the Los Angeles area: Cancridae. Jour. Paleont. 25(2):165-170.
- MIERS, E. J. 1879. On a collection of Crustacea made by Capt. H. D. St. John, R. N., in the Korean and Japanese Seas. Part I. Podophthalmia. Proc. Zool. Soc. London, 18-61.

- MILNE-EDWARDS, A. 1865. Etudes Zoologiques sur les Crustacés Recents de la Famille des Cancériens. *Nouv. Arch. Mus. Hist. Nat.*, 1:177-308.
- . 1865. Histoire des Crustacés Podophthalmaires fossiles, Tome 1, p. 1-390, pls. 1-36, Masson, Paris.
- NATIONS, J. D. 1968. A new species of cancroïd crab from the Pliocene of California. *Jour. Paleont.* 42(1):33-36.
- NOETLING, F. 1910. The fauna of the Miocene beds of Burma. *Geol. Surv. India, Paleont. Indica, N.S., Mem.* 1(3):1-378, pl. 1-25.
- PESTA, OTTO. 1918. Die Decapodenfauna der Adria. *Franz Deuticke, Leipzig and Vienna*, 500 pp.
- PHILIPPI, R. A. 1887. Die Tertiären und quartären versteinerten Chiles. *Leipzig, F. A. Brockhaus*, 266 p., 58 tables.
- RATHBUN, M. J. 1908. Fossil crabs from California. *U.S. Nat. Mus., Proc.* 35:341-349, pls. 45-49.
- . 1917. Description of a new species of crab from the California Pliocene. *U.S. Nat. Mus., Proc.* 53:451-452, pl. 59.
- . 1918. Decapod crustaceans from the Panama Region. *U.S. Nat. Mus., Bull.* 103:123-184, pls. 54-66.
- . 1919. West Indian Tertiary decapod crustaceans. *Carnegie Inst., Pub.* 291:157-184, pls. 1-9.
- . 1920. Additions to West Indian Tertiary decapod crustaceans. *U.S. Nat. Mus., Proc.* 58:381-384, 1 pl.
- . 1923a. Decapod crustaceans from the Upper Cretaceous of North Carolina. *North Carolina Geol. and Econ. Surv.* 5:403-408, pls. 101-102.
- . 1923b. Fossil crabs from the Republic of Haiti. *U.S. Nat. Mus., Proc.* 63(9):1-6, pls. 1-2.
- . 1926. The fossil stalk-eyed Crustacea of the Pacific Slope of North America. *U.S. Nat. Mus., Bull.* 138:1-155.
- . 1928. Two new crabs from the Eocene of Texas. *U.S. Nat. Mus., Proc.* 73(6):1-6, pls. 1-3.
- . 1929. A new crab from the Eocene of Florida. *U.S. Nat. Mus., Proc.* 75:(15):1-4, pls. 1-3.
- . Fossil decapod crustaceans from Mexico. *U.S. Nat. Mus., Proc.* 78(8):1-10, pls. 1-6.
- . 1930a. The cancroïd crabs of America of the Families Euryalidae, Portunidae, Atelecyclidae, Cancridae and Xanthidae. *U.S. Nat. Mus., Bull.* 152:1-609.
- . 1930b. A correction. *Science* 72:279.
- . 1932. A new species of *Cancer* from the Pliocene of the Los Angeles Basin. *Wash. Acad. Sci., Jour.* 22(1):19-20.
- . 1935. Fossil Crustacea of the Atlantic and Gulf Coastal Plain. *Geol. Soc. America, Spec. Paper* 2:1-160.
- SAKAI, TUNE. 1939. Studies on the crabs of Japan, IV. Brachygnatha, Brachyrhyncha, 741 pp.
- . 1965. The crabs of Sagami Bay, collected by His Majesty the Emperor of Japan. *Maruzen Co., Ltd., Tokyo*, 206 pp.
- . 1969. Two new genera and twenty-two new species of crabs from Japan. *Proc. Biol. Soc. Washington*, 82:243-280.
- SCHMITT, W. L. 1921. The marine decapod Crustacea of California. *Univ. Calif., Publ. Zool.* 23:1-470 p., 50 pls., 165 text-figs.
- . 1965. Crustaceans. *Univ. Mich. Press*, 204 p., 75 figs.
- SIMPSON, G. G. 1941. Range as a zoological character. *Amer. Jour. Sci.* 239(11):785-804, 2 tables.
- SIMPSON, G. G., ANNE ROE AND R. C. LEWONTIN. 1960. *Quantitative Zoology*. Harcourt, Brace and World, Inc., New York, 440 pp.
- SOPER, E. K., AND U. S. GRANT, IV. 1932. Geology and Paleontology of a portion of Los Angeles, California. *Geol. Soc. America, Bull.* 43:1041-1068, 7 figs.
- STENZEL, H. B. 1944. Decapod crustaceans from the Cretaceous of Texas. *Univ. Texas Publ., Contrib. to Geol.* 4401:401-453, 16 text-figs., 12 pls.
- . 1952. Decapod crustaceans from the Woodbine Formation of Texas. *Jour. Paleont.* 8(1):38-56, pls. 6 and 7.
- SVERDRUP, H. U., M. W. JOHNSON AND R. H. FLEMING. 1942. The oceans, their physics, chemistry and general biology. *Prentice-Hall, New Jersey*, 1087 pp.
- TAKEDA, M. AND S. MIYAKE. 1972. New crabs from the sea around the Tsushima Islands. *Nat. Sci. Mus., Tokyo, Bull.* 15(2):254.
- VAN COUVERING, M., AND H. B. ALLEN. 1943. Devils Den Oil Field. *Calif. Div. Mines, Bull.* 118:496-504, figs. 211-213, 1 table.
- VAN STRAELEN, V. 1938. Crustacés Decapodes Cénozoïques des Indes Orientales Néerlandaises. *Leidsche Geol. Meded., Leyden*, 10:90-103.
- . 1939. *Martinezicancer schencki*, Brachyüre nouveau du Pliocène de Californie. *Mus. Royal Hist. Nat. Belg., Bull.* 15(63): 1-3, figs. a, b.
- WEAVER, C. E. 1944. Correlation of Cenozoic formations of western North America. *Geol. Soc. America, Bull.* 55:569-598.
- WEYMOUTH, F. W. 1910. Synopsis of the true crabs (Brachyura) of Monterey Bay, California. *Leland Stanford Junior Univ. Pub., Univ. Ser.* 4:1-64, 14 pls., 9 text-figs.
- WEYMOUTH, F. W., AND D. C. G. MACKAY. 1936. Analysis of the relative growth of the Pacific edible crab, *Cancer magister*. *Proc. Zool. Soc., London*, pt. 1:257-280, 10 text-figs., 10 tables.
- WILLETT, GEORGE. 1937. An Upper Pleistocene Fauna from the Baldwin Hills, Los Angeles County, California. *San Diego Soc. Nat. Hist., Trans.* 8(30):379-406, pls. 25, 26.
- WILSON, IVAN F. 1943. Geology of the San Benito Quadrangle, California. *Calif. Jour. Mines and Geol.*, 39(2):183-270.
- WOOLWARD, HENRY. 1896. On some podophthalmous Crustacea from the Cretaceous Formation of Vancouver and Queen Charlotte Islands. *Quart. Jour. Geol. Soc. London*, 52:221-228.
- ZULLO, V. A. 1969a. Thoracic cirripedia of the San Diego Formation, San Diego County, California. *Los Angeles Co. Mus., Contrib. Sci.* 159:1-25, figs. 1-77.
- . 1969b. A Late Pleistocene marine invertebrate fauna from Bandon, Oregon. *Calif. Acad. Sci., 4th Ser., Proc.* 36(12):347-361.