

- Strombidae and Aporrhaidae. *American Journal of Conchology* 4:137–149, pls. 13, 14.
- GARDNER, J. S. 1884. British Eocene Aporrhaidae. *Geological Magazine, New Series, Decade 3*, 1:529–534, pl. 17.
- GITTON, J. L., P. LOZOUET & P. MAESTRATI. 1986. Biostratigraphie et paléocologie des gisements types du stampien de la région d'Etampes (Essonne). France, Editions du Bureau des Recherches Géologiques et Minières, *Géologie de la France* no. 1–1986, 101 pp., 4 pls.
- GLIBERT, M. 1957. Pélécytopes et gastropodes du Rupélien supérieur et du Chattien de la Belgique. Institut royal des sciences naturelles de Belgique, *Mémoire* 137, 98 pp.
- GRAY, J. E. 1850. Systematic arrangement of the figures. Pp. 63–124 in M. E. Gray, 1850, *Figures of Molluscos Animals Selected from Various Authors; Etched for the Use of Students, Vol. 4*. Longman, Brown, etc: London, 219 pp.
- GRÜNDEL, J. 1997. Die Gastropodenfauna des Rupels von Amsdorf (Westlich Halle). *Palaeontographica, Abteilung A, Paläozoologie—Stratigraphie*: 1–36, 6 pls.
- HALLER, B. 1888. Die Morphologie der Prosobranchier. *Morphologisches Jahrbuch* 14: 54–169, pl. 3–9.
- HARRIS, G. D. 1899. The Lignitic Stage. Part II. Scaphopoda, Gastropoda, Pteropoda, and Cephalopoda. *Bulletins of American Paleontology* 3(11):1–128, pls. 1–12. [Reprinted by Kraus Reprint, 1964]
- KOLLMANN, H. A. & J. S. PEEL. 1983. Paleocene gastropods from Nigssuaq, West Greenland. København, Grønlands Geologiske Undersøgelse, *Bulletin* 146, 115 pp., 257 figs.
- LINNAEUS, C. 1758. *Systema naturae per regna tria naturae. Editio decima, reformata*. Stockholm, Vol. 1, *Regnum animale*. 824 pp.
- LOZOUET, P. 1986. Les gastropodes prosobranches de l'Oligocene Supérieur du bassin de l'Adour (Systematique, paléoenvironnements, paléoclimatologie, paléobiogéographie). Thèse Mémoire de l'école Pratique des Hautes Études (Paris), 475 pages, 34 pls.
- MACNEIL, F. S. & D. T. DOCKERY, III. 1984. Lower Oligocene Gastropoda, Scaphopoda, and Cephalopoda of the Vicksburg Group in Mississippi. *Mississippi Bureau of Geology Bulletin*, 124, 415 pp., 72 pls.
- MANTELL, G. 1829. A tabular arrangement of the organic remains of the County of Sussex. *Geological Society of London Transaction* (2)3:201–216.
- MEEK, F. B. 1864. Check-list of the invertebrate fossils of North America; Cretaceous and Jurassic. *Smithsonian Miscellaneous Collections* 7, no. 8 (Publ. 177), 40 pp.
- MEEK, F. B. & F. V. HAYDEN. 1856. Description of new species of Gastropoda from the Cretaceous formations of Nebraska Terr. *Academy of Natural Sciences of Philadelphia, Proceedings for 1856* 9:117–148.
- MICHAUD, A. G. L. 1828. Description de plusieurs espèces de coquilles vivantes de la Méditerranée. *Bulletin d'Histoire Naturelle de la Société Linnéenne de Bordeaux* 2:119–122.
- PERRON, F. E. 1978. Seasonal burrowing behavior and ecology of *Aporrhais occidentalis* (Gastropoda, Strombacea). *Biological Bulletin* 154:463–471.
- PONDER, W. F. & A. WARÉN. 1988. Classification of the Caenogastropoda and Heterostropha—a list of the family-group names and higher taxa. *Malacological Review* 4(Suppl.): 288–326.
- POPENOE, W. P. 1983. Cretaceous Aporrhaidae from California: Aporrhainae and Arrhoginae. *Journal of Paleontology* 57: 742–765, 6 figs.
- RAFINESQUE, C. S. 1815. *Analyse de la nature ou tableau de l'univers et des corps organisés*. Palermo. 224 pp.
- RISSE, J. A. 1810–1826. *Histoire naturelle des principales productions de l'Europe méridionale et particulièrement de celles des environs de Nice et des Alpes maritimes*. 5 Vols. Paris.
- ROSENKRANTZ, A. 1970. Marine Upper Cretaceous and lowermost Tertiary deposits in West Greenland. *Meddelanden fra Dansk Geologiske Forening* 19:406–453.
- ROSSBACH, T. J. & J. G. CARTER. 1991. Molluscan biostratigraphy of the Lower River Bend Formation at the Martin Marietta Quarry, New Bern, North Carolina. *Journal of Paleontology* 65:80–118.
- ROY, K. 1994. Effects of the Mesozoic marine revolution on the taxonomic, morphologic, and biogeographic evolution of a group: aporrhaid gastropods during the Mesozoic. *Paleobiology* 20:274–296.
- ROY, K. 1996. The roles of mass extinction and biotic interaction in large-scale replacements: a reexamination using the fossil record of stromboidean gastropods. *Paleobiology* 22:436–452.
- SAUL, L. R. 1998. Eight aporrhaid gastropod species from the Cretaceous of the Pacific Slope of North America and clarification of the type species of *Perissoptera*. *The Nautilus* 111:119–142.
- SCHLOTHEIM, E. F. VON 1820–1823. *Die Petrefactenkunde auf ihrem jetzigen Standpunkte durch die Beschreibung seiner Sammlung versteinerter und fossilier überreste des Theirund Pflanzensreichs der Vorwelt erläutert*. Becker'schen Buchhandlung, Gotha, lxii + 437 pp. Atlas pls. 15–29 [1820]; pls. 1–21 [1822]; pls. 22–37 [1823].
- SOHL, N. F. 1960. Archaeogastropoda, Mesogastropoda, and stratigraphy of the Ripley, Owl Creek, and Prairie Bluff Formations. *United States Geological Survey Professional Paper* 331-A, 152 pp., 18 pls.
- SOHL, N. F. 1967. Upper Cretaceous gastropods from the Pierre Shale at Red Bird, Wyoming. *United States Geological Survey Professional Paper* 393-B: B1–B46, 11 pls.
- SOWERBY, J. 1822. *The Mineral Conchology of Great Britain, Vol. 4*. (pt. 61): London, 69–76, pls. 349–354.
- STANTON, T. W. 1893. The Colorado Formation and its invertebrate fauna. *United States Geological Survey, Bulletin* 106, 288 pp., 45 pls.
- STANTON, T. W. 1920. The fauna of the Cannonball Marine Member of the Lance Formation. *United States Geological Survey Professional Paper* 128-A. *Shorter Contributions to General Geology, 1920*: 1–60, pls. 1–9.
- VERMEIJ, G. J. & E. M. VOKES. 1997. Cenozoic Muricidae of the western Atlantic region. Part XII—The subfamily Ocenebrinae (in part). *Tulane Studies in Geology and Paleontology* 29:69–118.
- WARD, L. W. 1992. Molluscan biostratigraphy of the Miocene, middle Atlantic Coastal Plain of North America. *Virginia Museum of Natural History Memoir* 2, [vi] 159 pp, 26 pls.
- WARD, L. W., D. R. LAWRENCE & B. W. BLACKWELDER. 1978. Stratigraphic revision of the middle Eocene, Oligocene, and lower Miocene—Atlantic Coastal Plain of North Carolina. *United States Geological Survey Bulletin* 1457-F:F1–F23.
- WENZ, W. 1940. Superfamilia Strombacea. Pp. 905–948 in O. H. Schindewolf (ed.), *Handbuch de Paläozoologie, Band 6, Prosobranchia, Teil 3*. Gebrüder Borntraeger: Berlin. 1639 pp. [reprinted 1960–1961].
- WHITFIELD, R. P. 1877. Preliminary report on the geology of the Black Hills, containing descriptions of new species of fossils from the Potsdam, Jurassic, and Cretaceous formations of

the Black Hills of Dakota. United States Geographical and Geological Survey of the Rocky Mountain Region (Powell Survey), 49 pp.

WHITFIELD, R. P. 1880. Paleontology of the Black Hills of Dakota. Pp. 325–468 in H. Newton & W. P. Jenney, Report on the Geology and Resources of the Black Hills of Dakota. United States Geographical and Geological Survey of the Rocky Mountain Region (Powell Survey).

WRIGLEY, A. 1938. English Eocene and Oligocene Strombidae and Aporrhaidae. Malacological Society of London Proceedings 23:61–88, pls. 4–6.

APPENDIX

Locality descriptions for fossils illustrated.

866 TU: "Silverdale Beds." Silverdale, North Carolina (marl pit on north side of Webb Creek and east side of unnumbered county road, Silver-

dale, Onslow County, North Carolina. Coll. Dick & Liz Petit, 1968–1970. Belgrade Formation, Haywood Landing Member. latest OLIGOCENE, Chattian.

25073 LACMIP [= UCLA loc. 5073]: limey concretions in shale exposed on north bank Grand River about 762 m (2500') S, 427 m (1400') E of NW cor. sec. 24, T. 21 N, R. 24 E, Bullhead Quad. (USGS 1956), Corson County, South Dakota. Coll: T. Susuki, 1963. Fox Hills Formation, Trail City Member, *Protocardia* zone. CRETACEOUS, Maastrichtian.

28428 LACMIP: Clarendon, Wiltshire, England. Coll.? & id. Arthur Wrigley—sent by Wrigley to UCLA (exUCLA 11439). London Clay. EOCENE, Ypresian.