



FIGURE 11—*Longoconcha* and *Retipirula* from the Pacific slope of North America. 1–8, *Longoconcha eumeka* n. sp., 1–4, paratype, SDNHM 33992, 1–3, $\times 1.3$, 1, apertural view, 2, oblique apertural view showing folds, 3, abapertural view, 4, uppermost 6 mm of apertural view spire, $\times 9.7$; 5, paratype, SDNHM 70968, $\times 0.66$; 6, paratype, SDNHM 25959, $\times 0.76$; 7, 8, holotype, SDNHM 70974, $\times 0.75$. 9, 10, *Retipirula calidula* n. sp. rubber peels, 9, holotype LACMIP 13218, $\times 1.4$; 10, paratype, LACMIP 13219, $\times 1.6$. 11, 12, *Retipirula crassitesta* (Gabb, 1869), $\times 1.7$, hypotype LACMIP 13221, 11, apertural view, 12, back view. 13, 14, *Retipirula pinguis* n. sp., holotype LACMIP 13222, $\times 1.5$ 13, apertural view, 14, back view.

1964), is commonly arched anteriorly with three folds near midpoint of the aperture. In some species the folds are not visible from the aperture. Sculpture of juvenile whorls is strongly axial, becoming more strongly spiral, to the extent, in some species, of loss of axial ribs in adults. The species described herein average

four (with individual variants having three to six) weaker spirals on the ramp, one or two stronger noded spirals at the shoulder position, two to ten widely spaced spirals across the body of the whorl, followed by several more closely spaced, unnoded, weaker spirals on the siphonal neck.

The oldest *Volutoderma* of this study, *V. querna* of Coniacian age, in addition to its small protoconch, differs from modern volutes in having a growth-line sinus on the ramp. This sinus is positioned closer to the posterior suture than is that of *Carota dilleri*, but not at the posterior suture as in *V. averillii*. It is the sinus that Bandel (2003) has indicated requires the inclusion of Volutoderminae in Pyrifusidae Bandel and Dockery, 2001. Dall (1907) claimed that the posterior sinus of *Volutoderma* allowed for protrusion of part of the mantle. Whatever the function of the posterior outer lip sinus, early Late Cretaceous volutes display a migration of this sinus toward the posterior suture and away from its earlier more abapical position. First appearance of the genus *Volutoderma* is in the Coniacian.

During the Campanian, in the southern part of its range, *Volutoderma* diversified and formed three morphological lineages. 1) "Typical" *Volutoderma* consists of *Volutoderma querna* n. sp., *V. averillii* (Gabb), *V. blakei* n. sp., *V. jalama* n. sp., and *V. perissa* n. sp. These species all have three strong, medially placed folds on the columella visible at the aperture. 2) "Angelica" includes *V. angelica* n. sp., *V. elderi* n. sp., and *V. ynezae* n. sp. which have three, more oblique, weaker folds on the columella that are not visible from the aperture. 3) "Magna" includes *V. santana* Packard, *V. magna* Packard, and *V. antherena* n. sp. which have a vasiform shape with a nearly straight columella.

SPECIES OF THE "TYPICAL" LINEAGE

Diagnosis.—*Volutoderma* species with elongate fusiform shell, having three strong columellar folds visible from the aperture and both axial and spiral sculpture. Axial sculpture diminished in maturity but spirals noded on whorl face, unnoded on siphonal area.

VOLUTODERMA QUERNA n. sp.

Figure 4.3–4.13

Diagnosis.—Relatively short *Volutoderma* with nearly conical spire and growth line noticeably broadly sinused on ramp; early whorls with broad sloping ramp. Parietal shield not extending beyond suture of previous whorl, anterior sinus barely deflected to left.

Description.—Shell large, elongate fusiform with conical spire, spire about one third of height, pleural angle 40°–45°, diameter about one third of height; early whorls with low collar at suture, slightly concave on ramp, without well-marked shoulder, convexly inflated about periphery, contracted anteriorly, anterior siphon somewhat deflected to left (apertural view); ramp concavity filled by posteriorward extension of callus shield and crossed by raised traces of outer lip. Growth line with broad posterior sinus on ramp, prosocline across suture, joining parietal-shield extension without additional notch, and slightly convex across periphery of whorl. Sculpture of 12–17 moderately strong, rounded axials, best developed between ramp and anterior siphon, overridden by narrower spiral cords tending to form nodes at axials, spiral interspaces twice as wide as cords, two to six narrow cords on ramp, four to seven noded cords around periphery, six to nine unnoded anterior cords. Aperture elongate, ellipsoidal, sharply narrowed posteriorly; outer lip a little reflexed, especially posteriorly with broad notch in ramp; ramp crossed by strong imbricate growth lines that join posterior edge of parietal shield; columella rather straight with three strong, medially placed folds; inner lip with moderate callus shield extending posteriorly to subsutural collar.

Etymology.—Latin, *querna*, of oak for its occurrence along Oak Run, Shasta County, northern California.

Types.—Holotype LACMIP 13154 from LACMIP 24106, Clover Creek, Area 4, Santonian. Paratypes LACMIP 13155–13158 from LACMIP 10816, Oak Run, Area 4, Coniacian; LACMIP 13159 from LACMIP 10821, Pine Timber Gulch, Area 4, late Coniacian; 13160–13163 from LACMIP 24106, Clover Creek, Area 4, Santonian; 13164 from LACMIP 10797, Clover Creek, Area 4, Santonian; LACMIP 13165 from LACMIP 23624, Chico Creek, Area 6, Santonian; 13166 from LACMIP 23633 Chico Creek, Area 6, Santonian.

Material examined.—About 157 specimens; one specimen from LACMIP loc. 24672, Area 3; ninety-eight specimens, from Area 4, (twenty-four from Member IV, thirty-three from Member V, forty-one from Member VI); fifty-four specimens from Area 6; six specimens from Area 9.

Occurrence.—Age range Coniacian to Santonian. (Area 3) Hornbrook Formation, Osburger Gulch Member; lower Coniacian. (Area 4) Redding Formation, Members IV and lower V, Coniacian; upper V and VI, Santonian.

(Area 6) Chico Creek; Chico Formation, Musty Buck Member, Santonian. (Area 9) ?basal Forbes Formation; upper Santonian.

Discussion.—The extension of the parietal shield beyond the posterior end of the aperture results in the covering of some sculpture and of the subsutural collar. The posterior extension of the shield fills the concavity of the ramp and gives the spire a nearly straight-sided profile. Extension of the outer lip leaves raised, irregularly spaced growth-line ridges crossing this callus-filled area at a small opisthocline angle to the shell axis.

Most specimens of *V. querna* are incomplete as to height; complete diameters are common. Most adult nearly complete specimens are between 70 and 80 mm in height and are, on average, a third smaller than mature *V. averillii*. *Volutoderma querna* paratype LACMIP 13159 from LACMIP 10821 is incomplete but has a diameter of approximately 52 mm, nearly twice as great as that of the next largest specimen; and its postulated height would have been in excess of 150 mm. The spire and ramp of the last whorl have been eroded away; three strong columellar folds are visible; the expanded part of the last whorl is well preserved with five widely spaced noded cords and perhaps two low unnoded cords. Shell of the final 30 mm of the whorl to the outer lip is thickened and has strong growth lines which nearly obliterate the spiral cords. An incomplete specimen from LACMIP loc. 10794 displays five columellar folds, but all other specimens of *Volutoderma* in which the folds are known have three.

Both strength of sculpture and number of cords and ribs are variable, but *V. querna* commonly has fewer, somewhat stronger axial ribs, a straighter sided, more conical spire, a more steeply sloping ramp, and less bend to the anterior siphon than does *V. averillii*. *Volutoderma querna* grades into *V. averillii*, and specimens of latest Santonian and earliest Campanian age, such as those from the Haslam Formation at Brannan Lake, are difficult to assign.

VOLUTODERMA AVERILLII (Gabb, 1864)

Figure 5

Fusus averillii GABB, 1864, p. 83, pl. 18, fig. 34; J. C. MERRIAM, 1895, p. 40.

Volutilithes navarroensis Shumard. GABB, 1864, p. 102, pl. 19, fig. 56. Not *Volutilithes navarroensis* Shumard, 1861, p. 192.

Volutilithes navarroensis Shumard. ? GABB, 1869, p. 120.

Volutoderma navarroensis GABB, 1877, p. 289 (type of the genus).

Fulguraria Navarroensis Shumard. WHITEAVES, 1879, p. 117, pl. 15, fig. 3, 3a; BOLTON, 1965, p. 21. Not *Volutilithes navarroensis* Shumard, 1861.

Fulguraria Gabbii WHITE, 1889, p. 23, pl. 3, fig. 1 [not, p. 46].

Rostellites gabbii (White). WHITEAVES, 1903, p. 356. Not *Rostellites gabbii* (White) of WARING, 1917, p. 68, pl. 9, fig. 8 = *Volutoderma blakei* n. sp.

Volutoderma californica DALL, 1907, p. 10, fn 1, new name for *V. navarroensis* Gabb not Shumard. ANDERSON, 1958, p. 173, pl. 52, fig. 1, 1a.

Volutoderma suciana DALL, 1907, p. 10, fn 4, new name for *Fulguraria Navarroensis* Whiteaves.

Volutoderma gabbii (White). ARNOLD, 1910, figure only, pl. 1, fig. 1 (reprint of White, 1889, pl. 3, fig. 1) not p. 11. Reprinted in ARNOLD AND ANDERSON, 1910, figure only, pl. 23, fig. 1 (reprint of White, 1889), not p. 60 (figure reprinted as *Volutoderma* in FOWKES, 1982, p. 15). ANDERSON, 1958, p. 174 in part "White's holotype"; not specimen from "Sidney Richardson farm" = *Volutoderma blakei* n. sp.

Volutoderma averillii (Gabb). STEWART, 1927, p. 409, pl. 22, fig. 10, 11; SCHENCK AND KEEN, 1940, p. 25, pl. 17, fig. 4 (reprint of Arnold, 1910, pl. 1, fig. 1); WENZ, 1943, p. 1,312, fig. 3,736 (reprint of Stewart, 1927, pl. 22, figs. 10, 11). SHIMER AND SHROCK, 1944, p. 508, pl. 209, fig. 21 (reprint of Gabb, 1864, fig. 56); ELDER AND SAUL, 1993, p. 184, pl. 2, fig. 21.

Longoconcha navarroensis (Shumard). LUDVIGSEN AND BEARD, 1994, p. 96, fig. 61. Reprinted LUDVIGSEN AND BEARD, 1997, p. 115, fig. 72. Not SHUMARD, 1861.

Diagnosis.—*Volutoderma* with large, very fusiform, shell tapering similarly both anteriorly and posteriorly from the periphery; early whorls with concave ramp and noticeable shoulder; longitudinal ribs fading on large individuals, growth line slightly notched at suture; parietal-callus shield extending posteriorly onto

previous whorl and across face of last whorl; anterior siphon deflected to left.

Description.—Shell large, elongate fusiform with somewhat turruculate spire; spire near one third of height, pleural angle about 43° in adult, near 20° in juvenile, shell diameter about one third of height; early whorls with collared suture, concave ramp, and well developed shoulder, convexly inflated about the periphery, contracted anteriorly, anterior siphon deflected to left (apertural view). Growth line of some young specimens with shallow anti-spiral sinus on ramp, of large specimens without sinus and having slight posterior notch at the suture; slight reflection of outer lip resulting in raised-collabral ridges near suture on ramp crossing onto previous whorl and joining posterior end of parietal shield. Sculpture of 12–17 moderately strong, rounded axials, best developed between ramp and anterior siphon, overridden by narrower spiral cords tending to form nodes at ribs; cord interspaces about three times width of cords, commonly two to five narrower cords on ramp, five to six noded cords about the periphery, six to eleven unnoded anterior cords. Aperture elongate ellipsoidal, sharply narrowed posteriorly, with posterior sinus at the suture; columella arched anteriorly with three strong medially placed folds; inner lip with callus shield extending posteriorly onto previous whorl and medially across face of last whorl.

Types.—Of *Fusus averillii*, holotype UCMP 11964, recognized by J. C. Merriam (1895), Tuscan Springs, on Little Salt Creek, Tehama County, northern California, Area 5, Chico Formation, early Campanian. Of *Volutilithes navarroensis* Gabb, 1864, not Shumard; lectotype ANSP 4298 (not 4299 as in Stewart, 1927, p. 409), (Stewart, pl. 22, fig. 11), paralectotype ANSP 4298a (pl. 22, fig. 10) designated by Stewart (1927), Area 5. Of *Volutoderma californica* Dall, 1907, new name for *Volutilithes navarroensis* Gabb, 1864, lectotype ANSP 4298 designated by Stewart (1927), Area 5.

Of *Fulguraria Navarroensis* Whiteaves, 1879, not Shumard; hypotypes GSC 5786, a–e, (Bolton, 1965, p. 21) Sucia Island, San Juan County, Washington, Area 2, lower Cedar District Formation, late early Campanian. Of *Volutoderma suciana* Dall, 1907, syntypes GSC 5786, a–c new name for *Fulguraria Navarroensis* Whiteaves, 1879, lectotype GSC 5786c, here designated (fide Bolton, 1965 figured by Whiteaves, 1879, pl. 15, fig. 3) measured by Dall (1907, p. 10, fn 4); paralectotypes GSC 5786, a–b, d–e (fide Bolton, 1965, GSC 5786 figured by Whiteaves, 1879, pl. 15, fig. 3a).

Of *Fulguraria Gabbi* White, 1889; lectotype USNM 20112 here designated, near Centerville, Butte Creek, Butte County, California, Area 6, Chico Formation, early early Campanian.

Of *Volutoderma californica* Dall (Anderson, 1958), hypotype CASG 10673 from CASG loc. 1125B, Tuscan Springs, Area 5, Chico Formation, early early Campanian.

Of *Volutoderma averillii* (Gabb, 1864) (Elder and Saul, 1993), hypotype LACMIP 12131 from LACMIP loc. 10865, Tuscan Springs, Area 5. Hypotypes this paper, UCR 3435/1 from near Pentz, Area 7; LACMIP 13168 and 13173 from LACMIP 10865 Tuscan Springs, Area 5; 13169, and 13170 from LACMIP 23648, Chico Creek, Area 6; 13271 and 13272 from LACMIP 24082 Tuscan Springs, Area 5; RBCM.EH2004.013.0001 from Nanaimo River Road at the Boulder Creek bridge, Area 1, Haslam Formation, early early Campanian.

Of *Longoconcha navarroensis* (Shumard), not Shumard, 1861, (Ludvigsen and Beard, 1994, 1997), hypotypes VIPM 053 and 054, from Brannan Lake, west of Nanaimo, Area 1, Haslam Formation, early Campanian.

Material examined.—About 168 specimens; thirteen from Area 1, twenty-five from Area 2, seventeen from AREA 5, seventy-seven from Area 6, one from Area 7, five from Area 16, Bell Canyon, twenty-eight from Area 18.

Occurrence.—Early Campanian, zones of *Submortonicerias chicoense* and *Baculites chicoensis* to *Hoplitoplacenticerias vancouverense*, *Canadoceras newberryanum*, and *Baculites inornatus* (see Fig. 2). (Area 1) Haslam, Pender, and lower Cedar District Formations; lower Campanian. (Area 2) lower Cedar District Formation; upper lower Campanian. (Area 5) Chico Formation undifferentiated, lower Campanian. (Area 6) Chico Formation, in Butte Creek upper Musty Buck and Ten Mile Members; in Chico Creek, Ten Mile Member; lower lower Campanian. (Area 7) Chico Formation, Pentz Road Member; lower lower Campanian. (Area 16) Bell Canyon, Chatsworth Formation; upper lower Campanian. (Area 18) Ladd Formation, upper Holz Shale Member; lower Campanian.

Discussion.—*Volutoderma averillii* commonly reached a larger size than *V. querna*. The spire of *V. averillii* is more turruculate than that of *V. querna* or *V. blakei*. The anterior canal of *V. averillii* is also more strongly bent to the left, and the columella is more arched anteriorly. In mature specimens, the growth line of *V. averillii* is not sinused on the ramp but is notched at the suture. The parietal callus shield extends posteriorly beyond the previous suture and covers a broader area across the whorl than in *V. querna* n. sp.

Volutoderma averillii has spire whorls that are more convex

and less slender than those of *V. blakei*, but mature *V. blakei* are larger and stockier than *V. averillii*. Nodular ends of the axial ribs of *V. averillii* are more visible on its spire whorls.

Several juvenile specimens of *V. averillii* from Chico Creek LACMIP loc. 23643 have very early whorls preserved. Shell material is, in most specimens, decalcified enough that the surface flakes away, thereby removing surface details. Two specimens appear to have at least partial protoconchs, which are low turbinate and of about two and one half whorls. The change from protoconch to telococonch is inferred to be at a diameter of 0.6–0.8 mm where the well-rounded whorl profile alters to noticeably taller and straighter-sided and axial ribs become present. By the second or third juvenile whorl, the strong axial ribs are overridden by finer spirals, whose width is about that of the interspaces. At heights of less than 40 mm, spirals are of nearly equal strength and undifferentiated into noded and unnoded (Fig. 5.1, 5.2). At a height of 40 mm, shells are slenderly fusiform with strong axial and spiral sculpture in which interspaces are wider than ribs and cords. Strength of axial ribs is diminished at 80 mm, but shells remain slender.

Larger specimens of *V. averillii* expand in girth, and a robust specimen from the Chico Formation on Butte Creek “near Centerville” was named *Fulguraria gabbi* White, 1889. North of Centerville, the basal conglomerate of the Chico Formation, i.e., the Ponderosa Way Member, rests on pre-Cretaceous metamorphic rocks. Ammonites from localities south of Centerville, along Butte Creek, identified by Matsumoto (1960, p. 19), included *B. chicoensis* and *C. newberryanum*, which are indicative of the early Campanian. Other large, robust specimens of *V. averillii* sensu stricto (e.g., Fig. 5.13, 5.14), with which White’s specimen is conspecific, are known from well-constrained early Campanian Chico Formation outcrops along Butte Creek (Area 6) and near Pentz (Area 7). Despite the imprecise locality and stratigraphic data for White’s locality, his specimen is considered by us to be from the *Baculites chicoensis* Zone and of early Campanian age.

In contrast, Anderson (1958, p. 174) considered *V. gabbi* (White) similar to a specimen collected by S. Richardson from the Hornbrook Formation, Siskiyou County, northern California (CASG loc. 455A = CASG 61798) and of early Senonian age because, according to Anderson, it was from “near *Desmophyllites siskiyouensis* n. sp., *Prionocycloceras crenulatum* (Anderson, 1902) and above the level of *Nowakites klamathonia* n. sp.” Anderson included in CASG loc. 455A specimens collected over time by several individuals, and Matsumoto (1960, p. 2, 3) noted that the fossils were derived from more than one stratigraphic interval, i. e., *P. crenulatum* and *N. klamathonia* (= *Kosmaticeras* fide Matsumoto, 1959b, p. 64) are of Coniacian age and from lower Hornbrook Formation. Whereas “*D. siskiyouensis* n. sp.” = *Desmophyllites dyphylloides* (Forbes, 1846) is late Campanian (Matsumoto, 1959b, p. 10). With the assistance of J. DeMouthe, S. Richardson’s Hornbrook *Volutoderma* specimen was found in the CASG collection (CASG 61798.03). It is *Volutoderma blakei* and has in its matrix a small *Metaplacenticerias pacificum*. Both *M. pacificum* and *M. californicum* (Anderson, 1902) were described from upper Hornbrook strata now referred to as the Rancheria Gulch Sandstone of the Blue Gulch Mudstone Member (e.g., Nilsen, 1984a, 1984b). Its ammonites are indicative of early late Campanian age, an age in agreement with that of *D. dyphylloides*.

White’s (1889, pl. 3, fig. 1) *Volutoderma gabbi* was reprinted by Arnold (1910), Arnold and Anderson (1910), and Fowkes (1982) to accompany discussions of specimens from the Joaquin Ridge Sandstone Member of the Panoche Formation (Area 11). Reports (Arnold, 1910; Arnold and Anderson, 1910) of *V. gabbi* and *V. averillii* from the Joaquin Ridge Sandstone Member are probably based on misidentifications. The Joaquin Ridge Sandstone overlies the *Metaplacenticerias pacificum*-bearing Alcalde Shale, and no undoubted *V. averillii* has been found from so

young a horizon. North of Coalinga near Joaquin Ridge, the Uhalde Shale at CASG 2361 has yielded *Volutoderma perissa* n. sp. Westward of Coalinga, specimens of *V. blakei* have been recovered from redeposited cobbles in conglomerate lenses within the Alcalde Shale.

Anderson (1958) argued that the type specimen of *Fusus averillii* was immature and could not be shown to be the same species as the much larger lectotype for *Volutoderma californica*. However, in support of Stewart's (1927) synonymizing, growth series from the Ten Mile Member of the Chico Formation on Chico Creek (e.g., LACMIP loc. 23643) indicate that Stewart was correct in considering *V. navarroensis* (Gabb), *V. gabbi*, and *V. californica* to be junior synonyms of *V. averillii*.

Dall (1907) thanked J. F. Whiteaves of the Geological Survey of Canada for the loan of original material for *Fulguraria navarroensis* Shumard in Whiteaves, 1879 (Whiteaves listed 10 specimens). Bolton (1965) listed six hypotypes, and five were loaned by the Geological Survey of Canada for this study. According to Bolton GSC 5786c, the tallest of these specimens, is Whiteaves' plate 15, figure 3 (i.e., Fig. 5.11, 5.12). Although Dall (1907) named these specimens *Volutoderma suciana* and referred to Whiteaves, 1879, plate 15, figure 3, 3a, he did not designate a type specimen for *V. suciana*. He considered it more closely related to *V. gabbi* than to *V. californica*, both of which we consider to be *V. averillii*. He differentiated *V. suciana* by its large size and lack of axial ribs from *V. californica*, not realizing that these are mature characteristics of the species.

VOLUTODERMA BLAKEI new species

Figure 6.1–6.4

Rostellites gabbi (White). WARING, 1917, p. 68, pl. 9, fig. 8. Not *Fulguraria Gabbi* White, 1889 or *Rostellites gabbi* (White) of Whiteaves, 1903.

Volutoderma gabbi (White). ANDERSON, 1958, p. 174, in part (specimen found by S. Richardson); Not *Fulguraria Gabbi* White, 1889.

Diagnosis.—Large *Volutoderma*, shell slender in youth but with expanded penultimate and ultimate whorls; both spiral and axial sculpture at small and intermediate shell sizes, but primarily spiral cords at maturity; growth line sharply sinused at suture.

Description.—Shell large, elongate fusiform, slender in youth becoming stout as adult; spire high, near one third of shell height; suture just anterior to shoulder; pleural angle 32°; diameter of shell less than one third of height; early whorls with steeply sloping ramp and noticeable shoulder, shoulder becoming effaced and marked only by sculpture; ultimate whorl with enlarged diameter; anterior siphonal neck long, gently flexed. Growth line sharply sinused at suture, nearly straight across the body whorl. Early whorls with strong axial ribs; spirals present by diameter of 4 mm; next whorl with four spirals on well-defined steep ramp and two spirals overriding posterior ends of eight to ten axial ribs forming shoulder; about seven to nine wide-spaced, noded spirals on body whorl; unnoded, more closely spaced spirals on siphonal neck. Aperture elliptical, long, and narrow, with narrow posterior sinus at suture; anterior sinus long, open; columella with three folds visible from aperture; inner lip callused, callus shield extending posteriorly to shoulder.

Etymology.—The species is named for William Blake, a member of the original California Geological Survey.

Types.—Holotype LACMIP 13174 from LACMIP 22415, Bee Canyon, Lomas de Santiago, Area 18. Paratypes LACMIP 13175 from LACMIP 8186, Aliso-Santiago Creek divide, Area 18; LACMIP 13176 from LACMIP 10105 Aliso-Santiago Creek divide, Area 18; LACMIP 13177 from LACMIP 26471 Cooper Canyon, Area 11; LACMIP 13178 from LACMIP 22415, Bee Canyon, Area 18; LACMIP 13359 and 13177 from LACMIP 17829, Bee Canyon, Area 18; CASG GeTy 10760 from CASG 61798.03 near Montague, Area 3; CASG 68821.01 from CASG 68821 Dayton Canyon, Area 16.

Material examined.—About 84 specimens; one from CASG loc. 61798, Area 3; two from Cooper Canyon cobble in conglomerate, Area 11 (a cast in sandstone may also be this species); nine from Area 16; one from Area 17; forty-three from Area 18, (one from Schulz Ranch Conglomerate Member, forty-two from Pleasants Sandstone Member).

Occurrence.—Early middle to early late Campanian, zones of *Hoplitoplacenticerans?* *bowersi* and *Metaplacenticerans pacificum*. (Area 3) Hornbrook Formation, Blue Gulch Mudstone Member, Rancheria Gulch Sandstone beds; lower upper Campanian. (Area 11) Panoche Formation, cobble in Alcalde Member; middle Campanian. (Area 16) Chatsworth Formation; middle Campanian. (Area 17) Tuna Canyon Formation; lower upper Campanian. (Area

18) Williams Formation, Schulz Ranch Conglomerate Member, middle Campanian, and Pleasants Sandstone Member, lower upper Campanian.

Discussion.—The wide steep ramp of *V. blakei* gives its early whorls a more straight-sided aspect, whereas the concave ramp and strong shoulder nodes make early whorls of *V. averillii* more turriculate. *Volutoderma blakei* has a larger number of noded spirals over the whorl periphery than do *V. averillii* or *V. querna*. *Volutoderma averillii* has a more strongly flexed anterior canal than does *V. blakei*. Several specimens of *V. blakei* with a diameter of over 70 mm are from LACMIP loc. 22415, but none has both a complete spire and anterior siphon. A probable height for LACMIP 13174 with a diameter of 73 mm, estimated by adding spire height and siphonal length from two other specimens, would be about 206 mm.

Volutoderma blakei occurs with *Hoplitoplacenticerans?* *bowersi* and with *Metaplacenticerans pacificum* in southern California. Recognition of *V. blakei* with *M. pacificum* in the Blue Gulch Mudstone Member of the Hornbrook Formation near Montague, northern California, is discussed under *V. averillii*. Although most specimens from the lower Cedar District Formation on Sucia Island are *V. averillii*, some have taller spires than typical *V. averillii* and reduced shoulder development similar to *V. blakei* (see CASG 228.06).

VOLUTODERMA JALAMA new species

Figure 6.5–6.13

Volutoderma sp. cf. *V. averillii* (Gabb, 1864). DAILEY AND POPENOE, 1966, fig. 3.

Diagnosis.—*Volutoderma* with relatively short, strongly sculptured shell; spire narrowly tapered with concave profile, whorl abruptly widened at maturity to much wider adult pleural angle. Posterior sinus strongly V-shaped, creating raised strong spiral track at suture.

Description.—Shell moderately large, plumply fusiform, almost biconic in youth becoming roundly stout as adult; spire near one third of shell height; suture just anterior to shoulder; juvenile pleural angle 37°, adult pleural angle 50°–60°; maximum diameter of shell near one third of height; early whorls with steeply sloping ramp, shoulder marked mainly by change in sculpture; ultimate whorl with enlarged diameter. Growth line sharply V-sinused at suture, nearly straight across side of whorl. Both axial and spiral sculpture relatively strong; axial ribs persistent on ultimate whorl; three spiral cordlets on ramp, about five to seven wide-spaced noded spiral cords on whorl; unnoded, more closely spaced spirals on siphonal neck; 13–16 axial ribs strong at shoulder fading toward siphonal neck. Track of V-shaped posterior notch leaving a strong spiral at suture. Aperture elliptical; columella with three folds visible from aperture; inner lip callused, parietal callus shield extending posteriorly to shoulder.

Etymology.—Named for Jalama Creek, Santa Barbara County, California.

Types.—Holotype LACMIP 13179 from LACMIP 6243. Paratypes LACMIP 13180 from LACMIP 6242; LACMIP 13181 from LACMIP 10695; LACMIP 13182 from LACMIP 10701; LACMIP 13183 from LACMIP 24120; LACMIP 13184 from LACMIP 24144. All localities in Area 15.

Material examined.—Twenty-one specimens from Area 15 and one poorly preserved specimen from Area 12 (LACMIP loc. 24183).

Occurrence.—Late late Campanian, zone of *Baculites rex*. (Area 15) Jalama Formation. (Area 12) Panoche Formation.

Discussion.—*Volutoderma jalama* is very similar to *V. blakei* n. sp. but differs from the latter in being relatively stouter for its height, having stronger, rougher-appearing sculpture, a deeper, very V-shaped posterior siphonal notch that forms a strong welt-like cord along the suture (Fig. 6.12), and in having succeeding whorls envelop more of the previous whorl. The sharp, narrow spire of *V. jalama* does not have the steep sutural slope of the *V. angelica* group but has a sutural slope more similar to that of *V. perissa* n. sp.

VOLUTODERMA PERISSA n. sp.

Figure 7.1–7.4

? *Volutoderma* cf. *magna* Packard. ANDERSON AND HANNA, 1935, p. 26.
Volutoderma cf. *V. californica* Dall. SUNDBERG AND RINEY, 1984, fig. 3.8.
Volutoderma sp. cf. *V. magna* Packard. ELDER, SAUL, AND POWELL, 1998, p. 160, pl. 1, fig. 11.

Diagnosis.—Elongate, relatively narrow whorled *Volutoderma*, shell lacking shoulder angulation and having aciculate spire. Axial sculpture consisting mainly of aligned nodes on cords.

Description.—Shell large, elongate ovoid with aciculate spire; spire profile concave; ramp demarcated by closer spacing of four weaker cords; body whorl unshouldered; pleural angle of aciculate spire 22°, of mature spire about 45°; diameter near one third height of shell; anterior siphon very moderately flexed. Growth line strongly notched at suture. Sculpture of weak axial costae, expressed mainly by aligned nodes on cords; about eight noded, widely spaced, narrow cords on last whorl, and five or more unnoded cords on siphonal neck. Aperture elongate, ellipsoidal, widest near base of penultimate whorl; columella with three strong folds visible from aperture, just adapical to widest part of aperture; inner lip callused, pallial callus shield covering and extending beyond apertural face of body whorl, posteriorly covering suture and apertural face of penultimate whorl.

Etymology.—Greek, *perissos*, odd, strange.

Types.—Holotype SDNHM 33994 from SDNHM 3405, Area 19. Paratypes CASG 2361.07 from 2361, Area 11; LACMIP 13185 and LACMIP 13186 from LACMIP 2855, Area 19; LACMIP 13187 and LACMIP 13188 from LACMIP 23268, Area 19; LACMIP 13189 from LACMIP 27136, Area 20; SDNHM 25968 from SDNHM 3162, Area 19; SDNHM 33993 and SDNHM 70972 from SDNHM 3405, Area 19. Hypotype UCMP 39906 from USGS M8830, Area 8.

Material examined.—Thirty-two specimens; one from Area 8, Anchor Bay; two from Area 11, Joaquin Ridge, north of Coalinga; one from Area 16, "Lang Ranch," west end Simi Hills; two from Area 19, Carlsbad Research Center; one from Area 20, Punta Banda; two from Area 21, near San Antonio del Mar; twelve from Area 22, one from Cañon San Fernando, eleven from Arroyo Santa Catarina.

Occurrence.—Late Campanian/Early Maastrichtian, commonly in association with *Baculites lomaensis*. (Area 8) Gualala Formation, Anchor Bay Member. (Area 11) Panoche Formation, Uhalde Shale Member. (Area 16) "Lang Ranch," west end Simi Hills, upper Chatsworth Formation. (Area 19) Point Loma Formation. (Area 21) and (Area 22) Rosario Formation.

Discussion.—*Volutoderma perissa* resembles other "Typical" *Volutoderma* in having three strong columellar folds that are clearly visible within the aperture, although the most anterior fold extends forward less than the other two. *Volutoderma perissa* differs from the other "Typical" species in its aciculate spire, convex ramp rounding into the whorl side without an angulate shoulder, and more enveloping mature whorls. Its spire is most similar to that of *V. jalama* but is more aciculate.

A protoconch is not preserved on the studied specimens. The earliest whorl preserved on SDNHM 33993 has a diameter of 1 mm; diameter of its fifth whorl is 5 mm; of its eighth whorl is 15 mm; height of these eight whorls is 24.5 mm. Sculpture on the early whorls appears to have been covered by callus deposits.

Identification of incomplete specimen UCMP 39906 is somewhat questionable, and its locality USGS M8830, Area 8, Gualala Formation at Anchor Bay is the most northern locality known for this species. Otherwise the most northern locality for *Volutoderma perissa* is at CASG loc. 2361, Area 11, the type locality of *Baculites lomaensis* Anderson, northwest of Coalinga, Fresno County, California.

Arnold (1910, p. 11, pl. 1; in Arnold and Anderson, 1910, p. 60, pl. 23) listed *Volutoderma gabbi* (White) from his "loc. 3" northwest of Coalinga, Area 11. As noted earlier, the figure is a reprint of White's 1889 *Volutoderma gabbi* = *V. averillii* from Butte Creek. Included on Arnold's plate with the *Volutoderma* are three misidentified bivalves which were also from "loc. 3," northwest of Coalinga: "*Mactra ashburneri* Gabb, 1864" = *Cymbophora triangulata* (Waring, 1917), "*Meekia sella* Gabb, 1864" = *Meekia (Mygallia) daileyi* Saul and Popenoe, 1962, and "*Tellina? ooides* Gabb, 1864" = *Willimactra (Petromactra) truncata* (Gabb, 1864). These bivalves indicate a younger age than that of *V. averillii*. All three bivalves occur in the Jalama Formation (Saul and Popenoe, 1962; Saul, 1973, 1974), which has yielded *V. jalama* n. sp. The bivalves are also present in the Rosario Formation of Baja California associated with *V. perissa* n. sp., which occurs in the Uhalde Shale Member of the Panoche Formation north of Coalinga, Fresno County.

Anderson and Hanna (1935) reported two imperfect specimens of *Volutoderma* cf. *V. magna* from near Santa Catarina Landing, but available specimens from the Rosario Formation near Santa Catarina Landing are either *L. eumeka* n. sp. or *V. perissa*.

SPECIES OF THE "ANGELICA" LINEAGE

Diagnosis.—*Volutoderma* species having ovoid, greatly enlarged last whorl; high, narrow spire with steep sutural slope; and three, oblique columellar folds not visible from aperture; sculpture dominantly spiral; axial sculpture reduced in maturity, becoming nearly obsolete in some.

VOLUTODERMA ANGELICA new species

Figure 8.1, 8.2

Diagnosis.—Large *Volutoderma* with high, pointed spire and roundly expanded adult whorls; early whorls weakly sculptured, expanding slowly with steep sutural slope; parietal callus shield thick, expanded roundly over face of last whorl.

Description.—Shell large, fusiform, round-shouldered, with ramp delimited only by sculpture; spire high, narrow with steep sutural slope, and rounded whorl profiles, spire height about one fifth of shell height; pleural angle 23°. Penultimate and ultimate whorls enlarged, roundly expanded posteriorly and contracted anteriorly; diameter of shell about one third of height. Growth line with strong notch at suture; reflection of outer lip results in raised collabral ridges near suture on ramp crossing onto previous whorl and joining posterior end of parietal shield. Sculpture of about 16 broad, weak, rounded axial costae overridden by widely spaced, narrow spiral cords, spiral cords forming elongate nodes at intersections with axials on strongly sculptured specimens; ramp defined by finer, up to five, close-spaced cords and absence of axials. Aperture elongate, expanded medially; columella somewhat arched anteriorly, callused, with three, medially placed, oblique folds, not visible from the aperture; parietal shield very thick, spreading roundly over face of last whorl and expanded posteriorly to suture. Outer lip reflected, especially posteriorly, lip extended adapically to notch-like posterior sinus, and forming thickened right edge of shield.

Etymology.—Named for Los Angeles, Latin, *angelicus*, angelic.

Types.—Holotype LACMIP 13190. Paratypes LACMIP 13191–13193; LACMIP 13194–13196. All types from LACMIP 10715 = LACMIP 6965, Dayton Canyon, Area 16.

Material examined.—About thirty-five specimens; thirty-one from Area 16, Dayton Canyon; four questionably identified specimens from (Area 18) Santa Ana Mountains (two from upper Holz Shale; two from Schultz Conglomerate).

Occurrence.—Early early? Campanian to early middle Campanian, associated with *Hoplitoplacenteras? bowersi* Anderson, 1958, *Baculites* cf. *B. inornatus* Meek, 1862 (Matsumoto, 1960) and *B. anceps pacificus* Matsumoto and Obata, 1963 (Saul and Alderson, 1981) in Dayton Canyon. (Area 16) lower Chatsworth Formation; lower middle Campanian. (Area 18) questionably identified specimens upper Holz Shale Member, Ladd Formation, lower Campanian; partial specimens, Williams Formation, Schulz Ranch Conglomerate Member; ?lower middle Campanian).

Discussion.—*Volutoderma angelica* resembles *V. magna* in having a narrow but high spire with steep sutural slope and rapidly enlarging penultimate and ultimate whorls, but none of the specimens from Dayton Canyon develops the ascending, callus-bordered, U-shaped posterior sinus or the expanded and enveloping outerlip of *V. magna*. Additionally, specimens of *V. angelica* have a more rounded body whorl and a columella that is arched anteriorly. The early whorls of *V. angelica* have a more convex profile and commonly show three noded cords. Sculpture is best preserved on paratype 13194. A few poorly preserved specimens from the uppermost Holz Shale (e.g., Area 18, LACMIP loc. 10723) resemble *V. angelica* in having a convex profile and three noded cords on early whorls but do not have as steep a sutural slope as *V. angelica* from Dayton Canyon.

Volutoderma angelica is most similar to *V. elderi* n. sp., which is more weakly sculptured, has fewer, more widely spaced spiral cords, and nearly lacks axial costae and nodes on the spirals.

At Dayton Canyon, *V. angelica* occurs with ammonites previously identified as *Metaplacenteras californicum* (?) (Anderson, 1902) or *M. pacificum* and *M. sanctamonicae* Waring, 1917 (Popenoe, 1954). As pointed out to us by J. M. Alderson, the most

abundant Dayton Canyon ammonite is *Metaplacenticerases? bowersi* Anderson, 1958 (type locality Dayton Canyon) which Matsumoto (1984) referred to *Hoplitoplacenticerases? bowersi* (Anderson, 1958). Dayton Canyon outcrops yield both *V. blakei* n. sp. and *V. angelica*, but *V. angelica* is far more common.

Schoellhamer et al., (1981) listed two specimens from the Schulz Ranch Conglomerate Member of the Williams Formation (LACMIP loc. 15900 = SA66 = F43 of Schoellhamer et al., 1981) as *Volutoderma santana*. They do not have the strong sculpture, shouldered whorls or wide ramp cords of *V. santana*, but have instead sculpture, whorl shape, and steep sutural slope more similar to *V. angelica*, and suggest a correlation of the Schulz Ranch Conglomerate Member with the Chatsworth Formation of Dayton Canyon.

VOLUTODERMA ELDERI new species

Figure 8.3–8.7

Diagnosis.—Large, round-shouldered *Volutoderma* sculptured by widely spaced, narrow cords and having high, narrow spire with very steep sutural slope.

Description.—Shell large, fusiform, round-shouldered, with ramp delimited only by sculpture; spire high, narrow, with steep sutural slope, and rounded-whorl profile; spire height about one third of shell height; pleural angle without penultimate whorl 23°, with penultimate 38°. Penultimate and ultimate whorls enlarged, roundly expanded posteriorly, and contracted anteriorly; diameter of shell about one third of height. Growth line with strong notch at suture. Sculpture of three to four weak cords on ramp (may be effaced), nine to twelve widely spaced, narrow spiral cords, commonly unnodded on last whorl; interspaces four to six times width of cords; axial ribs weak, present in juveniles, lost on mature shells. Aperture elongate, expanded medially; columella barely arched anteriorly, callused, with three, medially placed, oblique folds not visible from aperture (Fig. 8.6); parietal shield roundly expanded, thin over face of whorl, thick near posterior end of aperture, thickest adapical to aperture. Outer lip thin, deeply sinused at suture.

Etymology.—Named for W. P. Elder, whose work on a Cretaceous Pacific-Slope time scale was very helpful.

Type specimens.—Holotype LACMIP 13197 from LACMIP 10120, Santiago-Aliso Canyon divide, Area 18. Paratypes LACMIP 13198–13200 from LACMIP 10105, Area 18; LACMIP 13201 from LACMIP 8186, Area 18; LACMIP 13202 from LACMIP 21530, Area 18; LACMIP 13203 from LACMIP 24207, Area 18; LACMIP 13358 from LACMIP 17828, Area 18.

Material examined.—At least twenty-nine specimens; one partial specimen from Area 16, head of Dayton Canyon; one poorly preserved specimen from Area 17; twenty-seven partial specimens from Area 18.

Occurrence.—Late middle to early late Campanian, *Metaplacenticerases pacificum* Zone. (Area 16) Chatsworth Formation; late middle Campanian. (Area 17) Tuna Canyon Formation; early late Campanian. (Area 18) Williams Formation, Pleasants Sandstone Member; early late Campanian.

Discussion.—Juveniles of *V. elderi* are very tall and slender. Early whorls up to a diameter of about 9 mm have axial ribs, but these fade on the following two whorls. The ultimate whorl of the most complete specimen, LACMIP 13358, is flattened, making measurement of diameter impossible. A diameter of 22 mm, measured on its relatively uncrushed penultimate whorl, is too low as the greatest diameter of that whorl was doubtless within the area crushed. Anterior siphonal areas and the outer lip are missing from all available specimens. Outer lip of paratype LACMIP 13358 has been broken back far enough to expose the columellar folds (Fig. 8.6). *Volutoderma elderi* is very similar to *V. angelica* n. sp. from which it differs in having a less distinctly marked ramp that is somewhat more sloping than in *V. angelica*, and spirals that are more widely spaced and weakly noded to the extent of being unnodded on the ultimate whorl. *Volutoderma elderi* resembles *V. ynezae* n. sp. in lacking nodes on the spirals when mature but differs in having fewer more widely spaced spirals and in lacking the subangulate shoulder of *V. ynezae*.

VOLUTODERMA YNEZAE new species

Figure 8.8–8.11

Volutoderma magna? PACKARD, 1922. DAILEY AND POPENOE, 1966, figs. 2 and 3, listed only.

Diagnosis.—*Volutoderma* with an aciculate spire, strong, rounded shoulder, 16–17 spiral cords, and no axial ribs on mature whorls. Columellar folds not visible from aperture.

Description.—Shell large, almost strombiform; spire high, narrow with steep sutural slope, pleural angle not measured, maximum diameter of shell near shoulder; early whorls not available; last whorl roundly inflated, slightly angular at shoulder, wrapping posteriorly, and enveloping previous whorl; suture of body whorl climbing on last half whorl to previous suture at aperture; ramp flatly sloping, nearly smooth, complete anterior siphon not available. Growth line strongly notched at suture, relatively straight across whorl. Sculpture on adult of 16–17 narrow spiral cords with wider interspaces, interspaces two or three times width of cords. Aperture made wide by expansion of last whorl; columella arched anteriorly; inner lip heavily callused, wrapping onto back of siphonal neck anteriorly; outer lip forming deep U-shaped notch at suture; parietal shield apparently thin, expanded roundly over apertural face of last whorl.

Etymology.—Named for the Santa Ynez Mountains.

Types.—Holotype LACMIP 13204 from LACMIP 10690, Jalama Creek, Area 15. Paratypes LACMIP 13205 from LACMIP 10690, Area 15; LACMIP 13206 from LACMIP 24118, Area 15; LACMIP 13207 from 24128, Area 15.

Material examined.—Eighteen specimens from Area 15.

Occurrence.—Late late Campanian, *Baculites rex* Zone. (Area 15) Jalama Formation.

Discussion.—As in *V. magna*, *V. elderi* n. sp., and *V. angelica* n. sp., the columellar folds of *V. ynezae* are deep within the aperture and out of sight. The profile of *V. ynezae* is similar to that of *V. magna* in developing a shoulder but is more roundly expanded anterior to the shoulder, giving *V. ynezae* the appearance of being less tall. None of the *V. ynezae* specimens has as widely expanded a final whorl as *V. magna*, and all *V. ynezae* have a more arched columella. The columella of *V. ynezae* is also more strongly arched than that of *V. elderi*. *Volutoderma ynezae* has less axial sculpture on its mature whorl than any of the other volutes discussed in this paper. Some specimens of *V. ynezae* have no axial ribs or nodes on less-than-mature whorls. Spiral cords on *V. ynezae* are commonly more numerous and closer together than cords on *V. angelica* and *V. elderi*.

The pleural angle of the immature whorls of *V. ynezae* appears to be have been narrow and the spire apparently had a steeply sloping sutural angle, but all available specimens of *V. ynezae* have broken spires and retain barely more than the mature whorls, which provide a wide pleural angle.

Submature specimens of *V. ynezae* resemble *V. perissa* in blimplike shape and perhaps an aciculate spire, but *V. ynezae* is less elongate, has spirals that are closer together and less noded, and has weaker columellar folds that are not visible from the aperture.

A submature specimen, from LACMIP 24118, which had not yet developed a shoulder, has a thin parietal shield roundly expanded over the apertural face of its last whorl. The posterior suture of its last whorl has climbed toward the previous suture but not up to it; despite the specimen's immaturity, a strong callus is present at the posterior siphonal notch. The spiral cords of this specimen are faintly noded.

SPECIES OF THE "MAGNA" LINEAGE

Diagnosis.—Strongly sculptured *Volutoderma* that tend to be vasiform with a relatively straight columella; two strongly noded cords on the shoulder, and two? to eight more anterior, strongly noded cords; growth line strongly notched on ramp; three columellar folds present but not all visible from aperture.

VOLUTODERMA SANTANA Packard, 1922

Figure 9

Volutoderma santana PACKARD, 1922, p. 432, pl. 36, fig. 3; ANDERSON, 1958, p. 175; ELDER AND SAUL, 1993, p. 184, pl. 2, fig. 20.

Diagnosis.—A relatively short-spined almost pyriform *Volutoderma* with shouldered whorls and strongly noded sculpture, un-noded cords on ramp wider than interspaces. Anterior canal long, curved to left. Columella with three folds, two visible at aperture.

Description.—Shell medium-sized, subpyriform, height near two and one half times diameter, with turriculate spire about one fourth of shell height, pleural angle 53°, early whorls with collar at suture, whorl concave on ramp with well-marked shoulder, whorl convexly inflated about periphery, adult whorls contracted anteriorly, anterior siphon somewhat deflected to left (apertural view). Growth line notched at suture, barely notched prosoclinally across ramp and nearly straight to siphonal tip. Sculpture of about 9–12 widely spaced, strong, short, rounded ribs per whorl; ribs best developed just anterior to shoulder, overridden by strong spiral cords forming nodes at intersections with ribs; commonly two strongest cords just anterior to shoulder, four to five noded cords anterior to shoulder cords; about four unnoded spirals on siphonal neck, ramp with two to four weaker unnoded spirals, posterior two or three of which being as wide as interspaces; cords at nodes about equal in width to interspaces. Aperture elongate, ellipsoidal, narrowed posteriorly with notch at suture; columella slightly arched anteriorly with two of three strong, medially positioned folds visible within aperture; inner lip with moderate callus shield extending posteriorly to suture.

Type specimens.—Holotype UCMP 12294 from UCMP 2135, Area 18. Hypotypes LACMIP 12130 from LACMIP 26467, Area 10; LACMIP 13208 from LACMIP 10084; LACMIP 13209 from LACMIP 10080; LACMIP 13210 from LACMIP 10093; and LACMIP 13211 from LACMIP 15904, Area 18.

Material examined.—Eighty specimens: one from Area 10, north of Pigeon Point; seventy-seven from Area 18, Santa Ana Mountains. Questionably identified two: one from Area 16, Dayton Canyon, and one from Area 18, Williams Formation, Schulz Ranch Conglomerate Member.

Occurrence.—Early Campanian and possibly early middle Campanian. (Area 10) Pigeon Point Formation; lower or lower middle Campanian. (Area 16) lower Chatsworth Formation, lower middle Campanian. (Area 18) Ladd Formation, upper Holz Shale Member; lower Campanian; and possibly Williams Formation, Schulz Ranch Conglomerate Member; lower middle Campanian.

Discussion.—Packard's (1922) referral of the holotype of *V. santana* to the "*Turritella pescaderoensis* zone" is evidently an error. Popenoe (1937, 1942) showed that species of Packard's "*Turritella pescaderoensis* zone" came from the Williams Formation, Pleasants Sandstone Member, of middle to early late Campanian age, but Popenoe (1942) identified as *V. santana* specimens from the upper Holz Shale Member of the Ladd Formation. No examples of *V. santana* are known from the Pleasants Sandstone Member. Matsumoto (1960, p. 143) listed *Baculites* cf. *B. inornatus* Meek, 1862 from UCMP loc. 2135, the type locality of *V. santana*, and recognized *B. inornatus* from three uppermost Holz Shale localities. Packard's holotype of *V. santana* lacks spire and anterior canal, but its shoulder strength and type of sculpture, especially on the ramp, identify it with the more than 70 specimens from the upper Holz Shale here identified as *V. santana*.

Three specimens from the Schulz Ranch Conglomerate Member of the Williams Formation LACMIP locs. 15900 and 15904 were listed in Schoellhamer et al. (1981) as *V. santana*. However, the two specimens from LACMIP loc. 15900, although very incomplete, have more characteristics of *V. angelica* n. sp. than of *V. santana*. *Volutoderma angelica* has otherwise only been found in the Chatsworth Formation in strata yielding *Hoplitoplacenticeraster? bowersi* (see Fig. 2). A mangled Chatsworth Formation specimen from LACMIP 10715 has nodes like those of *V. santana*. Possible co-occurrence of these two volutes, if correctly identified, gives *V. santana* a range into the early middle Campanian and an early middle Campanian age for the Schulz Ranch Conglomerate.

Volutoderma santana has a more southern distribution than *V. averillii*, whose range extends from British Columbia (Fig. 1) south to Orange County, California (Fig. 3). The northernmost occurrence of *V. santana* is in the southern sequence of the Pigeon Point Formation (Elder and Saul, 1993, pl. 2, fig. 20; this paper, Figs. 1, 3, Area 10).

Juvenile specimens of *V. santana* have the broadest apex of any *Volutoderma* considered here. Juveniles of *V. santana* are readily distinguished from juveniles of *V. averillii*, *V. querna*, *V. angelica*, and *V. magna* by having strong shoulder nodes, very short axial costae with wide interspaces, and a low spire with a

wider pleural angle than these other species. Mature *V. santana* has larger nodes and fewer costae per whorl than *V. averillii*, *V. querna* n. sp., and *V. angelica*. It is most similar to submature *V. magna* in sculpture and whorl shape, but lacks the expanded development of the ultimate whorl of mature *V. magna*. The ramp sculpture of *V. santana* differs from other *Volutoderma* in consisting of cords that are wider than their interspaces.

VOLUTODERMA MAGNA Packard, 1922

Figure 10.1–10.6

Volutoderma magna PACKARD, 1922, p. 432, pl. 37, fig. 1; ANDERSON, 1958, p. 175.

Volutoderma magna (?) PACKARD, 1922, p. 432, pl. 38.

Not *Volutoderma* sp. cf. *V. magna* PACKARD, ELDER, SAUL, AND POWELL, 1998, p. 160, pl. 1, fig. 11 = *V. perissa* n. sp.

Probably not *Volutoderma* cf. *magna* PACKARD, ANDERSON AND HANNA, 1935, p. 26.

Diagnosis.—*Volutoderma* with large strombiform shell, widely expanded outer lip, growth line strongly notched at suture, posterior notch defined by thick, vertical, U-shaped callus; spire high, pointed, with slender whorls and steep sutural slope; columella straight and heavily callused.

Description.—Shell large, strombiform (Packard, 1922, pl. 38), with greatest diameter at posterior part of whorl; adult whorls subshouldered with steeply sloping, slightly concave ramp rising to collar at suture; spire high, pointed, with slender, weakly sculptured, unshouldered whorls, and steep sutural slope; spire height less than a quarter of shell height; pleural angle about 35°, pleural angle of young specimens 27°. Penultimate and penultimate whorls expanded posteriorly to form rounded shoulder, penultimate and ultimate whorls enlarged, ultimate whorl enveloping penultimate whorl; diameter of shell about one half of height; anterior siphon somewhat deflected to left (apertural view). Growth line with strong, callused, U-shaped posterior notch at suture; slight reflection of outer lip resulting in raised collabral ridges near suture on ramp, crossing onto previous whorl, and joining posterior end of parietal shield. Sculpture of 17–18 commonly weak axial ribs overridden by 8–11 widely spaced, spiral cords forming nodes at intersections with costae, ramp with four weak cords, siphonal neck with at least four cords. Aperture elongate, widely expanded in maturity, columella rather straight, heavily callused, with three moderately oblique, medially placed folds, deep within aperture; parietal shield spreading over face of last whorl and expanded posteriorly onto previous whorl.

Types.—Holotype UCMP 12274 from UCMP 2166, Area 18. Hypotypes UCMP 12278 from UCMP 2166, Area 18; LACMIP 13212 from LACMIP 15901, Area 18; LACMIP 13213 from LACMIP 17419, Area 17; LACMIP 13214 from LACMIP 10096, Area 18.

Material examined.—Twenty-four specimens; twelve from Area 18; twelve from Area 17.

Occurrence.—Late middle to early late Campanian, *Metaplacenticeraster* Zone. (Area 17) Tuna Canyon Formation, upper middle Campanian; (Area 18) Williams Formation, Pleasants Sandstone Member, lower upper Campanian.

Discussion.—Packard (1922) hesitated to include the specimen that was larger and nearly devoid of shell, UCMP 12278, in *V. magna*, but it is a more mature example of the species in which the last whorl has wrapped over the moderately shouldered penultimate whorl, leaving one of the slender spire whorls exposed. *Volutoderma magna* resembles *V. angelica* n. sp. in having a high slender spire with a steep sutural angle but differs from *V. angelica* in developing a vertical U-shaped gutter at the posterior end of the aperture, in having a straighter more heavily callused columella, in having more strongly noded spiral cords, and in having the whorl's periphery more posteriorly placed.

The overall aspect of the shell of *V. magna* with its high-pointed spire, very widely expanded outer lip, and posterior expansion of the whorl diameter, which narrows anteriorly without much flexure to the anterior siphon, is somewhat like a very elongate top. These are the largest volutes in the Santa Ana Mountains. Although Packard (1922) noted the prominent shoulder of *V. magna* as a characteristic distinguishing *V. magna* from "*V. gabbi*," the spire whorls of *V. magna* are not shouldered, nor is the last whorl. The penultimate and penultimate whorls expand posteriorly, forming a rounded shoulder, as in the holotype of *V.*

magna, but the final whorl wraps over the penultimate whorl and produces an unshouldered profile.

Anderson (1958) questioned the Campanian age of *V. magna*, but *V. magna* is associated in both Area 18, the Santa Ana Mountains, and Area 17, the Santa Monica Mountains with *Metaplasticeras pacificum*, an ammonite indicative of late middle to early late Campanian age (Fig. 2). Also *Anapachydiscus californicus* (Yabe in Yabe and Shimizu, 1921) [= *Pachydiscus (Parapachydiscus) fascicostatum californicum* Yabe], which was described from UCMP 2166, the type locality of *V. magna*, is Campanian (Matsumoto, 1960).

Specimens listed by Dailey and Popenoe (1966) as *V. magna* are *V. ynezae* n. sp. Anderson and Hanna (1935) referred two "imperfect specimens" of *Volutoderma* from near Santa Catarina Landing, Baja California, to "*Volutoderma* cf. *magna* Packard," but specimens from near Santa Catarina examined for this study are the new species *Volutoderma perissa* and *Longoconcha eumeka*.

VOLUTODERMA? ANTHERENA new species

Figure 10.7, 10.8

Diagnosis.—Vasiform volute with four strong spiral cords heavily noded at intersections with broad axial costae; shell slightly constricted just posterior to columellar folds; siphonal area with noded cord anterior to constriction. Spire conical. Growth line with notch at suture, prosocert across the ramp, sinused at the posterior row of nodes, and relatively straight toward the anterior sinus.

Description.—Shell moderately large, vasiform, basally contracted; whorl broadest at shoulder; spire short, pleural angle 73°; anterior siphon broken, extant part straight. Growth line with strong U-shaped notch at suture, projected to form antisinus on ramp, looped into sinus across shoulder nodes, projected into minor antisinus, and nearly straight across whorl side. Suture collared, suture waved by attachment anteriorly to second noded spiral of shoulder. Sculpture of 12 short, strong axial costae extending from shoulder to fourth strong spiral cord; spiral cords narrower than axial costae, bearing large nodes at intersections with costae; spirals one and two closely spaced at shoulder, making double node at shoulder; spirals two, three, and four widely spaced; interspace between fourth cord and fifth cord broad, three times width of third interspace, with narrow sunken spiral band adjacent to fifth cord; fifth cord narrower and more finely noded than previous four cords, aligned with posterior fold of columella; fifth interspace about four times width of fifth cord; sixth cord half as wide as fifth, ?noded. Aperture elongate, medially broadened; columella straight with two barely visible columellar folds posterior to the midpoint; inner lip lightly callused; parietal callus roundly expanded over face of ultimate whorl, but not overlapping suture, extending adapically to first row of shoulder nodes.

Etymology.—Greek, *antherene*, wild bee, for Bee Canyon.

Type.—Holotype, an incomplete specimen LACMIP 13226 from LACMIP 8182, area 18.

Occurrence.—Early late Campanian, *Metaplasticeras pacificum* Zone with *Baculites rex*. One specimen, (Area 18) Williams Formation, Pleasants Sandstone Member.

Discussion.—The species is based on an incomplete specimen missing its early whorls, the end of the anterior siphon, and the outer lip. Surface preservation is poor; traces of fine spirals are not clear enough to describe; the nodes are worn; greatest diameter might be at either the shoulder or the third spiral cord. This singular specimen resembles *V. santana* in having a double row of nodes at the shoulder, but *V. antherena* is stouter with a stronger shoulder, has stronger sculpture with fewer axials and spirals. It resembles *V. magna* in having an apparently straight columella.

The species is questionably assigned to *Volutoderma* because there is no apparent expansion of the parietal callus apexward across the suture, and thus no corrugated subsutural deposit along the suture as in all other species of *Volutoderma* and in *Longoconcha*. Absence of the subsutural deposits may result from poor preservation. The growth line of *V. antherena* differs from that of *Volutoderma* in having, in addition to a sinus adjacent to the suture, an antisinus and a second shallow sinus at the second spiral cord.

VOLUTODERMA? new species

Figure 7.5, 7.6

Description.—Shell medium-sized, fusiform?, consisting of five or six round-shouldered whorls, spire short, suture on second row of nodes, one row of nodes showing on penultimate whorl, steep subsutural collar with two spirals at suture followed by narrow concave ramp. Pleural angle about 55°. Sculpture of at least 12 rounded costae and narrower spiral cords forming strong nodes at intersections, subsutural collar with two weakly noded cords, five to six strongly noded cords over inflated part of whorl, weaker cords on siphonal area. Aperture elongate, parietal shield roundly expanded on face of last whorl.

Type.—Hypotype LACMIP 13217 from LACMIP loc. 30141.

Occurrence.—Early late Maastrichtian. One small specimen, Area 13, LACMIP loc. 30141, Bee Rock Road, about a mile north of Nacimiento River, San Luis Obispo County, California; El Piojo Formation.

Discussion.—The specimen is small, incomplete, somewhat post-depositionally flattened, and has about six whorls, the last one with about five noded spirals. Distortion has doubtless made the diameter measurement of 30.5 mm uncharacteristically large. At least one fold can be discerned on the columella. The anterior siphon is broken off, and the specimen is too incomplete to serve as a holotype. One noded spiral cord shows on the penultimate spire whorl adapical to the suture. Two cords are at the suture on the subsutural collar, but *Volutoderma?* has a concave ramp that was probably without cords. Absence of ramp cords is more typical of *Retipirula* than of *Volutoderma*, but the slightly elongated whorl shape is more similar to *Volutoderma*. A larger, more mature, more complete specimen of this species might make its referral to a genus more positive. Taliaferro (1944) listed *Volutoderma averilli* from his Asuncion Formation on the north side of the Nacimiento River. Unfortunately, no specimen bearing any resemblance to *V. averilli*, *Volutoderma?*, or *Retipirula* from the Asuncion Formation was found in the UCMP or the CAS collections. *Coralliochama* sp. was collected from the El Piojo Formation at UCMP A-3435.

Volutoderma? is less elongate and has more convexly rounded early whorls, and a less aciculate, shorter spire, than *V. perissa* n. sp. *Volutoderma?* differs from *Retipirula calidula* n. sp. in having a relatively narrower, higher spire, a less roundly inflated body whorl, a less concave ramp, fewer costae on the last whorl, and weaker nodes on the subsutural collar. Whereas the whorl periphery of *Volutoderma?* is nearer the strong cord bounding the abapical margin of the ramp, the periphery of *R. calidula* is more abapically placed near the second strong cord from the ramp.

Volutoderma? occurs with *Turritella chaneyi* C. W. Merriam, 1941, and *T. webbi paynei* Saul, 1983, whereas *Retipirula calidula* n. sp. occurs with *T. chaneyi orienda* Saul, 1983 and *T. webbi* Saul, 1983.

Genus LONGOCONCHA Stephenson, 1941

Type species.—By original designation *Volutoderma tennesseensis* Wade, 1926; Tennessee, Ripley Formation, Late Campanian.

Diagnosis.—Elongate, narrowly fusiform to subcylindrical volutodermine volutes, having flanks of mature whorls flattened, spire high and evenly tapering. Three or more strong columellar folds.

Occurrence.—Late Cretaceous, Coniacian to Maastrichtian; North America: Atlantic and Gulf Coast of the United States; Europe: Germany and France; North Africa (Sohl, 1964); ?India (?*Volutoderma* of Bandel, 2000); southern California and northern Baja California, Mexico (herein).

Discussion.—The Texas species, *Volutulithes navarroensis*, was placed by Stephenson (1941) in *Longoconcha*, which he defined as a new subgenus of *Volutoderma*, differing from *Volutoderma* in being markedly more slender, having the flanks of the mature whorls conspicuously more flattened, and the ultimate whorl less constricted anteriorly. Stephenson designated *Volutoderma tennesseensis*, as type species and included *V. navarroensis*, *Volutoderma protracta* Dall, 1907, and *Volutoderma (Longoconcha) dalli* Stephenson, 1941 in *Longoconcha*.

Sohl (1964) listed species of *Longoconcha*, eight from Gulf and Atlantic states and two from Europe and Africa. All of his listed species are Campanian and Maastrichtian, but he mentioned an undescribed Coniacian species from the Eutaw Formation of Alabama. He considered *Longoconcha* to be generically distinct from *Volutoderma*, and in so doing reduced the geographic range of *Volutoderma* to the eastern Pacific. We concur with Sohl in considering *Volutoderma* to be endemic to the eastern Pacific.

In addition to its elongate, slim, subcylindrical shell with an evenly tapering spire, and very elongate, rather flat-sided whorls, *Longoconcha* commonly has spiral cords that are not noded and some species have secondary as well as primary spiral sculpture on the body whorl. Some features considered characteristic of *Longoconcha* by Sohl (1964), a slender spire, axial ribs present on spire but absent on the mature whorls, and spiral cords not noded on mature whorls, are variously present in Pacific Slope *Volutoderma* species. The characters currently considered diagnostic of *Volutoderma*, a corrugated subsutural collar, strong, raised imbricate growth lines, a heavy inner lip callus, and a spreading parietal callus shield are based on better preserved material not then available to Sohl. Only one Pacific slope species, *Longoconcha eumeka* n. sp., is here included in *Longoconcha*.

LONGOCONCHA EUMEKA new species

Figure 11.1–11.8

? *Volutoderma* cf. *magna* Packard. ANDERSON AND HANNA, 1935, p. 26.

Diagnosis.—Strongly sculptured *Longoconcha* with well-developed, rounded shoulders. Spire scaliform, penultimate whorl with three or four strong spiral cords exposed posterior to suture.

Description.—Shell elongate fusiform to subcylindrical with high turriculate spire and rather flat-sided adult whorls; ratio diameter to height nearly one to four; anterior siphon barely? deflected to left; ramp crossed by raised traces of outer lip, forming U-shaped posterior notch as it joins parietal shield. Spire less than one third of shell height; pleural angle of young specimen 30°, of mature specimen 39°. Protoconch (?) of two and a half whorls, turbinate, diameter 0.4 mm, ?smooth. Growth line with U-shaped sinus at suture becoming nearly vertical across body of whorl, turning slightly opisthocline? on the siphonal area. Mature sculpture of about 14 round-topped axial costae, weak across ramp, strongest at fifth to sixth spiral and becoming weak at eleventh to twelfth spiral, lacking or nearly so on siphonal area. Immature sculpture of spire whorls three to eleven: whorls three to six with strong arcuate (?collabral) costae and only faint indications of spirals; whorl seven with weak spirals; whorl eight with strongest spiral near midwhorl; whorl 11 with fourth and/or fifth spiral from suture strong, spirals one and two forming low collar, spiral three weak, spirals five or six to eleven or twelve forming nodes over axials, five or six weaker spirals on siphonal area. Aperture elongate, slightly wider posteriorly; outer lip edge denticulated at spiral cords, reflected to form narrow rim, extended posteriorly to form U-shaped trough extending above suture and joining parietal callus. Columella straight, callused, with three strong folds; parietal shield expanded over about two thirds of face of body whorl and extending posteriorly beyond aperture, in some specimens as far as suture.

Etymology.—Greek, *eumekes*, of good length.

Types.—Holotype SDNHM 70974 from SDNHM 3405. Paratypes SDNHM 25952 from SDNHM 3162-L; SDNHM 33992 from SDNHM 3405; SDNHM 70968 from SDNHM 4071; SDNHM 25959 from SDNHM 3162-C; LACMIP 13215 from LACMIP 23268; LACMIP 13216 from LACMIP 5991.

Material examined.—Fifty-seven specimens; one specimen from Area 11 upper Panoche Formation near Coalinga; three specimens from Area 16, at "Lang Ranch"; fifteen specimens from Area 19; two specimens from Area 21; thirty-six specimens from Area 22 (one from Cañon San Fernando, thirty-five from Arroyo Santa Catarina).

Occurrence.—Late late Campanian to early Maastrichtian, zones of *Baculites lomaensis*, *Baculites occidentalis*, and *Pachydiscus catarinae*. (Area 11) type locality of *Baculites lomaensis* Panoche Formation, "Ragged Valley Shale" = Uhalde Shale Member; lower Maastrichtian. (Area 16) upper Chatworth Formation. (Area 19) Point Loma Formation. (Areas 21 and 22) Rosario Formation. (Areas 16, 18, 19, 21, and 22) are upper upper Campanian?/lower Maastrichtian.

Discussion.—Although this species has a high spire, its suture is not descending as in *V. angelica* n. sp., *V. magna*, and *V. ynezae* n. sp., nor is the spire as aciculate as that of *V. perissa* n. sp.

Early whorls of *L. eumeka* have predominantly strong axial sculpture without the overriding spirals common on species of *Volutoderma*. Axial sculpture is dominant on about six juvenile whorls of *V. averillii*, but dominant on 11 of *L. eumeka*. *Volutoderma blakei* n. sp. and *V. elderi* n. sp. also have fewer juvenile whorls with dominantly axial sculpture than *L. eumeka*. *Longoconcha eumeka* resembles Gulf Coast species of the genus in shape and mature sculpture, but neither Stephenson (1941) nor Sohl (1964) mentioned such early sculpture on *Longoconcha* species, and their illustrations do not enlarge the early whorls to show details. *Longoconcha eumeka* has a greater height to diameter ratio than any Pacific slope *Volutoderma*. In general aspect, *L. eumeka* is more similar to the Gulf Coast species of *Longoconcha* than to *Volutoderma* and is more similar to the type species *L. tennesseensis* than to *L. navarroensis* largely because of the strength of the sculpture. *Longoconcha eumeka* differs from *L. tennesseensis* in commonly having more costae per whorl and in having them more persistent on mature specimens. On *L. eumeka*, secondary cords are generally not developed between the primary cords of the ultimate whorl. *Longoconcha dalli* Stephenson, 1941, Navarro Formation, Kemp Clay, Texas, *L. protracta* Dall, 1907, Ripley Formation, Alabama, and *L. quadrilirata* (Sohl, 1964) Ripley Formation, Mississippi, also lack secondaries. On some specimens of *L. eumeka* the ultimate whorl is "fimbriated" by remains of reflexed outer lip edges, which obscure the normal sculpture.

Longoconcha eumeka has been found over a greater geographical range than other species described herein, with the exception of *V. averillii*. *Longoconcha eumeka* is also fairly common in some areas. That it has not previously been described may result from the poor preservation of many of the specimens. In collections, *L. eumeka* has commonly been misidentified as a *Volutoderma*, and if assigned to a species, assigned either to *V. averillii* or *V. magna*.

Genus RETIPIRULA Dall, 1907

Type species.—*Turbinella crassitesta* Gabb, 1869 (p. 157, pl. 26, fig. 37) by original designation Dall, 1907, p. 5, fn 2. Gabb's specimens were from "near Martinez," Contra Costa County, northern California.

Diagnosis.—Pyriiform volutes with axial and spiral ribbing forming nodes at their intersections and having two to four columellar plaits. Aperture extended posteriorly into rounded posterior notch at suture; inner lip callus expanded, forming parietal shield on whorl face extending adapically over previous suture. Protoconch turbinate.

Occurrence.—Lake County, northern California to Baja California, Mexico; latest Cretaceous and early late Paleocene.

Discussion.—*Retipirula* was considered to be a subgenus of *Volutocorbis* Dall, 1890, by Wenz (1943) and Zinsmeister (1977), but the more U-shaped posterior notch of the aperture, the deposits extending posteriorly onto the previous whorl and obscuring the suture and part of the previous whorl, and the parietal shield are suggestive of a relationship to *Volutoderma* rather than to *Volutocorbis*. Additionally, the anterior sinus of *Retipirula* is flexed leftward (apertural view), but the sinus of *Volutocorbis* is essentially straight. The size difference between the strongly noded spiral cords of the whorl face and the finer smoother cords of the siphonal area is more distinct in *Retipirula* than in *Volutoderma* or in *Volutocorbis*, in both of which the change from coarse to fine is more gradational. *Retipirula* lacks the spiral cordlets commonly present on the ramp of *Volutoderma*.

Stewart (1927) chose a lectotype for the type species of *Retipirula* from among Gabb's three incomplete specimens at the Academy of Natural Sciences Philadelphia. Gabb's figure is a synthetograph, and as can be seen by comparison with Figure 11.11 and 11.12 and Zinsmeister's illustration (1977, p. 178, text-fig. 1d), Gabb drew a more posteriorly sharply ovoid aperture rather than the somewhat rounder, posteriorly more broadly

notched, and more heavily callused aperture of *Retipirula*. Gabb's specimens came from "near Martinez," and subsequent specimens from that vicinity have been collected from the lower Vine Hill Formation below the occurrence of P4 Zone foraminifers (C. E. Weaver, 1953, chart, p. 28–29). Occurrences of *R. crassitesta* in the Santa Susana Formation of the Simi Hills, Ventura County, California, are also in lower Selandian strata below those yielding foraminifera of the P4 Zone.

With recognition of *R. calidula* n. sp. and *R. pinguis* n. sp., *Retipirula* is no longer a monotypic genus. Its range is extended to include the youngest Cretaceous.

RETIPIRULA CALIDULA new species

Figure 11.9, 11.10

Diagnosis.—Relatively high-spired *Retipirula* with cords on siphonal area only slightly smaller than those of whorl face.

Description.—Shell medium-sized, pyriform, consisting of about five roundly expanded whorls, diameter to height ratio about 2.4; spire short with two rows of nodes on each whorl; suture on second row of nodes; ramp steep, concave. Pleural angle about 70°. Sculpture of about 12 rounded axial ribs and narrower spiral cords forming strong nodes at intersections; subsutural collar strongly noded; ramp without cords, five to six strongly noded cords over inflated part of whorl, weaker cords on siphonal area. Aperture elongate; outer lip forming U-shaped notch at suture, covering suture, and extending to posterior noded spiral of previous whorl, parietal shield roundly expanded on face of ultimate whorl. Columella with at least two folds.

Etymology.—Latin, *calidula*, warm, referring to Warm Springs Mountain.

Types.—Holotype LACMIP 13218 and paratypes 13219, 13220 from LACMIP loc. 21591.

Material examined.—Three specimens from LACMIP loc. 21591.

Occurrence.—Latest Maastrichtian. (Area 14) Warm Springs Mountain, basal San Francisquito Formation.

Discussion.—Available specimens of *R. calidula* are all crushed and consist of molds or casts, but the following differences between *R. calidula* and *R. crassitesta* (Fig. 11.11, 11.12) are apparent. *Retipirula calidula* is larger with one more strong, noded spiral cord about the body whorl, has a wider ramp and a higher spire, the whorls of which show one additional noded spiral. On the spire of *R. calidula* the suture is noticeably anterior to the most posterior of the strong spirals causing a tabulation, but on the spire of *R. crassitesta* the suture is closely adjacent to the second noded cord of the previous whorl. Perhaps because of its larger size and higher spire this new species bears a greater resemblance to *Volutoderma* than does *R. crassitesta*.

At its type locality on Warm Springs Mountain, *Retipirula calidula* occurs with *Turritella webbi* and *Turritella chaneysi orienda* (Fig. 2).

RETIPIRULA CRASSITESTA (Gabb, 1869)

Figure 11.11, 11.12

Turbinella crassitesta GABB, 1869, p. 157, pl. 26, fig. 37.

Retipirula crassitesta (Gabb, 1869), DALL, 1907, p. 50; WARING, 1917, p. 83, pl. 13, figs. 1, 2; NELSON, 1925, p. 432, pl. 59, fig. 6a, 6b; STEWART, 1927, p. 406, pl. 25, fig. 11; CLARK, 1929, pl. 2, fig. 10 (reprint of Nelson, 1925). *Volutocorbis (Retipirula) crassitesta* (Gabb, 1869). WENZ, 1943, fig. 3,749 (reprint of Stewart, 1927). ZINSMEISTER, 1977, p. 177, text-fig. 1, a–d. ZINSMEISTER AND PAREDES-MEJIA, 1988, pl. 2, fig. 6.

Diagnosis.—Medium-sized, pyriform shell with moderately elevated, tabulated spire. Whorls roundly inflated with sculpture of six to seven spiral cords strongly noded at intersections with collabral costae; distinctly finer unnoded cords on siphonal neck.

Description.—Shell medium-sized, pyriform; spire moderately elevated, tabulated by ramp; ramp slightly concave, barely sloping, bordered distally by noded cord; whorl side roundly inflated, periphery at third to fourth noded cord, roundly constricted to broad siphonal neck; suture at second noded cord of previous volution; growth line opisthocline across ramp to first row of nodes, barely flexed across whorl flank. Sculpture of spiral cords and collabral costae, narrower than interspaces, six to seven cords, strong and sharply noded at intersections with costae; up to 10 finer unnoded cords on siphonal neck; costae weaker than cords, extending from first row of nodes to fifth or sixth row. Aperture elongate, widest medially, with slight posterior notch and moderate width anterior canal; outer lip evenly curved; inner lip callused with

parietal callus expanded apically up to first cord of previous whorl; three oblique columellar folds near base of columella, two nearly equal, moderately strong, and apically one much weaker fold.

Types.—Hypotype LACMIP 13221 from LACMIP 22307.

Type locality.—"Martinez," Contra Costa County, California.

Material examined.—More than 200 specimens from the lower Santa Susana Formation, Simi Valley and Simi Hills, Area 16, Ventura County, California.

Occurrence.—Late early and early late Paleocene. Clear Lake, Lake County, "Martinez," Paleocene. Vicinity of Martinez, Contra Costa County, lower Vine Hill Formation; upper lower Paleocene. Rock Creek, Valerino area, Los Angeles County, San Francisquito Formation; upper lower Paleocene. (Area 14) San Francisquito Formation; lower Paleocene. (Area 16) Santa Susana Formation; upper lower Paleocene. (Area 17) Santa Susana Formation; upper lower Paleocene. (Area 22) south of Arroyo Santa Catarina (LACMIP loc. 26364), Sepultura Formation; upper lower Paleocene.

Discussion.—As noted by Zinsmeister (1977), *R. crassitesta* resembles *Volutocorbis virginiae* Zinsmeister, 1977, but the sculpture on whorls of *R. crassitesta* is coarser and becomes abruptly finer on the siphonal neck, its columella is more flexed, and its anterior siphon more strongly deflected leftward. A fortuitously broken specimen that shows the columella well back from the aperture has three folds on the columella, two strong and one weak.

RETIPIRULA PINGUIS new species

Figure 11.13, 11.14

Diagnosis.—*Retipirula* with very low spire.

Description.—Shell medium-sized, roundly ovoid, flattened apically and elongated abapically by short, slightly twisted, anterior siphonal neck; spire barely elevated; suture undulated on first noded spiral cord of previous whorl; whorls expanding flatly; ramp barely concave and slightly sloped to first noded cord; whorl side roundly inflated, widest medially between third and fourth cord, constricted into broad siphonal neck. Protoconch turbinata, of three whorls. Growth line opistocyt across ramp, slightly flexed across whorl flank, occasionally strong enough to disrupt costae. Sculpture of spiral cords and collabral costae; spiral cords narrower than interspaces, about seven strong spirals sharply noded at intersections with costae, eight (?) finer unnoded cords on siphonal neck; collabral costae extending from first noded cord to seventh cord. Aperture elongate, widest medially with slight posterior siphonal notch and moderate width anterior canal; outer lip evenly curved; inner lip callused with parietal callus expanded apically to first cord of previous whorl in early whorls and across ramp of previous whorl in mature specimens; four oblique folds near base of columella, two nearly equal, moderately strong, near base of columella, one weak, and one very weak anterior fold.

Etymology.—Latin, *pinguis*, fat.

Types.—Holotype LACMIP 13222 from LACMIP 22701, Area 16. Paratypes 13223, 13224 from LACMIP 22701; 13225 from LACMIP 22574, Area 16.

Material examined.—Twenty-five specimens from Area 16; six specimens from redeposited cobble in Santa Susana Formation conglomerate bed on north side Simi Valley (LACMIP 7142), Santa Susana Mountains; nineteen specimens from Simi Hills, south side of Simi Valley.

Occurrence.—Early Paleocene. (Area 16) Santa Susana Mountains and Simi Hills, lower Santa Susana Formation; lower Paleocene.

Discussion.—*Retipirula pinguis* resembles *R. crassitesta* in overall sculpture and shape, but the whorl of *R. pinguis* is more roundly inflated posteriorly with more, narrower cords and costae, the intersections of which are marked by smaller nodes than on *R. crassitesta*. Whorls of *R. pinguis* overlap the posterior row of nodes of the previous whorl and have a broad, almost horizontal ramp which results in a shorter spire with a much wider pleural angle than in *R. crassitesta*.

The occurrence of *R. pinguis* with *Turritella reversa* Waring, 1917, in redeposited boulders in Santa Susana Formation conglomerates, suggests that *R. pinguis* is of early Paleocene age.

ACKNOWLEDGMENTS

We thank the members of the Victoria Paleontology Society, Saanichton, British Columbia, for sending us excellent specimens with good locality data from the Nanaimo Group, Georgia Basin, British Columbia; B. J. Dougherty of the Geological Survey of Canada for the loan of Whiteaves' specimens from the Nanaimo Group, Sucia Island, which became the type lot for *V. suciana* Dall, 1907; D. L. Lindberg of the University of California, Berkeley,

for the loan of types of *Volutoderma santana* Packard and *V. magna*; T. A. Deméré of the Natural History Museum of San Diego for the loan of excellent specimens of volutes from the Point Loma Formation; J. D. Cooper of California State University, Fullerton, for the loan of Orange County specimens; J. DeMouthe of California Academy of Sciences for access to CASG collections and loan of specimens; M. Kooser of University of California, Riverside for access to UCR collections and loan of specimens; R. Hilton of Sierra College for field assistance in the Pentz area and E. Göhre for field assistance and loan of specimens; L. Groves of Malacology at the Natural History Museum of Los Angeles County for loans and library assistance. We are also indebted to the late H. G. Richards of the Academy of Natural Sciences of Philadelphia for the loan of Gabb's specimens identified as *Volutolithes navarroensis* Shumard; and the late J. W. Durham and J. H. Peck of the University of California, Berkeley for the opportunity to study and make plaster casts of the type specimens of *Fusus averillii* Gabb, *Volutoderma santana* Packard, and *V. magna* Packard. We thank P. D. Ward for correcting ranges of ammonites, especially *Baculites* associated with *Volutoderma*.

REFERENCES

- ABBASS, H. L. 1963. A monograph on the Egyptian Cretaceous gastropods. United Arab Republic, Geological Survey and Mineral Research Department, Palaeontological Series, Monograph 2, 146 p.
- ANDERSON, F. M. 1902. Cretaceous deposits of the Pacific Coast. California Academy of Sciences Proceedings, series 3, 2:1–154, pls. 1–12.
- ANDERSON, F. M. 1958. Upper Cretaceous of the Pacific Coast. Geological Society of America Memoir 71, 378 p., 75 pls.
- ANDERSON, F. M. AND G. D. HANNA. 1935. Cretaceous geology of Lower California. California Academy of Sciences, Proceedings, series 4, 23(1): 1–34.
- ARNOLD, R. 1908. Descriptions of new Cretaceous and Tertiary fossils from the Santa Cruz Mountains, California. U.S. National Museum Proceedings, 34:345–390, pls. 31–37.
- ARNOLD, R. 1910. Paleontology of the Coalinga District, Fresno and Kings Counties, California. United States Geological Survey Bulletin, 396, 173 p., 30 pls.
- ARNOLD, R. AND R. ANDERSON. 1910. Geology and oil resources of the Coalinga District, Fresno and Kings counties, California. United States Geological Survey Bulletin, 398, 354 p., 52 pls.
- BANDEL, K. 2000. Some gastropods from the Trichinopoly Group Tamil Nadu, India and their relation to those from the American Gulf Coast. Geological Society of India Memoir, 46:65–111.
- BANDEL, K. 2003. Cretaceous volutid Neogastropoda from Western Desert of Egypt and their place within the Neogastropoda (Mollusca). Mitteilungen Geologisch-Paläontologisches Institut Universität Hamburg, 87:73–98.
- BANDEL, K. AND D. T. DOCKERY. 2001. The Sarganidae (Pyrifusoidea, La-trogastropoda), their taxonomy and paleobiogeography. Journal of the Czech Geological Society, 46:335–351.
- BOLTON, T. E. 1965. Catalogue of type invertebrate fossils of the Geological Survey of Canada. Volume II. Geological Survey of Canada, Ottawa, 344 p.
- BOUCHET, P., J. FRYDA, B. HAUSDORF, W. PUNDER, Á. VALDÉS, AND A. WARÉN. 2005. Working classification of the Gastropoda, p. 239–284. In P. Bouchet and J.-P. Rocroi (eds.), Classification and Nomenclator of Gastropod Families. Malacologia, 47(1–2).
- BRÉBION, P. 1956. Gasteropoda Crétacés du Bas-Congo, recoltés par H. Pierard. Annales du Musée Royal du Congo Belge Tervuren, Belgique, Sciences Géologiques, 17:81–93.
- BUTLER, R. F., G. E. GEHRELS, AND K. P. KODAMA. 2001. A moderate translation alternative to the Baja British Columbia hypothesis. GSA Today, 11(6):4–9.
- CLARK, B. L. 1929. Stratigraphy and faunal horizons of the Coast Ranges of California. Privately published, Berkeley, 132 p., 50 pls.
- COBBAN, W. A. AND W. J. KENNEDY. 1995. Maastrichtian ammonites chiefly from the Prairie Bluff Chalk in Alabama and Mississippi. The Paleontological Society Memoir 44 (Journal of Paleontology 69, no. 5 supplement III of III), 40 p.
- COSSMANN, M. 1896. Essais de Paléontologie Comparé. Privately published, Paris, tome 2, 176 p.
- COSSMANN, M. 1909. Essais de Paléontologie Comparé. Privately published, Paris, tome 8, 248 p.
- CUVIER, G. L. C. 1797. Tableau Élémentaire de l'Histoire Naturelle des Animaux [des Mollusques]. Baudouin, Paris, 710 p.
- DAILEY, D. H. AND W. P. POPENOE. 1966. Mollusca from the Upper Cretaceous Jalama Formation, Santa Barbara County, California. University of California Publications in Geological Sciences, 65, 41 p.
- DALL, W. H. 1890. Contributions to the Tertiary fauna of Florida with especial reference to the Miocene Silex beds of Tampa and the Pliocene beds of the Caloosahatchie River, Pt. 1. Wagner Free Institute of Science Transactions, 3:1–200.
- DALL, W. H. 1907. Notes on some Upper Cretaceous Volutidae, with descriptions of new species and a revision of the groups to which they belong. Smithsonian Miscellaneous Collections, 50 (no. 1704):1–23.
- DARRAGH, T. A. AND W. F. PONDER. 1998. Family Volutidae, p. 833–835. In P. L. Beesley, G. J. B. Ross, and A. Wells (eds.), Mollusca: The Southern Synthesis. Fauna of Australia, 5. CSIRO Publishing, Melbourne.
- DARTVELLE, E. AND P. BRÉBION. 1956. Mollusques fossiles du Crétacé de la Côte occidentale d'Afrique du Cameroun à l'Angola. Annales du Musée Royal du Congo Belge Tervuren, Belgique, Sciences Géologiques, 17:viii + 1–128, pls. 1–8.
- DICKINSON, W. R. AND R. F. BUTLER. 1998. Coastal and Baja California paleomagnetism reconsidered. Geological Society of America Bulletin, 110: 1268–1280.
- DJALILOV, M. R. 1977. Cretaceous Gastropods from Southeastern Central Asia. Akademia Nauk Tadzhiik SSR, 202 p. (In Russian)
- D'ORBIGNY, A. 1842–1843. Paléontologie Française. Ser. 1. Animaux Invertébrés. Terrain Crétacés, 2, 456 p.
- DZHALILOV, 1977, see Djalilov
- ELDER, W. P. AND L. R. SAUL. 1993. Paleogeographic implications of molluscan assemblages in the Upper Cretaceous (Campanian) Pigeon Point Formation, California, p. 171–186. In G. Dunne and K. McDougall (eds.), Mesozoic Paleogeography of the Western United States-II, Pacific Section SEPM, Book 71.
- ELDER, W. P. AND L. R. SAUL. 1996. Taxonomy and biostratigraphy of Coniacian through Maastrichtian *Anchura* (Gastropoda: Aporrhaidae) of the North American Pacific Slope. Journal of Paleontology, 70:381–399.
- ELDER, W. P., L. R. SAUL, AND C. L. POWELL. 1998. Late Cretaceous and Paleocene molluscan fossils of the Gualala Block and their paleogeographic implications, p. 149–168. In W. P. Elder (ed.), Geology and Tectonics of the Gualala Block, Northern California, Pacific Section SEPM, Book 84.
- FISHER, W. L., P. U. RODDA, AND J. W. DIETRICH. 1964. Evolution of *Athleta perrosa* stock (Eocene, Gastropoda) of Texas. University of Texas Publications, 6413, 117 p.
- FORBES, E. 1846. Report on the Cretaceous fossil invertebrates from southern India, collected by Mr. Kaye and Mr. Cunliffe. Geological Society of London Transactions [1845], series 2, 7:97–174.
- FOWKES, E. J. 1982. An Educational Guidebook to the Geologic Resources of the Coalinga District, California. West Hills College, Coalinga, California, 260 p.
- FRAKES, L. A. 1999. Estimating the global thermal state from Cretaceous sea surface and continental temperature data, p. 49–57. In E. Barrera and C. C. Johnson (eds.), Evolution of the Cretaceous Ocean-Climate System. Geological Society of America Special Paper 332.
- GABB, W. M. 1864. Description of the Cretaceous fossils. California Geological Survey, Palaeontology, 1:57–243, pls. 9–32.
- GABB, W. M. 1866–1869. Cretaceous and Tertiary fossils. California Geological Survey, Palaeontology, 2, 299 p.
- GABB, W. M. 1877. Notes on American Cretaceous fossils, with descriptions of some new species. Academy of Natural Sciences of Philadelphia Proceedings, 28:276–324.
- GRADSTEIN, F. M., J. G. OGG, AND A. SMITH. 2004. Geologic Time Scale. Cambridge University Press, Cambridge, 589 p.
- HAGGART, J. W. 1986. Stratigraphy of the Redding Formation of North-Central California and its bearing on Late Cretaceous paleogeography, p. 161–178. In P. L. Abbott (ed.), Cretaceous Paleogeography and Stratigraphy. SEPM Pacific Section, symposium volume, Book 46.
- HAGGART, J. W. AND P. D. WARD. 1984. Late Cretaceous (Santonian–Campanian) stratigraphy of the northern Sacramento Valley, California. Geological Society of America Bulletin, 95:618–627.
- HOLZAPFEL, E. 1888–1889. Die Mollusken der Aachener Kreide. Palaeontographica, 34:29–80 (1888); 35:139–268 (1889).
- ICZN. 1999. International Commission on Zoological Nomenclature. International Code of Zoological Nomenclature, (fourth edition). International Trust for Zoological Nomenclature, c/o Natural History Museum, London, 306 p.
- JOHNSON, C. C. 1999. Evolution of Cretaceous surface current circulation patterns, Caribbean and Gulf of Mexico, p. 329–343. In E. Barrera and C. C. Johnson (eds.), Evolution of the Cretaceous Ocean-Climate System. Geological Society of America Special Paper 332.
- KIEL, S. AND K. BANDEL. 2003. New taxonomic data for the gastropod fauna of the Umzamba Formation (Santonian–Campanian, South Africa) based on newly collected material. Cretaceous Research, 24:449–475.
- KIEL, S. AND F. J. KRÜGER. 2006. Gastropoda aus dem Mittelsanton (Oberkreide) von Lengede (Niedersachsen). Braunschweiger Naturkundliche Schriften, 7:677–696.
- KIEL, S. AND M. C. PERRILLIAT. 2004. New gastropods from the Maastrichtian of the Mexcala Formation in Guerrero, southern Mexico, Pt. III, Higher Caenogastropoda. Neues Jahrbuch für Geologie und Paläontologie, Abhandlungen, 231:191–217.
- KODAMA, K. P. AND P. D. WARD. 2001. Compaction-corrected paleomagnetic

- paleolatitudes for Late Cretaceous rudists along the Cretaceous California margin: Evidence for less than 1500 km of post-Late Cretaceous offset for Baja British Columbia. *Geological Society of America Bulletin*, 113:1171–1178.
- KOLLMANN, H. A. 2005. Révision critique de la Paléontologie Française d'Alcide d'Orbigny, Volume III Gastropodes Crétacés. Publication dirigée par J.-C. Fischer. *Museum d'Histoire naturelle de Vienne, Autriche*, Backhuys Publishers, Leiden, Pays-Bas, France, 239 p. + Réédition du Tome Second (Terrains crétacés, Gastropodes de la Paléontologie française d'Alcide d'Orbigny (édition original: 1842–1843) 454 p. + Atlas of pl. 149–236.
- LUDVIGSEN, R. AND G. BEARD. 1994. *West Coast Fossils*. Whitecap Books, Vancouver/Toronto, 194 p.
- LUDVIGSEN, R. AND G. BEARD. 1997. *West Coast Fossils* (second edition). Harbour Publishing, Madiera Park, British Columbia, 216 p.
- LUND, S. P. AND D. J. BOTTJER. 1992. Paleomagnetic evidence for microplate tectonic development of southern and Baja California, p. 231–248. *In* J. P. Dauphin and B. R. T. Simoneit (eds.), *The Gulf and Peninsular Province of the Californias*. American Association of Petroleum Geologists Memoir 47.
- MATSUMOTO, T. 1959a. Upper Cretaceous ammonites of California, Pt. I. *Kyushu University Faculty of Science Memoir, series D, Geology* 8:91–171.
- MATSUMOTO, T. 1959b. Upper Cretaceous ammonites of California, Pt. II. *Kyushu University Faculty of Science Memoir, series D, Geology, Special Volume 1*, 172 p.
- MATSUMOTO, T. 1960. Upper Cretaceous ammonites of California, Pt. III. *Kyushu University Faculty of Science Memoir, series D, Geology, Special Volume 2*, 204 p., 2 pls.
- MATSUMOTO, T. 1984. Some ammonites from the Campanian (Upper Cretaceous) of Northern Hokkaido. *Palaeontological Society of Japan Special Papers*, 27, 93 p.
- MATSUMOTO, T. AND I. OBATA. 1963. A monograph of the Baculitidae from Japan. *Kyushu University, Faculty of Science, Memoirs, series D, Geology*, 13:1–116.
- MEEK, F. B. 1858. Descriptions of new organic remains from the Cretaceous rocks of Vancouver's Island. *Albany Institute, Transactions*, 4:37–49 (1857).
- MEEK, F. B. 1862. Descriptions of new Cretaceous fossils collected by the North-Western Boundary Commission, on Vancouver and Suia Islands. *Academy of Natural Sciences of Philadelphia Proceedings*, 13:314–318.
- MERRIAM, C. W. 1941. Fossil turritellas from the Pacific Coast region of North America. *University of California Publications, Department of Geological Sciences Bulletin*, 16:1–214.
- MERRIAM, J. C. 1895. A list of type specimens in the Geological Museum of the University of California, etc. *University of California, Geology Department*, 3 printed, unnumbered pages. [Reprinted by A. W. Vogdes, 1896, *California State Mining Bureau, Bulletin 10*, and 1904, *California State Mining Bureau, Bulletin 3*.]
- MORTON, B. 1986. The diet and prey capture mechanism of *Melo melo* (Prosobranchia: Volutidae). *Journal of Molluscan Studies*, 51:156–160.
- MORTON, S. G. 1834. Synopsis of the Organic Remains of the Cretaceous Group of the United States. Illustrated by nineteen plates, to which is added an appendix containing a tabular view of the Tertiary fossils discovered in America. Key and Biddle, Philadelphia, 88 p.
- MÜLLER, J. 1851. Monographie der Petrefacten der Aachener Kreideformation, Pt. 2. J. J. Beaufort, Bonn, 88 p.
- NELSON, R. N. 1925. A contribution to the paleontology of the Martinez Eocene of California. *University of California Publications, Department of Geological Sciences Bulletin* 15:397–466.
- NILSEN, T. H. 1984a. Tectonics and sedimentation of the Upper Cretaceous Hornbrook Formation, Oregon and California, p. 101–118. *In* J. K. Crouch and S. B. Bachman (eds.), *Tectonics and sedimentation along the California margin*. Society of Economic Paleontologists and Mineralogists, Pacific Section, Book 38.
- NILSEN, T. H. 1984b. Stratigraphy, sedimentology, and tectonic framework of the Upper Cretaceous Hornbrook Formation, Oregon and California, p. 51–88. *In* T. H. Nilsen (ed.), *Geology of the Upper Cretaceous Hornbrook Formation, Oregon and California*. Society of Economic Paleontologists and Mineralogists, Pacific Section, 42.
- PACKARD, E. L. 1922. New species from the Cretaceous of the Santa Ana Mountains, California. *University of California Publications, Department of Geological Sciences, Bulletin*, 13:413–462.
- PERVINQUIÈRE, L. 1912. Études de Paléontologie Tunisienne II, Gastropodes et lamellibranches des terrains Crétacés. *Direction Générale des Travaux Publics, Carte Géologique de la Tunisie, Paris*, 352 p. (Atlas), 23 pls.
- PLSBRYS, H. A. AND A. A. OLSSON. 1954. Systems of the Volutidae. *Bulletins of American Paleontology*, 35(152):275–306.
- PONDER, W. F. AND A. WARÉN. 1988. Appendix. Classification of the Caenogastropoda and Heterostropha—A list of the family-group names and higher taxa, p. 288–328. *In* W. F. Ponder (ed.), *Prosobranch Phylogeny*. Malacological Review, Supplement 4.
- POPENOE, W. P. 1937. Upper Cretaceous Mollusca from southern California. *Journal of Paleontology*, 11:379–402.
- POPENOE, W. P. 1942. Upper Cretaceous formations and faunas of Southern California. *American Association of Petroleum Geologists Bulletin*, 26:162–187.
- POPENOE, W. P. 1954. Mesozoic formations and faunas, southern California and northern Baja California, p. 15–21. *In* California Division of Mines, *Bulletin 170, Chapter 3, (Historical Geology)*.
- POPENOE, W. P. 1983. Cretaceous Aporrhaidae from California: Aporrhainae and Arrhoginae. *Journal of Paleontology*, 57:742–765.
- POPENOE, W. P. AND L. R. SAUL. 1987. Evolution and classification of the Late Cretaceous—early Tertiary gastropod *Perissitys*. *Natural History Museum of Los Angeles County, Contributions in Science*, 380, 37 p.
- RAFINESQUE, C. S. 1815. *Analyses de la Nature ou Tableau de l'Univers et des Corps Organisés*. Barravecchia, Palermo, 224 p.
- SAUL, L. R. 1973. Evidence for the origin of the Macridae (Bivalvia) in the Cretaceous. *University of California Publications in Geological Science*, 97, 59 p.
- SAUL, L. R. 1974. Described or figured West Coast species of *Cymbophora*. *Journal of Paleontology*, 48:1068–1095.
- SAUL, L. R. 1978. The North Pacific Cretaceous Trigonid Genus *Yaadia*. *University of California Publications in Geological Science*, 119, 65 p.
- SAUL, L. R. 1983. *Turritella* zonation across the Cretaceous–Tertiary boundary, California. *University of California Publications in Geological Science*, 125, 164 p.
- SAUL, L. R. AND J. M. ALDERSON. 1981. Late Cretaceous Mollusca of the Simi Hills: An introduction, p. 29–42. *In* M. H. Link, R. L. Squires, and I. P. Colburn (eds.), *Simi Hills Cretaceous turbidites*, Southern California. SEPM, Pacific Section, 1981 Fall Field Trip Guidebook. Los Angeles, California.
- SAUL, L. R. AND D. J. BOTTJER. 1982. Late Cretaceous megafossil locality map, northern Santa Ana Mountains, California, p. 77–79. *In* D. J. Bottjer, I. P. Colburn, and J. D. Cooper (eds.), *Late Cretaceous Depositional Environments and Paleogeography, Santa Ana Mountains, Southern California*. SEPM, Pacific Section, Annual Convention Field Guidebook & Volume. Los Angeles, California.
- SAUL, L. R. AND W. P. POPENOE. 1962. *Meekia*, enigmatic Cretaceous pelecypod genus. *University of California Publications in Geological Science*, 40:289–344.
- SAUL, L. R. AND W. P. POPENOE. 1993. Additions to Pacific Slope Turonian Gastropoda. *The Veliger*, 36:351–388.
- SCHENCK, H. G. AND A. M. KEEN. 1940. *California Fossils for the Field Geologists*. Privately published and Stanford University Press, 86 p.
- SCHOELLHAMER, J. E., J. G. VEDDER, R. F. YERKES, AND D. M. KINNEY. 1981. Geology of the northern Santa Ana Mountains, California. *U.S. Geological Survey, Professional Paper 420-D*, 107 p.
- SHIMER, H. W. AND R. R. SHROCK. 1944. *Index Fossils of North America*. John Wiley and Sons, New York, 837 p.
- SHUMARD, B. F. 1861. Descriptions of new Cretaceous fossils from Texas. *Boston Society of Natural History Proceedings*, 8:188–205.
- SMITH, J. P. 1900. The development and phylogeny of *Placenticeras*. *California Academy of Sciences Proceedings, series 3*, 1:80–240.
- SOHL, N. F. 1964. Neogastropoda, Opisthobranchia and Basommatophora from the Ripley, Owl Creek, and Prairie Bluff formations. *U.S. Geological Survey Professional Paper*, 331-B:i–iv, 153–344, pls. 19–52.
- SOHL, N. F. 1987. Cretaceous Gastropods: Contrasts between Tethys and the temperate provinces. *Journal of Paleontology*, 61:1085–1111.
- SPATH, L. F. 1926. On new ammonites from the English Chalk. *Geological Magazine*, 63:77–83.
- SQUIRES, R. L. 1981. Geologic map of Upper Cretaceous Chatsworth Formation, Simi Hills, California (with fossil localities), map in pocket. *In* M. N. Link, R. L. Squires, and I. P. Colburn (eds.), *Simi Hills Cretaceous turbidites*, Southern California. SEPM, Pacific Section, Guidebook Fall Field Trip 1981. Los Angeles, California.
- SQUIRES, R. L. AND L. R. SAUL. 2006. Cretaceous *Acila* (*Truncacila*) (Bivalvia: Nuculidae) from the Pacific Slope of North America. *The Veliger*, 48(1):32–53.
- STEPHENSON, L. W. 1941. The larger invertebrate fossils of the Navarro Group of Texas. *University of Texas Publications*, no. 4101, 641 p.
- STEPHENSON, L. W. 1952. Larger invertebrate fossils of the Woodbine Formation (Cenomanian) of Texas. *U.S. Geological Survey, Professional Paper 242*, 226 p.
- STEWART, R. B. 1927. Gabb's California fossil type gastropods. *Academy of Natural Sciences of Philadelphia, Proceedings*, 78:287–447.
- STOLICZKA, F. 1867–1868. *Cretaceous Fauna of Southern India, 2. The Gastropoda*. *Geological Survey of India, Memoirs, Palaeontologia Indica, series 5*, 497 p.