



- Saul, L. R., 1983, Notes on Paleogene turritellas, venericardias, and molluscan stages of the Simi Valley area, California, in Squires, R. L., and Filewicz, M. V., eds., Cenozoic geology of the Simi Valley area, southern California: Pacific Section, Society of Economic Paleontologists and Mineralogists, Volume and Guidebook, p. 71-80.
- Squires, R. L., 1984, Megapaleontology of the Eocene Llajas Formation, Simi Valley, California: Los Angeles County Natural History Museum, Contributions in Science, Number 350, p. 1-76.
- \_\_\_\_\_, 1987, Eocene molluscan paleontology of the Whitaker area, Los Angeles and Ventura Counties, California: Los Angeles County Natural History Museum, Contributions in Science, Number 388, p. 1-96.
- \_\_\_\_\_, Eocene macropaleontology of northern Lockwood and Ventura County, California: Los Angeles County Natural History Museum, Contributions in Science (in press)

GASTROPODA	OLD STAGE RANGE	NEW STAGE RANGE
Amaurellina caleocia Vokes	D	C-D
Architectonica (Stellaxis) cognata Gabb	D	C-D
Bittium? dumblei (Dickerson)	D	C-D
Clavilithes tabulatus (Dickerson)	C	C-D
Conus caleocius Vokes	D	C-D
Conus remondii Gabb	D	C-D
Galeodea (Caligaleodea) californica Clark	D	C-D
Galeodea (Gomphopages) meganensis Vokes	M	M-C
Homalopoma umpquaensis domenginensis Vokes	D	C-D
Lyria andersoni Waring	D	C-D
Lyrischapa lajollaensis (Hanna)	D	C-D
Natica (Naticarius) uvasana Gabb	Tr-Te	C-Te
Olequahia domenginica (Vokes)	D	C-D
Paraserapha erraticus (Cooper)	D-Tr	C-Tr
Strepsidura ficus (Gabb)	D-Te	C-Te
Trypanotoma stocki (Dickerson)	D	C-D
Turritella andersoni Dickerson	Upper M?-C	C
Turritella buwaldana Dickerson	Upper M?, C?, D-Te	Upper M?, C-Te
Turritella uvasana hendoni s.l. Merriam	Mid. Ecc.	C-Mid. Ecc.
Xenophora stocki Dickerson	D	C-D

M = "Meganos Stage"; C = "Capay Stage"; D = "Domengine Stage"; Tr = "Transition Stage"; Te = "Tejon Stage".

Table 1. Refinements of the West Coast provincial molluscan stage ranges of 20 marine gastropods. These refinements are based on recent research by Squires (1984, 1987, in press).

BIVALVIA	OLD STAGE RANGE	NEW STAGE RANGE
Barbatia (Cucullaearpa) cliffensis Hanna	D	C-D
Claibornites diegoensis (Dickerson)	D	C-D
Crassatella uvasana Conrad	Te	D-Te
Corbula (Caryocorbula) dickersoni Weaver & Palmer	D-Te	C-Te
Glyptoactis (Claibornicardia) sandiegoensis (Hanna)	D	C-D
Glyptoactis (Glyptoactis) domenginica (Vokes)	D-Tr	C-Tr
Miltha packi (Dickerson)	D-Te	C-Te
Pitar (Capitaria) uvasanus (Conrad)	Tr-Te	D-Te
Pitar (Lamelliconcha) joaquinensis Vokes	D	C-D
Solena (Eosolen) novacularis (Anderson & Hanna)	D-Te	C-Te
Venericardia (Pacificor) aragonia joaquinensis (Vokes)	D	C-D

M = "Meganos Stage"; C = "Capay Stage"; D = "Domengine Stage"; Tr = "Transition Stage"; Te = "Tejon Stage".

Table 2. Refinements of the West Coast provincial molluscan stage ranges of 11 marine bivalves. These refinements are based on recent research by Squires (1984, 1987, in press).