



FIGURE 2-1-9, *Haimea bajasurensis* n. sp., CSUN loc. 1492. 1, holotype, IGM 5934, aboral view, $\times 3.5$; 2, paratype, IGM 5935, aboral view, $\times 3.2$; 3, paratype, IGM 5936, oral view, $\times 4.5$; 4, 5, paratype, IGM 5935, $\times 2.7$; 4, left side; 5, right side; 6, holotype, IGM 5934, anterior view, $\times 3.5$; 7, paratype, IGM 5936, posterior view, $\times 4.5$; 8, 9, paratype, IGM 5937, $\times 3.5$; 8, lower interior showing lantern supports; 9, upper interior.

extending approximately two-thirds of distance from apical margin to ambitus, other petaloids extending only 60 percent of this distance; straight, open, pore pairs conjugate, interporiferous zone at greatest width at extremity of petaloid, slightly narrower than poriferous zone on specimens with large and ovate genital pores, slightly wider than poriferous zone on spec-

imens with small and circular genital pores; petaloid III with two to three more pore-pairs in poriferous zone than petaloids II and IV; interambulacra two columns in each area, wider than ambulacra; plate sutures smooth; interambulacra at the peristome slightly externally inflated and form bourrelets.

Beyond petals and extending to peristome, ambulacra with