

- to California. U.S. Geological Survey Bulletin 1664A:1-13, pls. 1-4.
- NEWMAN, W. A. 1985. The abyssal hydrothermal vent invertebrate fauna: a glimpse of antiquity? *Biological Society of Washington Bulletin* 6:231-242.
- PROTHERO, D. R. & J. M. ARMENTROUT. 1985. Magnetostratigraphic correlation of the Lincoln Creek Formation, Washington: implications for the age of the Eocene/Oligocene boundary. *Geology* 13:208-211.
- RAU, W. W. 1986. Geologic map of the Humptulips quadrangle and adjacent areas, Grays Harbor County, Washington. Washington Division of Geology and Earth Resources *Geologic Map GM-33*, 1 sheet, scale 1:62,500.
- RISSE, A. 1826. *Histoire naturelle des principales productions de l'Europe Méridionale et particulièrement de celles des environs de Nice et des Alpes Maritimes* 4:1-439, pls. 1-12. Paris.
- SAITO, H. & T. OKUTANI. 1990. Two new chitons (Mollusca: Polyplacophora) from a hydrothermal vent site of the Iheya Small Ridge, Okinawa Trough, east China Sea. *Venus (Japanese Journal of Malacology)* 49(3):165-179, pls. 1-2.
- SIRENKO, B. I. 1988. A new genus of deep sea chitons *Ferrei-raella* gen n. (Lepidopleurida, Leptochitonidae) with a description of new ultra-abyssal species. *Zoologicheskii Zhurnal* 68(12):1776-1786, pls. 1-5 [in Russian].
- SNAVELY, P. D., JR., A. R. NIEM, N. S. MACLEOD, J. E. PEARL & W. W. RAU. 1980. Makah Formation—a deep marginal-basin sedimentary sequence of late Eocene and Oligocene age in the northwestern Olympic Peninsula, Washington. U.S. Geological Survey Professional Paper 1162-B: 1-28.
- SPENCER-JONES, D. 1971. Marginal Tertiary deposits of the Tyrendarra embayment—Grassdale and Hamilton district. Pp. 241-249 in H. Wopfner and J. G. Douglas (eds.), *The Otway Basin of Southeastern Australia*. Geological Surveys of South Australia and Victoria, Special Bulletin.
- SQUIRES, R. L. 1989. Middle Eocene rocky shoreline nearshore mollusks, Tejon Formation, Tehachapi Mountains, California. *Western Society of Malacologists, Annual Report* 21: 18.
- SQUIRES, R. L. 1995. First fossil species of the chemosynthetic-community gastropod *Provanna*: localized cold-seep limestones in upper Eocene and Oligocene rocks, Washington. *The Veliger* 38(1):30-36.
- SQUIRES, R. L. & J. L. GOEDERT. 1991. New late Eocene mollusks from localized deposits formed by subduction-related methane seeps, southwestern Washington. *Journal of Paleontology* 65(3):412-416, figs. 1-2.
- VAN BELLE, R. A. 1975. Sur la présence en Méditerranée de *Lepidopleurus alveolus* (Lovén, 1846). *Informations de la Société Belge de Malacologie, série 4*, 3:57-58, figs. 1-4.
- VAN BELLE, R. A. 1981. *Catalogue of Fossil Chitons (Mollusca: Polyplacophora)*. W. Backhuys: Rotterdam. 84 pp.
- WOLFF, T. 1979. Macrofaunal utilization of plant remains in the deep sea. *Sarsia* 64:117-136.
- WU, S-K. & T. OKUTANI. 1984. The deepsea chitons (Mollusca: Polyplacophora) collected by the R/V Soyo-Maru from Japan—I. *Lepidopleuridae*. *Venus (Japanese Journal of Malacology)* 43(1):1-31, pls. 1-4.