

intertidal zone.

Distribution: Previously recorded only from the type locality, Kyushu, Japan.

Genus *Sphaeroma* Latreille, 1802

Sphaeroma sieboldii Dollfus, 1889

Sphaeroma sieboldii: Richardson, 1909, p. 91; Thielemann, 1910, p. 51; Shiino, 1944, pp. 1-5, figs. 1-3; 1957, pp. 161-167, figs. 1-3; Kim and Kwon, 1983, p. 326.

Material examined: 6 ♂♂, 1 ♀, Pang'o, Anmyōndo I., Apr. 24, 1979, I. H. Kim; 1 ♀, Komso, Apr. 5, 1983, D. H. Kwon; 2 ♂♂, Panwöl, July 2, 1980, W. Kim.

Remarks: This species is active borer of stones and dwells in the hole. Found among oysters or in the hole of stones on mud flats. Shiino (1944, 1957) mentioned the boring habits of this species.

Distribution: Korea and Japan.

References

- Bruce, N. L., 1980. The systematics of some Japanese marine isopods (Fam. Sphaeromatidae) of the genera *Dynoides* Barnard, 1914 and *Cymodocella* Pfeffer, 1887, with descriptions of two new species. *Crustaceana* 38(2): 199~211.
- Choe, B. L. and D. H. Kwon, 1982. The marine invertebrate fauna from Deogjeog Islands. *Report on the Survey of Natural Environment in Korea. No. 1. Deogjeog Archipelago*, pp. 251~266.
- Gurjanova, E., 1933. Contribution to the Isopoda fauna of the Pacific Ocean. I. New species Valvifera and Flabellifera. *Expl. Mers d'U.R.S.S.* 17: 87~106.
- Hatch, H. H., 1947. The Chelifera and Isopoda of Washington and adjacent regions. *Univ. Washington Pub. Biol.* 10(5): 155~274.
- Hoestlandt, H., 1973. Etude systématique et génétique de trois espèces pacifiques Nord-Américaines du genre *Gnorimosphaeroma* (Isopodes Flabellifères). I. Considérations générales et systématique. *Arch. Zool. exp. gen.* 114: 349~395.
- Hoestlandt, H., 1975. Occurrences of the Isopoda Flabellifera *Gnorimosphaeroma rayi* Hoestlandt on the coast of Japan, Eastern Siberia and Hawaii, with a brief note on its genetic polychromatism. *Publ. Seto Mar. Biol. Lab.* 22(1/4): 31~46.
- Iverson, E. W., 1982. Revision of the isopod family Sphaeromatidae (Crustacea: Isopoda: Flabellifera). I. Subfamily names with diagnoses and key. *J. Crust. Biol.* 2(2): 248~254.
- Kim, H. S. and B. L. Choe, 1981. The fauna of marine invertebrate in Ulreung Is. and Dogdo Is... *Rep. KACN* 19: 193~200.
- Kim, H. S. and D. H. Kwon, 1982. Marine invertebrate fauna in the vicinity of Wando Island. *Report on the Survey of Natural Environment in Korea. No. 2. The islands adjacent to Wando*, pp. 187~206.
- Kim, H. S. and D. H. Kwon, 1983. Marine invertebrate fauna in the vicinity of Jindo Island. *Report on the Survey of Natural Environment in Korea. No. 3. Jindo Area, Dadohae Marine National Park*, pp. 313~336.
- Kussakin, O. G., 1974. Fauna and ecology of Isopoda (Crustacea) from the intertidal zone of the Kurile Islands. In: *Flora and fauna of the intertidal zone of the Kurile Islands. Novosibirsk, "Nauka"*, pp. 227~275.
- Kussakin, O. G., 1979. Marine and brackish-water Isopoda of the cold and temperate waters of the Northern Hemisphere. Suborder Flabellifera. *Opredeloteli po Fauna SSSR* 122: 1~470.
- Menzies, R. J., 1954. A review of the systematics and ecology of the genus "*Exosphaeroma*," with the description of a new genus, a new species, and a new subspecies (Crustacea, Isopoda, Sphaeromidae). *Am. Mus. Nov.* 1683: 1~24.
- Nishimura, S., 1968. *Gnorimosphaeroma lata* n. sp., a new marine isopod from Kii, Japan. *Publ. Seto Mar. Biol. Lab.* 16(4): 273~280.
- Nishimura, S., 1976a. Redescription of *Leptosphaeroma gottschei* Hilgendorf (Isopoda, Sphaeromatidae) with some biological notes. *Publ. Seto Mar. Biol. Lab.* 23(1/2): 169~177.

- Nishimura, S., 1976b. *Dynoidella conchicola*, gen. et sp. nov. (Isopoda, Sphaeromatidae), from Japan, with a note on its association with intertidal snails. *Publ. Seto Mar. Biol. Lab.* 23(3/5) 275-282.
- Nunomura, N. and S. Nishimura, 1976. Marine Isopoda from the rocky shore of Osaka Bay, Middle Japan (2). *Bull. Osaka Mus. Nat. Hist.* 30: 19-26.
- Richardson, H., 1904. Contributions to the natural history of the Isopoda. *Proc. U.S. Nat. Mus.* 27: 1-89.
- Richardson, H., 1907. Descriptions of new isopod crustaceans of the family Sphaeromidae. *Proc. U.S. Nat. Mus.* 31: 1-22.
- Richardson, H., 1909. Isopods collected in the Northwest Pacific by the U.S. Bureau of Fisheries steamer "Albatross" in 1906. *Proc. U.S. Nat. Mus.* 37: 75-129.
- Shen, C. J., 1929. Description of a new isopod, *Dynoides dentisinus*, from the coast of North China. *Bull. Fan. Mem. Inst. Biol.* 1: 65-75.
- Shen, C. J., 1933. Redescription of an imperfectly-known species of isopod of the family Sphaeromidae, *Holotelson tuberculatum* Richardson. *Ann. Mag. Nat. Hist., 10 ser.* 12: 269-279.
- Shiino, S. M., 1944. Studies on marine wood-boring crustaceans, II. On four species of boring sphaeromids (Hemibranchiatae). *Misc. Rep. Res. Inst. nat. Resour.* 7: 1-20.
- Shiino, S. M., 1957. The marine wood-boring crustaceans of Japan. II. (Sphaeromidae and Cheluridae). *Wasmann J. Biol.* 15(2): 161-197.
- Tattersall, W. M., 1921. Zoological results of a tour in the Far East. Mysidacea, Tanaidacea and Isopoda. *Mem. Asiat. Soc. Bengal* 6(7): 403-433, pls. 15-16.
- Thielemann, M., 1910. Beiträge zur Kenntnis der Isopodenfauna Ostasiens. Beiträge zur Naturgeschichte Ostasiens herausgegeben von Dr. F. Doflein. *Abh. Bayer. Akad. Wiss., math.-phys. Kl. K., Suppl.* 2(3): 1-109, pls. 1-2.
- Ueno, M., 1933. Freshwater Crustacea of Iturup. *Annot. Zool. Japon.* 14(1): 109-113.
- Ueno, M., 1935. Crustacea collected in the lakes of southern Sakhalin. *Annot. Zool. Japon.* 15(1): 88-94. figs. 1-4.
- Yun, S. G., 1982. Five species of sphaeromatid isopods (Flabellifera: Isopoda) from the southern coast of Korea. *Bull. Nat. Fish. Univ. Busan* 22(2): 1-23.

韓國産 진벌레科 (甲殼上綱, 等脚目, 有屬亞目)의 系統分類學的 研究

金 熏 洙* · 權 道 憲**

韓國産 等脚類의 系統分類學的 研究의 일환으로 1968年 부터 1983年 사이에 南韓의 34개소에 서 채집되어 서울大學校 自然科學大學 動物學科 標本室에 보관되어 있는 진벌레科 等脚類를 同定·分類한 결과 3亞科 6屬 12種이 밝혀졌다. 이 중에는 *Gnorimosphaeroma hoestlandti*, n. sp. 의 1新種과 *Gnorimosphaeroma chinense* (Tattersall)과 *Dynoides brevispina* Bruce의 2韓國未記錄種이 포함되어 있다. 한편 *Cassidias trituberculata* Thielemann, 1910과 *Gnorimosphaeroma kurilense* Kussakin, 1974는 각각 *Holotelson tuberculatus* Richardson, 1909와 *Gnorimosphaeroma chinense* (Tattersall, 1921)의 下位同物異名으로 思料된다. 이로써 韓國産 진벌레科 等脚類는 모두 3亞科 7屬 13種이 된다.

* 서울大學校 自然科學大學 動物學科
** 仁濟大學 生物學科 (動物分類學 전공)